




Contents lists available at SciVerse ScienceDirect

## Inorganic Chemistry Communications

journal homepage: [www.elsevier.com/locate/inoche](http://www.elsevier.com/locate/inoche)

Q41 **Unprecedented Pt(II) complex of an asymmetric 2,6-diacetylpyridine bis**  
 Q52 **(N-substituted thiosemicarbazone) ligand**

Q13 **Ana I. Matesanz, Pilar Souza\***

4 **Departamento de Química Inorgánica (Módulo de Química), Facultad de Ciencias, c/ Francisco Tomás y Valiente nº 7, Universidad Autónoma de Madrid, 28049-Madrid, Spain**

## ARTICLE INFO

## Article history:

Received 10 July 2012

Accepted 13 October 2012

Available online xxxxx

## Keywords:

Asymmetric N<sub>2</sub>S coordination

Bis(thiosemicarbazone)

2,6-diacetylpyridine

Platinum(II) complexes

Thiol function

X-ray diffraction

## ABSTRACT

Reaction of 2,6-diacetylpyridine bis(<sup>4</sup>N-*o*-tolylthiosemicarbazone), **H<sub>2</sub>L<sup>1</sup>**, with K<sub>2</sub>PtCl<sub>4</sub> and further recrystallization in DMSO/MeOH of the [**PL<sup>1</sup>**] complex obtained, led to the isolation of the novel platinum complex, [**PL<sup>2</sup>**], which was structurally characterized by single crystal X-ray diffraction. The molecular structure shows that the ligand has undergone an unexpected chemical transformation viz. reduction of one of the terminal phenyl rings into cyclohexyl. The resulted asymmetrical ligand acts a dianionic tetradentate donor, coordinating to the platinum(II) center in a square planar geometry through the N<sub>pyridinic</sub> atom and the N<sub>imino</sub> and the S atoms from one thiosemicarbazone arm, the fourth coordination position is occupied by the N<sub>hydrazinic</sub> atom of the other arm.

© 2012 Published by Elsevier B.V.

Thiosemicarbazones (R<sub>1</sub>R<sub>2</sub>C=N–NH–C(S)–NR<sub>3</sub>R<sub>4</sub>) are an important and versatile type of ligands due to the potential donor atoms that they possess, among which sulfur is of paramount importance in the metal–ligand linkage. Moreover the π delocalization and configurational flexibility create the possibility of a variety of coordination modes [1–3].

The coordination capacity of thiosemicarbazones can be further increased, if the parent aldehyde or ketone contains additional functional group in a position suitable for chelation. Particularly, compounds in which the thiosemicarbazone side-chain is attached in the α position to an N-heterocyclic ring, namely α-N-heterocyclic thiosemicarbazones, are strong metal chelating agents and have been reported to be among the most effective ribonucleotide reductase (RR) inhibitors yet identified [4]. Pyridine-2-carbaldehyde thiosemicarbazone was the first member of this class reported to have carcinostatic effects and since then many α-N-heterocyclic thiosemicarbazones and their metal complexes have shown anticancer activity against a wide spectrum of tumor cell lines. Currently, the 3-aminopyridine-2-carboxaldehyde thiosemicarbazone (Triapine, Vion Pharmaceuticals, New Haven, CT) is being evaluated in human phase II clinical trials as an antineoplastic therapeutic [5–7].

Although not as intensely studied as the mono(thiosemicarbazones), the α-N-heterocyclic bis(thiosemicarbazones) having the two thiosemicarbazone moieties positioned possess a variety of flexible donor sets and are capable of adopting various coordination modes, leading to enormous structural diversity of their complexes. For example, in the literature a series of dinuclear zinc complexes have been found derived from 2,6-diacetylpyridine bis(thiosemicarbazone) ligands

showing [7 + 7], [6 + 6], [6 + 4] and [4 + 4] coordination environments [8–11]. However no structural information about d<sup>8</sup> metal complexes bearing 2,6-diacetylpyridine bis(thiosemicarbazone) ligands has been encountered in the bibliography.


As a part of a program concerning the behavior of α-N-heterocyclic bis(thiosemicarbazones) we have published structural studies of a series of palladium(II) and platinum(II) complexes derived from 3,5-diacetyl-1,2,4-triazol-bis(thiosemicarbazones) [12–18]. In these complexes, the square-planar coordination geometry of the central metal ions is provided by one N atom of the heterocyclic ring, the N<sub>imino</sub> and S atoms of one thiosemicarbazone arm and the S of the other thiosemicarbazone arm (dimer complexes) or the P atom of the PPh<sub>3</sub> coligand (mononuclear complexes).

To extend the knowledge in this research field, particularly with respect to the coordination properties of the α-N-heterocyclic bis(thiosemicarbazones) and the stereochemistry and molecular structure of the complexes, we undertook the study of platinum(II) complexes derived from 2,6-diacetylpyridine bis(α-N-substituted thiosemicarbazones). Here we report on the synthesis, characterization and crystal structure of a new Pt(II) complex derived from an asymmetric 2,6-diacetylpyridine bis(α-N-substituted thiosemicarbazone) ligand.

The novel ligand 2,6-diacetylpyridine bis(<sup>4</sup>N-*o*-tolylthiosemicarbazone), **H<sub>2</sub>L<sup>1</sup>**, was synthesized by refluxing an ethanolic solution (20 mL) of 2,6-diacetylpyridine (1 mmol) with <sup>4</sup>N-*o*-tolylthiosemicarbazide (2 mmol), which was prepared as described in reference [18], for 5 h and then was left to stand in ambient temperature. The solution was reduced to half volume and the pale yellow solid formed was filtered, washed with cold EtOH and Et<sub>2</sub>O and dried in vacuo. The ligand was characterized by elemental analysis and FAB spectrometry as well as by IR and <sup>1</sup>H NMR spectroscopy [19].

\* Corresponding author. Tel.: +34 914975146; fax: +34 914974833.

E-mail address: [pilar.souza@uam.es](mailto:pilar.souza@uam.es) (P. Souza).



Scheme 1

Reaction of methanolic suspension (20 mL) of  $\text{H}_2\text{L}^1$  ligand (1.0 mmol) with  $\text{K}_2\text{PtCl}_4$  (1.0 mmol) in water for 5 h at room temperature yielded a brown precipitate which was filtered, washed with MeOH and  $\text{Et}_2\text{O}$ , purified by crystallization from DMSO and dried in vacuo. Analytical and spectroscopic characterization [20] was consistent with the formation of the expected neutral  $[\text{PtL}^1]$  complex (Scheme 1).

Further recrystallization of  $[\text{PtL}^1]$  from DMSO/MeOH led to the isolation of good quality single crystals which were studied by X-ray diffraction techniques [21]. The structural analysis allowed us to identify a new platinum(II) complex,  $[\text{PtL}^2]$ , in which the 2,6-diacetylpyridine bis( $^4\text{N}$ -*o*-tolylthiosemicarbazone) ligand has undergone an unexpected chemical transformation viz. reduction of one of the terminal phenyl rings into cyclohexyl.

Conventional hydrogenation reactions have implicit use of hydrogen as a reactant. However it is possible that in the presence of a metal complex, a donor hydrogen molecule transfers to a substrate which acts as an acceptor. The donor molecules which undergo dehydrogenation are often the reaction solvents and the catalysts are usually derived from platinum group metals with nitrogen or phosphorus donating ligands with the Wilkinson's catalyst  $[\text{RhCl}(\text{PPh}_3)_3]$  being the most prominent [22–24].

By inspection of the literature a few examples of thiosemicarbazone metal complexes exhibiting catalytic activity have been found. The fundamental features of these catalysts are the presence, in the complexes, of stabilizing five membered chelate rings as well as the presence of a labile coordinating bond prone to dissociate to provide an available coordination site [25–27].

In our case, the platinum complex  $[\text{PtL}^1]$  contains two five membered chelate rings as well as a more rigid and therefore less stable six membered chelate ring. Although further studies are necessary in order to identify the reductant, a possible candidate could be MeOH employed as solvent.

Since we have noticed that platinum(II) complexes derived of 3,5-diacetyl-1,2,4-triazol bis( $^4\text{N}$ -*o*-tolylthiosemicarbazone) ligand have shown a notable antitumor activity [17] we analyzed the cytotoxic properties of the new complex  $[\text{PtL}^2]$  by testing its antiproliferative activity in vitro against five human cancer cell lines: NCI-H460 (non-small cell lung cancer), HepG2 (hepatocellular carcinoma), MCF-7 (breast cancer), A2780 and A2780cisR (epithelial ovarian cancer) which are among the lines used in the NCI to identify novel potential anticancer drugs. Surprisingly, the platinum(II) complex  $[\text{PtL}^2]$  shows at 100  $\mu\text{M}$  concentration, a very low cellular growth inhibition (<50%) and therefore did not have evaluable cytotoxicity ( $\text{IC}_{50} > 100 \mu\text{M}$ ). The substitution of the N-heterocyclic ring (1,2,4-triazol versus pyridine) as well as the hydrogenation on the peripheral tolyl substituent seems to be factors that influence both structure and cytotoxicity.

The molecular structure of the neutral complex  $[\text{PtL}^2]$ , which crystallized with one DMSO molecule in the monoclinic  $P2_1$  space group, together with the atom labeling scheme is shown in Fig. 1.

The transformed asymmetrical ligand acts a dianionic tetradentate N,N,N,S-donor, coordinating to the platinum(II) center in a square


planar geometry through the  $\text{N}_{\text{pyridinic}}$  atom and the  $\text{N}_{\text{iminic}}$  and the S atoms from one thiosemicarbazone arm. The fourth coordination position is occupied by the  $\text{N}_{\text{hydrazinic}}$  atom of the other arm generating two typical five membered ( $\text{PtSCNN}$  and  $\text{PtNCCN}$ ) and one six membered ( $\text{PtNNCCN}$ ) chelate rings. Coordination by the  $\text{N}_{\text{hydrazinic}}$  instead of the  $\text{N}_{\text{iminic}}$  atom, although uncommon, has been found in the bibliography for some palladium and nickel bis(thiosemicarbazone) complexes [28–30].


The Pt–N [1.979(7), 2.021(8) and 2.023(8) Å] and Pt–S [2.289(2) Å] bond distances are comparable with those reported for Pt(II) thiosemicarbazone complexes.

Since the two thiosemicarbazone moieties coordinate in a different fashion it would expect that in the bidentate- $\text{N}^{\wedge}\text{S}$  arm, the C–S distance undergoes significant evolution from the thione to the thiol form [C–S distance of 1.752(9) Å] but it is important to note that the monodentate- $\text{N}_{\text{hydrazinic}}$  thiosemicarbazone arm also presents thiol C–S bond [1.81(3) Å] as well as an unexpected S–H bond.

It is well known that compounds containing thiosemicarbazone functional groups exhibit thiol–thione tautomerism, but unsubstituted and monosubstituted ones,  $>\text{C}=\text{N}-\text{NH}-\text{C}(\text{S})-\text{NHR}$ , are capable of stabilizing a third thiol form (Scheme 2). This is consistent with the C(15)–N(5) and C(17)–N(7) bond lengths [1.309(17) and 1.295(17) Å respectively] which correspond formally to double bonds while the C(17)–N(6) bond length is longer, 1.408(13) Å.

On the other hand, along the bidentate thiosemicarbazone arm and as consequence of the extensive delocalization of electron density, the

Fig. 1. Molecular structure of platinum(II) complex  $[\text{PtL}^2]$ .



Scheme 2

Q3

170 C–N and N–N bond distances are intermediate between formal single  
171 and double bonds.

172 Inspection of the angles formed between the platinum(II) ion and  
173 the coordinated atoms shows that the metal is contained within a  
174 slightly distorted square-planar environment. The distortion is caused  
175 by the restricted bite angle of the N(4), N(3), and S(1) donor set as  
176 reflected in the S(1)–Pt(1)–N(3) and N(3)–Pt(1)–N(4) angles (less  
177 than 90°). The angles N(4)–Pt(1)–N(6) and N(6)–Pt(1)–S(1) are  
178 therefore greater than 90°.

179 The crystal structure is stabilized by intermolecular hydrogen in-  
180 teraction involving the N(1) atom of the bidentate thiosemicarbazone  
181 arm and the oxygen atom of the DMSO solvent molecule. Within each  
182 molecule, the bis(thiosemicarbazone)-platinum moiety is close to  
183 planar, so the supramolecular association also involves  $\pi$ - $\pi$  stacking  
184 interactions between parallel layers of molecules (Fig. 2).

### 185 Acknowledgments

186 We are grateful to Ministerio de Economía y Competitividad,  
187 Instituto de Salud Carlos III of Spain (PI080525 and PI1100659) for  
188 financial support.

### 189 Appendix A. Supplementary material

190 Full crystallographic details have been deposited in CIF format  
191 with the Cambridge Crystallographic Data Centre. CCDC 890162 con-  
192 tains the supplementary crystallographic data for this paper. These  
193 data can be obtained free of charge via [http://www.ccdc.cam.ac.uk/  
194 conts/retrieving.html](http://www.ccdc.cam.ac.uk/conts/retrieving.html), or from the Cambridge Crystallographic Data  
195 Centre, 12 Union Road, Cambridge CB21EZ, UK; fax: +44 1223 336  
196 408; e-mail: [deposit@ccdc.cam.ac.uk](mailto:deposit@ccdc.cam.ac.uk). Supplementary data to this article  
197 can be found online at [http://dx.doi.org/10.1016/j.inoche.2012.10.  
198 022](http://dx.doi.org/10.1016/j.inoche.2012.10.022).




Fig. 2. View of platinum(II) complex [PtL<sup>2</sup>] showing the parallel disposition of the molecules as a result of  $\pi$ - $\pi$  stacking interactions.

### References

- 199
- [1] T.S. Lobana, R. Sharma, G. Bawa, S. Khanna, Bonding and structure trends of thiosemicarbazone derivatives of metals—an overview, *Coord. Chem. Rev.* 253 (2009) 977–1055.
  - [2] J.M. Vila, T. Pereira, A. Amoedo, M. Graña, J. Martínez, M. López-Torres, A. Fernández, The key role of sulfur in thiosemicarbazone compounds. Crystal and molecular structure of [Pd{4-MeOC<sub>6</sub>H<sub>4</sub>C(Me)=NN=C(S)NHPPh<sub>2</sub>}]<sub>2</sub>, *J. Organomet. Chem.* 623 (2001) 176–184.
  - [3] J.S. Casas, M.S. García-Tasende, J. Sordo, Main group metal complexes of semicarbazones and thiosemicarbazones. A structural review, *Coord. Chem. Rev.* 209 (2000) 197–261.
  - [4] A.C. Sartorelli, K.C. Agrawal, A.S. Tsiftoglou, E.C. Moore, Characterization of the biochemical mechanism of action of  $\alpha$ -(N)-heterocyclic carboxaldehyde thiosemicarbazone, *Adv. Enzym. Regul.* (1977) 117–139.
  - [5] M.C. Liu, T.S. Lin, A.C. Sartorelli, Synthesis and antitumor-activity of amino derivatives of pyridine-2-carboxaldehyde thiosemicarbazone, *J. Med. Chem.* 35 (1992) 3672–3677.
  - [6] R.A. Finch, M.C. Liu, A.H. Cory, J.G. Cory, A.C. Sartorelli, Triapine (3-aminopyridine-2-carboxaldehyde thiosemicarbazone; 3-AP): an inhibitor of ribonucleotide reductase with antineoplastic activity, *Adv. Enzym. Regul.* 39 (1999) 3–12.
  - [7] R.A. Finch, M.C. Liu, S.P. Grill, S.P.W.C. Rose, R. Loomis, K.M. Vasquez, Y.C. Cheng, A.C. Sartorelli, Triapine (3-aminopyridine-2-carboxaldehydethiosemicarbazone): a potent inhibitor of ribonucleotide reductase activity with broad spectrum antitumor activity, *Biochem. Pharmacol.* 59 (2000) 983–991.
  - [8] A.I. Matesanz, I. Cuadrado, C. Pastor, P. Souza, A novel sulfur-bridged dimeric zinc(II) complex with 2,6-diacetylpyridine bis(thiosemicarbazone), *Z. Anorg. Allg. Chem.* 631 (2005) 780–784.
  - [9] A. Bino, N. Cohen, Several coordination modes of the pentadentate ligand 2,6-diacetylpyridinebis(thiosemicarbazone), *Inorg. Chim. Acta* 210 (1993) 11.
  - [10] M.R. Bermejo, A.M. González-Noya, R.M. Pedrido, M.J. Romero, M. Vázquez, Route to cluster helicates, *Angew. Chem.* 117 (2005) 4254–4259.
  - [11] M.R. Bermejo, A.M. González-Noya, M. Martínez-Calvo, R. Pedrido, M.J. Romero, M. Vázquez, Checking the route to cluster helicates, *Eur. J. Inorg. Chem.* (2008) 3852–3863.
  - [12] P. Souza, A.I. Matesanz, C. Pastor, Preparation and structural characterisation of a novel palladium(II) binuclear complex containing triazole bithiosemicarbazone bridges, *Inorg. Chem. Commun.* 5 (2002) 344–346.
  - [13] A.I. Matesanz, J. Mosa, I. García, C. Pastor, P. Souza, Synthesis, characterization, crystal structure and electrochemistry of a novel palladium(II) binuclear complex containing 1,2,4-triazole bis(4-phenylthiosemicarbazone) bridges, *Inorg. Chem. Commun.* 7 (2004) 756–759.
  - [14] A.I. Matesanz, C. Pastor, P. Souza, Synthesis and structural characterization of a disulphide-bridged tetranuclear palladium(II) complex derived from 3,5-diacetyl-1,2,4-triazole bis(4-ethylthiosemicarbazone), *Inorg. Chem. Commun.* 10 (2007) 97–100.
  - [15] Ana I. Matesanz, Pilar Souza, Palladium and platinum 3,5-diacetyl-1,2,4-triazol bis(thiosemicarbazones): chemistry, cytotoxic activity and structure-activity relationships, *J. Inorg. Biochem.* 101 (2007) 245–253.
  - [16] A.I. Matesanz, C. Joie, P. Souza, Chemistry, antiproliferative activity and low nephrotoxicity of 3,5-diacetyl-1,2,4-triazol bis(<sup>4</sup>N-thiosemicarbazone) ligands and their platinum(II) complexes, *Dalton Trans.* 39 (2010) 7059–7065.
  - [17] A.I. Matesanz, C. Hernández, A. Rodríguez, P. Souza, Novel bis(thiosemicarbazones) of the 3,5-diacetyl-1,2,4-triazol series and their platinum(II) complexes: chemistry, antiproliferative activity and preliminary nephrotoxicity studies, *Dalton Trans.* 40 (2011) 5738–5745.
  - [18] A.I. Matesanz, C. Hernández, A. Rodríguez, P. Souza, 3,5-Diacetyl-1,2,4-triazol bis(<sup>4</sup>N-substituted thiosemicarbazone) palladium(II) complexes: synthesis, structure, antiproliferative activity and low toxicity on normal kidney cells, *J. Inorg. Biochem.* 105 (2011) 1613–1622.
  - [19] Ligand **H<sub>2</sub>L<sup>1</sup>**: Yield (73%). *Anal.* % found: C, 61.00; H, 5.55; N, 19.85; S, 12.90. C<sub>25</sub>H<sub>27</sub>N<sub>5</sub>S<sub>2</sub> requires: C, 61.35; H, 5.50; N, 20.05; S, 13.10. MS(FAB<sup>+</sup> with mNBA matrix), m/z = 490 for [H<sub>2</sub>L<sup>1</sup> + H]<sup>+</sup>. IR (KBr, cm<sup>-1</sup>),  $\nu$  = 3308, 3238, 3151 (s, NH);  $\nu$  = 1589 (s, CN),  $\nu$  = 1520 (s, CN-thioamide I),  $\nu$  = 878 (m, CS-thioamide IV). <sup>1</sup>H NMR (300 MHz, d<sup>6</sup>-DMSO, ppm),  $\delta$  = 10.65 [s, N(2) and N(6), 2H]; 10.10 [s, N(1) and N(7), 2H];  $\delta$  = 8.55 [d, C(11) and C(13), 2H]; 7.75 [t, C(12), 1H];  $\delta$  = 7.30–7.20 (m, aromatic-thiosemicarbazide, 8H);  $\delta$  = 2.50 (s, CH<sub>3</sub>-thiosemicarbazide, 6H);  $\delta$  = 2.25 (s, CH<sub>3</sub>-diacetylpyridine, 6H).
  - [20] Complex [PtL<sup>1</sup>]: Yield (55%). *Anal.* % found: C, 42.35; H, 3.95; N, 13.00; S 12.25; C<sub>25</sub>H<sub>31</sub>N<sub>7</sub>PtS<sub>2</sub>-DMSO requires C, 42.45; H, 4.30; N, 12.85; S 12.60%. MS(FAB<sup>+</sup> with mNBA matrix), m/z = 683 for [PtL<sup>1</sup> + H]<sup>+</sup>. IR (KBr, cm<sup>-1</sup>),  $\nu$  = 3353 (s, NH),  $\nu$  = 1586 (s, CN),

- 268  $\nu=1523$  (s, CN-thioamide I),  $\nu=849$  (w, CS-thioamide IV).  $^1\text{H}$  NMR (300 MHz,  $\text{d}^6\text{-DMSO}$ , ppm),  $\delta=10.77, 9.50$  [s, N(1) and N(7), 1H];  $\delta=8.52$  [d, C(11) and C(13), 2H];  $7.98$  [t, C(12), 1H];  $\delta=7.37\text{--}7.10$  (m, aromatic-thiosemicarbazide, 8H);  $\delta=2.76$  (s,  $\text{CH}_3$ -thiosemicarbazide, 6H);  $\delta=2.27, 2.21$  (s,  $\text{CH}_3$ -diacetylpyridine, 3H).
- 270  
271  
272 [21] Crystal data for complex  $[\text{PtL}^2]\cdot\text{DMSO}$ :  $\text{C}_{27}\text{H}_{37}\text{N}_7\text{O}_4\text{PtS}_3$ ,  $M=766.91$ , monoclinic, space group  $P2_1$ ,  $a=13.7849(6)$  Å,  $b=7.0373(3)$  Å,  $c=15.8575(6)$  Å,  $\alpha=90^\circ$ ,  $\beta=113.933(2)^\circ$ ,  $\gamma=90^\circ$ ,  $V=1406.05(10)$  Å<sup>3</sup>,  $T=296(2)$  K,  $Z=2$ ,  $D_c=1.811$  Mg/m<sup>3</sup>,  $F(000)=764$ ,  $\mu=5.249$  mm<sup>-1</sup>,  $\lambda=0.71073$  Å, 16,080 observed reflections, 5150 independent reflections [ $R(\text{int})=0.0495$ ]. The final agreement factors are  $R1=0.0411$ ,  $wR2=0.1076$  with  $I>2\sigma(I)$  and  $R$  indices (all data)  $R1=0.0571$ ,  $wR2=0.1382$ .
- 273  
274  
275  
276  
277 [22] K. Raghuraman, N. Pillarsetty, K.R. Prabhu, K.K. Katti, K.V. Katti, Unprecedented rhodium-mediated catalytic transfer hydrogenation of a phosphonate functionalized olefin in ecofriendly media, *Inorg. Chim. Acta* 357 (2004) 2933–2938.
- 278  
279  
280 [23] M. Albrecht, R.H. Crabtree, J. Mata, E. Peris, Chelatin bis-carbene rhodium(III) complexes in transfer hydrogenation of ketones and imines, *Chem. Commun.* (2002) 32–33.
- 281  
282  
283 [24] P.J. Dyson, Arene hydrogenation by homogeneous catalysts: fact or fiction? *Dalton Trans.* (2003) 2964–2974.
- 284  
285  
286 [25] I. Pal, S. Dutta, F. Basuli, S. Goverdhan, S. Peng, G.H. Lee, S. Bhattacharya, Unprecedented chemical transformation of semicarbazones mediated by Wilkinson's catalyst, *Inorg. Chem.* 42 (2003) 4338–4345.
- 287  
288
- [26] P. Pelagatti, A. Venturini, A. Leporati, M. Carcelli, M. Costa, A. Bacchi, G. Pelizzi, C. Pelizzi, Chemoselective homogeneous hydrogenation of phenylacetylene using thiosemicarbazone and thiobenzoylhydrazone palladium(II) complexes as catalysts, *Dalton Trans.* (1998) 2715–2721. 289–292
- [27] A. Bacchi, M. Carcelli, M. Costa, A. Leporati, E. Leporati, P. Pelagatti, C. Pelizzi, G. Pelizzi, Palladium(II) complexes containing a P, N chelating ligand part II. Synthesis and characterisation of complexes with different hydrazinic ligands. Catalytic activity in the hydrogenation of double and triple C–C bonds, *J. Organomet. Chem.* 535 (1997) 107–120. 293–297
- [28] C.A. Brown, W. Kaminsky, K.A. Claborn, K.I. Goldberg, D.X. West, Structural studies of 2,6-diacetyl- and 2,6-diformylpyridine bis(thiosemicarbazones), *J. Braz. Chem. Soc.* 13 (2002) 10–18. 298–300
- [29] J.I. Gradinaru, S.T. Malinowski, M.A. Popovici, M. Gdaniec, Template synthesis and crystal structure of an asymmetric square-planar complex: pyridine-2,6-dicarbonyl bis(S-methylisothiosemicarbazonato) nickel(II) iodide, *Crystallogr. Rep.* 50 (2005) 217–223. 301–304
- [30] T.R. Todorović, A. Bacchi, G. Pelizzi, N.O. Juranić, D.M. Sladić, I.D. Brčeski, K.K. Anelković, Synthesis and characterization of Zn(II) and Cd(II) complexes with 2,6-diacetylpyridine-bis(selenosemicarbazone). Crystal structure of a Ni(II) complex with a modified 2,6-diacetylpyridine-bis(selenosemicarbazone), *Inorg. Chem. Commun.* 9 (2006) 862–865. 305–309