

A search for pre-substellar cores and proto-brown dwarf candidates in Taurus: multiwavelength analysis in the B213-L1495 clouds

Aina Palau,^{1*} I. de Gregorio-Monsalvo,^{2,3} Ò. Morata,⁴ D. Stamatellos,⁵ N. Huélamo,⁶ C. Eiroa,⁷ A. Bayo,² M. Morales-Calderón,⁸ H. Bouy,⁶ Á. Ribas,⁹ D. Asmus¹⁰ and D. Barrado^{6,11}

¹*Institut de Ciències de l'Espai (CSIC/IEEC), Campus UAB, Facultat de Ciències, Torre C-5 parell 2, E-08193 Bellaterra, Spain*

²*European Southern Observatory, Alonso de Córdova 3107, Vitacura, Santiago, Chile*

³*Joint ALMA Office, Alonso de Córdova 3107, Vitacura, Santiago, Chile*

⁴*Academia Sinica, Institute of Astronomy and Astrophysics, PO Box 23-141, Taipei 106, Taiwan*

⁵*School of Physics & Astronomy, Cardiff University, 5 The Parade, Cardiff CF24 3AA, UK*

⁶*Dpto. Astrofísica, Centro de Astrobiología (INTA-CSIC), ESAC Campus, PO Box 78, E-28691 Villanueva de la Cañada, Spain*

⁷*Dpto. Física Teórica, Facultad de Ciencias, Universidad Autónoma de Madrid, E-28049 Madrid, Spain*

⁸*Spitzer Science Center, California Institute of Technology, 1200 E California Blvd., Pasadena, CA 91125, USA*

⁹*Herschel Science Centre, European Space Astronomy Centre (ESA), PO Box 78, E-28691 Villanueva de la Cañada, Madrid, Spain*

¹⁰*Institut für Theoretische Physik und Astrophysik, Christian-Albrechts-Universität zu Kiel, Leibnizstr. 15, 24098 Kiel, Germany*

¹¹*Centro Astronómico Hispano Alemán de Calar Alto (CAHA), C/ Jesús Durbán Remón 2-2, E-04004 Almería, Spain*

Accepted 2012 May 24. Received 2012 May 21; in original form 2012 March 20

ABSTRACT

In an attempt to study whether the formation of brown dwarfs (BDs) takes place as a scaled-down version of low-mass stars, we conducted Institut de Radioastronomie Millimétrique (IRAM) 30 m/MAMBO-II observations at 1.2 mm in a sample of 12 proto-BD candidates selected from *Spitzer*/IRAC data in the B213-L1495 clouds in Taurus. Subsequent observations with the Caltech Submillimetre Observatory at 350 μm , Very Large Array at 3.6 and 6 cm, and IRAM 30 m/EMIR in the ^{12}CO (1–0), ^{13}CO (1–0) and N_2H^+ (1–0) transitions were carried out towards the two most promising *Spitzer*/IRAC source(s), J042118 and J041757. J042118 is associated with a compact (<10 arcsec or <1400 au) and faint source at 350 μm , while J041757 is associated with a partially resolved (~ 16 arcsec or ~ 2000 au) and stronger source emitting at centimetre wavelengths with a flat spectral index. The corresponding masses of the dust condensations are ~ 1 and $5 M_{\text{Jup}}$ for J042118 and J041757, respectively. In addition, about 40 arcsec to the north-east of J041757, we detect a strong and extended submillimetre source, J041757-NE, which is not associated with near-infrared/far-infrared emission down to our detection limits, but is clearly detected in ^{13}CO and N_2H^+ at $\sim 7 \text{ km s}^{-1}$, and for which we estimated a total mass of $\sim 100 M_{\text{Jup}}$, close to the mass required to be gravitationally bound. In summary, our observational strategy has allowed us to find in B213-L1495 two proto-BD candidates and one pre-substellar core candidate, whose properties seem to be consistent with a scaled-down version of low-mass stars.

Key words: brown dwarfs – stars: formation – stars: individual: SSTB213 J041757 – stars: individual: SSTB213 J042118 – ISM: clouds – dust, extinction.

1 INTRODUCTION

The formation mechanism of brown dwarfs (BDs) is currently a matter of vigorous debate. Since the typical thermal Jeans mass in molecular cloud cores is around $1 M_{\odot}$, a cloud should not fragment in cores of substellar masses and the formation of BDs cannot be

directly explained as a scaled-down version of low-mass star formation. A possible scenario is that BDs form from fragmentation of turbulent cores. In this scenario, turbulence produces particular density enhancements, decreasing the Jeans mass and allowing the formation of cloud cores of very low masses (Padoan & Nordlund 2004). Another possible scenario is the formation of BDs through fragmentation of massive discs and subsequent ejection (Rice et al. 2003; Stamatellos & Whitworth 2009), or ejection could also take place in the very first stages of formation of a multiple system (e.g.

*E-mail: palau@ieec.uab.es

Reipurth & Clarke 2001; Umbreit et al. 2005; Basu & Vorobyov 2012), depriving the ejected objects from gaining further mass. Other scenarios propose that an initially typical core is photo-eroded due to the nearby presence of a massive star (Whitworth & Zinnecker 2004).

An observational way to distinguish among these different scenarios is the search and characterization of BDs in their very first evolutionary stages, what we call here the proto-BD stage [following the nomenclature of Pound & Blitz (1993, 1995), and equivalent to Class 0/I stages of low-mass star formation]. If BDs form in a similar way as their low-mass stellar counterparts, one would expect to find proto-BDs surrounded by substantial discs and envelopes, with the envelopes similar to the ones observed in the first stages of low-mass protostars (e.g. André, Ward-Thompson & Barsony 1993), which are strongly emitting in the millimetre/submillimetre range. In addition, proto-BDs should be found associated with outflow and ejection phenomena, whose parameters should be a scaled-down version of low-mass stars, and perhaps should even form planetary systems. As a matter of fact, a significant number of BDs with ages of a few Myr have been shown to have discs (e.g. Luhman, Liebert & Rieke 1997; White et al. 1999; Fernández & Comerón 2001; Barrado y Navascués & Martín 2003; Pascucci et al. 2003; Barrado y Navascués, Mohanty & Jayawardhana 2004; Natta et al. 2004; Apai et al. 2005; Luhman et al. 2005, 2007, 2008, 2010; Mohanty, Jayawardhana & Basri 2005; Jayawardhana & Ivanov 2006; Jayawardhana et al. 2006; Barrado y Navascués et al. 2007; Harvey et al. 2010; Monin et al. 2010; Joergens et al. 2012; Riaz et al. 2012b), but most of these discs were inferred from optical or near-infrared/mid-infrared (NIR/MIR) data, thus being T-Tauri analogues or Class II sources.

Different research groups have searched for proto-BDs in the millimetre/submillimetre range. One of the pioneering works was that of Pound & Blitz (1993, 1995), who find no clear-cut evidence of proto-BDs. These works were followed up by Greaves, Holland & Pound (2003), who survey an area of the Ophiucus star-forming region with the James Clerk Maxwell Telescope (JCMT) at 850 μm and find a dozen of possible planetary-mass isolated objects. However, their submillimetre sources could be tracing either truly proto-BDs or transient objects, and these objects remain to be confirmed through molecular line observations. Other candidates to be BDs in the Class 0/I phase are the so-called very low luminosity objects (VeLLOs), objects embedded in dense cores with internal luminosities ($<0.1 L_{\odot}$; Di Francesco et al. 2007; Dunham et al. 2008), such as L1014-IRS and L1148-IRS, which are associated with dusty and dense envelopes, centimetre emission (L1014: Shirley et al. 2007), and CO (1–0) and CO (2–1) outflows (Bourke et al. 2005; Kauffmann et al. 2011), all properties suggestive of accretion/ejection of matter. However, for L1014-IRS and the other few VeLLOs studied in detail (e.g. André, Motte & Bacmann 1999; Young et al. 2004; Kauffmann et al. 2005, 2008, Lee et al. 2009; Dunham et al. 2010; Kauffmann et al. 2011), their true *substellar* nature is not clear (e.g. Lee 2007), neither from the bolometric luminosity ($\gtrsim 0.05 L_{\odot}$) nor from the mass of the envelope, which could increase the present mass well above the substellar limit once accreted on to the object (Kauffmann et al. 2011). On the other hand, the studies of Klein et al. (2003) and Scholz, Jayawardhana & Wood (2006), who carry out surveys with the JCMT and/or the Institut de Radioastronomie Millimétrique (IRAM) 30 m telescope, were addressed towards samples of relatively old BDs (>1 Myr), selected from optical/infrared (IR) surveys, while the most embedded objects are expected at ages $\lesssim 0.5$ Myr. Finally, individual proto-BD candidates studied in detail in the millimetre/submillimetre range

either are more evolved than Class I sources (e.g. Phan-Bao et al. 2008; Harvey et al. 2012; Riaz et al. 2012a) or do not have the mass well constrained and could still be stellar (e.g. Scholz et al. 2008).

In this paper, we show the results of a millimetre/submillimetre study of a sample of 12 proto-BD candidates selected in the B213-L1495 dark clouds of the Taurus–Auriga complex using *Spitzer* data. We show the results of the observations at 1.2 mm towards this sample, as well as submillimetre data for the two most promising candidates found at 1.2 mm. One of the candidates was not previously reported, while the other candidate was already reported in Barrado et al. (2009), and here we present subsequent observations aimed at further constraining its nature, which additionally reveal a nearby starless core of substellar mass, i.e. a possible pre-substellar core.

2 SELECTION STRATEGY AND OBSERVATIONS

2.1 Selection strategy

We used IRAC and MIPS (*Spitzer*) data towards one of the youngest regions in Taurus, the B213-L1495 region. This region was selected from the Froebrich (2005) list of Class 0 sources, because it was one of the nearby (<300 pc) regions with a higher ratio of Class 0/I to Class II IRAC sources (see fig. 1 in Barrado et al. 2009). For B213-L1495, we downloaded the *Spitzer* archive data and selected 12 IRAC sources which were classified as Class 0/I according to IRAC colour–colour diagrams (Allen et al. 2004) and substellar (with a magnitude below the $0.075 M_{\odot}$ cut in the colour–magnitude diagram, after comparing with evolutionary models; Barrado et al. 2009). In addition, we rejected those objects classified as extragalactic candidates following the criteria of Harvey et al. (2006), Jørgensen et al. (2006) and Bouy et al. (2009). Fig. 1 shows the final sample overlaid on an extinction map of the region, elaborated for this work using the star count method on 2MASS data (Skrutskie et al. 2006).

2.2 IRAM 30 m ON–OFF at 1.2 mm and APEX at 870 μm for the sample of 12 proto-BD candidates

We observed with the IRAM 30 m¹ single-dish telescope (Pico Veleta, Spain) the 1.2 mm continuum emission of the 12 selected BD candidates using the 117-channel MAMBO-II bolometer array. The half-power beam width (HPBW) of the telescope at 1.2 mm is 11 arcsec. Observations were carried out in 2007 May, October, November and 2008 November 14, and the opacities ranged between 0.30 and 0.40. Each source was centred in the most sensitive pixel of the array (pixel 20). Typical pointing corrections were $\lesssim 5$ arcsec, providing a positional accuracy of ~ 3 arcsec. The background subtraction was performed by using the ON–OFF observing mode, with a wobbler throw of 32 arcsec. Every source was observed for at least one scan of 20 min with an integration time per subscan of 60 s. The data were reduced using the MOPSIC (Zylka 1998) pipeline provided by IRAM. In order to test the consistency of the data, we measured for each source the flux density for each scan separately and obtained values that were consistent

¹The IRAM 30 m telescope is operated by the Institut de Radioastronomie Millimétrique, which is supported by INSU/CNRS (France), MPG (Germany) and IGN (Spain).




Figure 1. Extinction map in 2MASS J -band derived with an adaptive star count method (Cambresy et al. 1997) in the B213-L1495 region (45 arcmin radius). Members and candidate members from Kenyon & Hartmann (1995) and references therein have been taken into account (removed from the star counting) when computing the A_J values. The resolution of the map varies from 30 to 50 arcsec, depending on the density of stars detected by 2MASS. Plus signs mark the positions of the selected *Spitzer*/IRAC sources observed with the IRAM 30 m telescope and the stars correspond to the sources additionally observed with the CSO.

within the uncertainties. In Table 1, we summarize the observations presented in this work, and in Table 2 we give the list of the 12 selected *Spitzer*/IRAC sources,² total integration time, opacity at 250 GHz and the measured flux density at 1.2 mm. The final rms noise at 1.2 mm was typically 0.7–1.5 mJy beam⁻¹, depending on the integration time. None of the 12 *Spitzer*/IRAC sources was detected above the 4σ level, corresponding to envelope/disc masses $\lesssim 3 M_{\text{Jup}}$ (for a dust temperature of 15 K and a dust opacity at 1.2 mm of 0.009 cm² g⁻¹; Ossenkopf & Henning 1994, table 2). However, for two of the observed *Spitzer*/IRAC sources, we measured fluxes above 2σ , namely J041757 and J042118, which were subsequently observed with the Caltech Submillimetre Observatory (CSO).

In addition, we used public archived data from the Large Bolometer Camera (LABOCA) for the Atacama Pathfinder Experiment (APEX) antenna.³ Observations at 870 μm were carried out in 2009 July under the project E-083.C-0453A.2009 (P.I.: Tafalla). The region mapped corresponds to Taurus B213 and B211 filaments, and was observed during a total time of 14 h, under good weather conditions [precipitable water vapour (PWV) ranged between 0.25 and 1.8 mm]. Data reduction was performed using BOA and MINICRUSH software packages following standard procedures. The LABOCA image covers five of the 12 *Spitzer*/IRAC sources of our sample, and none of the five sources were detected (see Table 2 for upper limits).

² Throughout this paper, we will refer to each source of Table 2 designated by ‘SSTB213 J04mmss.ss+ddmmss.s’ as ‘J04mmss’.

³ APEX is a collaboration between the Max-Planck-Institut fuer Radioastronomie, the European Southern Observatory and the Onsala Space Observatory.

2.3 CSO at 350 μm for J042118 and J041757

The CSO observations towards J042118 and J041757 were carried out on 2008 January 28, using the Submillimeter High Angular Resolution Camera (SHARC-II) bolometer array at 350 μm . The HPBW of the CSO at this wavelength is 8.6 arcsec, which was convolved to a final beam of 10.4 arcsec. Typical opacities at 225 GHz were around 0.04. The observations were carried out in the imaging mode using a Lissajous pattern of 20 arcsec of amplitude and 14.142 s of period, resulting in a final image size of 3.3×3.2 arcmin². In this observing mode, the telescope modulates the X and Y positions with a different sine wave for each axis (starting from the centre of the map). As a consequence, the rms is lower in the centre of the image than at the edges. Each scan was integrated for 10 min, and the total observing time per source was around 1–1.5 h. The absolute flux uncertainty was estimated from a measurement of the flux of HL Tau, and was $\lesssim 10$ per cent. The uncertainty in position was ~ 3 arcsec. The reduction of the SHARC-II data was carried out with the CRUSH software (Kovács 2008), version 1.62-4. For both sources, we assumed that the emission is compact (< 30 arcsec) and faint (< 1 Jy).⁴ The rms noises achieved at the map centres are 9 mJy for J042118 and 6 mJy for J041757.

2.4 VLA observations at 6 and 3.6 cm for J041757

We obtained radio continuum data at 6 (4.86 GHz) and 3.6 cm (8.44 GHz) towards the source J041757 using the Very Large Array (VLA) of the National Radio Astronomy Observatory (NRAO).⁵ Observations were performed on 2008 September 4 (project AP566) and on 2009 February 25 (project AD593), with the array in D and B configurations, respectively. We used the standard VLA continuum mode (4IF, 50 MHz per IF), and both right and left polarizations were processed. The phase centre of the observations was RA (J2000.0) = 04^h 17^m 57^s.8, Dec. (J2000.0) = +27° 41′ 05″.0 in both epochs. For the first set of data (D configuration of the array) and for each frequency, we used $\simeq 40$ min on source, and for the second set (B configuration) we integrated $\simeq 1.5$ h on source at each frequency. See Table 3 for further details on the observations.

Both sets of data were obtained during the Extended Very Large Array (EVLA) transition mode. Calibration and data reduction were performed using the Astronomical Image processing System (AIPS, developed by NRAO), and following NRAO official recommendations for the EVLA transition mode.⁶ As a result, data in D configu-

⁴ In Barrado et al. (2009), we show the CSO emission for J041757 assuming that the source is extended, i.e. allowing us to recover emission from structures > 30 arcsec. For this case, the source associated with J041757 is ~ 30 arcsec, and has a small extension to the north-east. However, the ‘compact’ option used in this work, although yielding more negative features, recovers a structure which is more consistent with independent observations of the region (as shown in this paper). For the J042118 region, the assumption of ‘extended’ emission (forcing the software to recover structures < 60 arcsec to avoid sky gradients) yields very similar results to the assumption of ‘compact’ emission.

⁵ The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

⁶ During the transition to the EVLA, the 50 MHz continuum mode was affected by high closure errors on EVLA–VLA baselines (affecting strongly the 3.6 cm receiver; for this reason, Barrado et al. (2009) report only the flux density at 6 cm). This problem is caused by non-matched bandpass shapes of different type of antennas. In this work, we tried to fix this problem by observing a strong source with known structure and determining baseline-based closure corrections using the AIPS task BLCAL.

Table 1. Summary of observations and catalogues used in this work.

Telescope/Instrument	Wavelength (μm)	Observing mode	Sources observed	Date (yy/mm/dd)	HPBW ^a (arcsec)	P.A. ^a ($^\circ$)
CFHT/MegaPrime-MegaCam	0.75, 0.90	Photometry	J041757	04/12/05 ^b	~ 1	–
APO/SDSS	0.89	Photometry	J042118	2004	~ 1	–
CAHA/Omega2000	1.25, 1.65, 2.17	Photometry	J041757	07/11/04 ^b	~ 1	–
UKIDSS/WFCAM	1.25, 1.63, 2.20	Photometry	J042118	07/09/28	~ 1	–
<i>Spitzer</i> /IRAC	3.6, 4.5, 5.8, 8.0	Photometry	Table 2	05/02/21 ^b	1.4–1.9	–
WISE/SC	12,22	Photometry	J041757, J042118	2010	6.5–12	–
<i>Spitzer</i> /MIPS	24,70	Photometry	Table 2	05/02/28 ^b	6–18	–
CSO/SHARC-II	350	Cont. imaging	J041757, J042118	08/01/28	10	–
APEX/LABOCA	870	Cont. imaging	5 sources, Table 2	09/07	18	–
IRAM 30 m/MAMBO-II	1200	ON–OFF	Table 2	07/05, 07/11	11	–
IRAM 30 m/EMIR	2700, ¹³ CO(1–0)	Line imaging	J041757	09/10/28	22	–
IRAM 30 m/EMIR	2600, ¹² CO(1–0)	Line imaging	J041757	09/10/28	22	–
IRAM 30 m/EMIR	3200, N ₂ H ⁺ (1–0)	Freq. switch.	J041757	10/08/13	27	–
VLA	36 000	Cont. imag., D ^c	J041757	08/09/04	10 \times 9	–68 $^\circ$
VLA	36 000	Cont. imag., B ^c	J041757	09/02/25	1.7 \times 1.6	–89 $^\circ$
VLA	60 000	Cont. imag., D ^c	J041757	08/09/04	17 \times 16	+77 $^\circ$
VLA	60 000	Cont. imag., B ^c	J041757	09/02/25	2.4 \times 1.8	+61 $^\circ$

^aHalf power beam width (HPBW) and position angle (P.A.) of the beam.

^bSome data already published in Barrado et al. (2009).

^c‘B’ or ‘D’ stands for the array configuration.

Table 2. IRAM 30 m/MAMBO-II at 1.2 mm and Archive APEX/LABOCA at 870 μm observations for the sample of proto-BD candidates.

Source	Position		Int. time ^a (min)	$S_{\nu}^{1.2\text{mm}b}$ (mJy)	M_{env}^c (M_{Jup})	$S_{\nu}^{870\mu\text{m}d}$ (mJy)
	α (J2000)	δ (J2000)				
SSTB213 J041726.38+273920.0	04 ^h 17 ^m 26 ^s .38	+27 $^\circ$ 39′ 20″.0	40	<4.2	<2.7	<84
SSTB213 J041740.32+282415.5	04 ^h 17 ^m 40 ^s .32	+28 $^\circ$ 24′ 15″.5	40	<4.2 ^e	<2.7	–
SSTB213 J041757.77+274105.0	04 ^h 17 ^m 57 ^s .77	+27 $^\circ$ 41′ 05″.0	100	<2.9 ^b	<1.8	<84
SSTB213 J041828.08+274910.9	04 ^h 18 ^m 28 ^s .08	+27 $^\circ$ 49′ 10″.9	20	<5.4	<3.4	<480
SSTB213 J041836.33+271442.2	04 ^h 18 ^m 36 ^s .33	+27 $^\circ$ 14′ 42″.2	20	<7.2	<4.6	<96
SSTB213 J041847.84+274055.3	04 ^h 18 ^m 47 ^s .84	+27 $^\circ$ 40′ 55″.3	20	<5.8	<3.7	<88
SSTB213 J041913.10+274726.0	04 ^h 19 ^m 13 ^s .10	+27 $^\circ$ 47′ 26″.0	20	<5.9	<3.7	–
SSTB213 J041938.77+282340.7	04 ^h 19 ^m 38 ^s .77	+28 $^\circ$ 23′ 40″.7	20	<5.0	<3.2	–
SSTB213 J042016.70+280033.7	04 ^h 20 ^m 16 ^s .70	+28 $^\circ$ 00′ 33″.7	20	<5.7	<3.6	–
SSTB213 J042019.20+280610.3	04 ^h 20 ^m 19 ^s .20	+28 $^\circ$ 06′ 10″.3	20	<5.4	<3.4	–
SSTB213 J042118.43+280640.8	04 ^h 21 ^m 18 ^s .43	+28 $^\circ$ 06′ 40″.8	40	<4.0 ^b	<2.5	–
SSTB213 J042123.70+281800.6	04 ^h 21 ^m 23 ^s .70	+28 $^\circ$ 18′ 00″.6	20	<5.6	<3.5	–

^aTotal integration time for IRAM 30 m/MAMBO-II ON–OFF observations.

^bUpper limits correspond to 4σ . For the case of J041757 and J042118, the weighted mean of the different subscans was 1.89 ± 0.72 and 2.15 ± 1.00 mJy, respectively, both above 2σ , while for the other sources the weighted mean was well below 2σ .

^cMasses of disc/envelope estimated from the 1.2 mm continuum emission, assuming that the emission is optically thin, a dust temperature of 15 K and a dust opacity at 1.2 mm of $0.009\text{ cm}^2\text{ g}^{-1}$ (Ossenkopf & Henning 1994).

^dUpper limits correspond to 4σ and are measured in the archive APEX/LABOCA image of the B213/B211 filament (see Section 2.2).

^eNo reliable flux due to the presence of a submillimetre source (J0417426+282442, detected by *AKARI* with a flux of 3.6 Jy at 140 μm) within the wobbler throw. We estimated that up to 6 mJy could have been subtracted to this source. This is the only source of our sample with an *AKARI* submillimetre source within the wobbler throw of the IRAM 30 m/MAMBO observations.

ration at both 6 and 3.6 cm were successfully calibrated. As for the second data set (B configuration), anomalous data affecting the flux calibrator were flagged at 6 cm.⁷ Thus, we used data in B configuration just to illustrate the morphology of the emission at higher

⁷For the B configuration data, anomalous jumps in the amplitudes and phases for certain EVLA antennas, specially during the end of the target and flux calibrator observation, prevented us from a reliable flux calibration. These anomalous data were flagged only for the 6 cm case, because the 3.6 cm observations were too short. However, flagging these data with

angular resolution, and for flux measurement purposes we used only the first data set (D configuration). For the D configuration data, we produced radio continuum maps with natural weighting to optimize the sensitivity of the images, achieving rms noises of $\simeq 0.025$ and

anomalous jumps improved the signal-to-noise ratio, while the morphology of the object was found unaffected, giving us confidence on the morphology of the source seen in the B configuration data at both 6 and 3.6 cm.

Table 3. Parameters of the VLA radio continuum observations towards J041757.

Config.	λ (cm)	Flux calib.	Flux (Jy)	Phase calib.	Flux ^a (Jy)
D	6	3C147	7.9	J0431+206	2.34 ± 0.03
D	3.6	3C147	4.7	J0431+206	1.590 ± 0.005
B	6	3C138	3.6	J0431+206	2.63 ± 0.05
B	3.6	3C138	2.4	J0431+206	1.06 ± 0.04

^aBootstrapped flux.

0.055 mJy beam⁻¹ at 6 and 3.6 cm, respectively. For the B configuration data, we produced maps with natural weighting and used a taper in the uv data of 150 k λ at 6 cm and 120 k λ at 3.6 cm. For the B configuration data, we refrain from giving the rms noises because of anomalous data for the flux calibrator.

2.5 IRAM 30m CO, ¹³CO and N₂H⁺ emission for J041757

J041757 was observed in the CO(1–0) (115.271 204 GHz) and ¹³CO(1–0) (110.201 370 GHz) transitions with the IRAM 30 m telescope using the EMIR (Carter et al. 2012) heterodyne receivers. The observations were carried out on 2009 October 28 and 29, with system temperatures of 200–400 K ($\tau \sim 0.16$ –0.35). We conducted on-the-fly maps in the position-switching mode, scanning the region in RA, and with steps of 4 arcsec in Dec., covering a size of 1.6×1.6 arcmin² centred on J041757. For each of the CO(1–0) and ¹³CO(1–0) transitions, we used two units of the Versatile Spectrometer Array (VESPA) backend, one in each polarization, of 40 kHz (0.11 km s⁻¹) of spectral resolution and a bandwidth of 40 MHz (~ 100 km s⁻¹). Simultaneously to the VESPA backend, we also used the Wideband Line Multiple Autocorrelator (WILMA) backend, providing a bandwidth of 9000 km s⁻¹ with ~ 5 km s⁻¹ of spectral resolution at the frequency of the CO(1–0) transition. The OFF position used, at (–7600, –287) arcsec from the map centre, was selected as free of CO from a survey in Taurus (Goldsmith et al. 2008) and was proved to be free of emission. Calibration following the standard procedures was carried out during observing time and using the MIRA package, and typical calibration errors are within 15–20 per cent. To make the final map, spectra were averaged every 8 arcsec in both RA and Dec. Typical rms noise of spectra at each point of the final map were around 0.24 and 0.12 K for CO(1–0) and ¹³CO(1–0), respectively, in the main beam brightness temperature scale. The main beam efficiency at the CO(1–0) line frequency is 0.82 ($F_{\text{eff}} = 94.01$ per cent, $B_{\text{eff}} = 77.53$ per cent) and at the ¹³CO(1–0) frequency is 0.83 ($F_{\text{eff}} = 94.18$ per cent, $B_{\text{eff}} = 78.13$ per cent).

Subsequent observations of the dense gas tracer N₂H⁺(1–0) (at 93.173 764 GHz) were carried out on 2010 August 13 under reasonable summer weather conditions (p_{wv} ~ 4 mm; system temperature ~ 130 K; opacity ~ 0.4), in frequency switching mode, and using VESPA with 40 MHz of bandwidth and 0.02 MHz of spectral resolution. We integrated at different positions in and around J041757, separated ~ 20 arcsec and covering offsets with respect to J041757 from (40,40) arcsec to (–20, –20) arcsec, reaching rms noises in each individual spectra of ~ 0.015 K (main beam brightness temperature). The main beam efficiency at the N₂H⁺(1–0) frequency is 0.85 ($F_{\text{eff}} = 94.76$ per cent, $B_{\text{eff}} = 80.15$ per cent). All data reduction was done using the GILDAS software package supported at IRAM.

3 RESULTS

3.1 J042118

Although the 1.2 mm emission for J042118 was very marginal (2.2σ , Table 2), the CSO observations at 350 μ m reveal one clear source with a flux level of $5\sigma \sim 46$ mJy (Fig. 2, top). The position of the submillimetre source, which is unresolved, is coincident with the position of the *Spitzer*/IRAC and United Kingdom Infrared Digital Sky Survey (UKIDSS) source, and there is marginal




Figure 2. CSO 350 μ m signal-to-noise ratio contours superimposed on an NIR image. Top: J042118. Contours are –4, –3 (dashed), 3, 4 and 5 (solid) times the rms noise of the map [noise is variable across the map, from 9 (centre) to 11 mJy beam⁻¹ (wedges)]. Grey-scale is the UKIDSS K -band image. Bottom: J041757. Black contours correspond to the CSO 350 μ m emission and are –5, –4, –3 (dashed), 3, 4, 5, 6, 7, 8, 9 and 10 (solid) times the rms noise of the map [noise is variable across the map, from 6 (centre) to 9 mJy beam⁻¹ (wedges)]. White contours correspond to the APEX 870 μ m emission and are 1, 2, 3, 4 and 5 times 19 mJy beam⁻¹. The APEX beam at 870 μ m is 18 arcsec. Grey-scale is the Centro Astronómico Hispano-Alemán (CAHA)/Omega2000 H -band image (see Barrado et al. 2009 for details on the observations), showing the two NIR sources, A and B, found near the IRAC source. In both panels, the CSO beam, shown in the bottom-right corner, is 10 arcsec, and the crosses indicate the position of the *Spitzer*/IRAC source.

Table 4. Parameters of the sources detected with CSO/SHARC-II at 350 μm .

Source	rms ^a (mJy)	Position ^b		Deconvolved ang. size (arcsec)	Deconvolved P.A. (°)	Deconvolved size (au)	$I_{\nu}^{\text{peak } c}$ (mJy beam ⁻¹)	S_{ν}^c (mJy)	Mass ^d (M_{Jup})
		α (J2000)	δ (J2000)						
J042118	8.9	04 ^h 21 ^m 18 ^s .38	28° 06′ 40″.1	10.5 × 8.7	58.8	1500 × 1200	46 ± 9	46	0.3–3
J041757	6.1	04 ^h 17 ^m 57 ^s .67	27° 41′ 05″.8	19.5 × 12.7	−67.3	2700 × 1800	37 ± 6	165	1–10
J041757-NE	8.6	04 ^h 18 ^m 00 ^s .30	27° 41′ 36″.3	29.2 × 15.4	29.7	4100 × 2100	84 ± 9	356	2–21

^aThe rms noise at the position of the source. Total integration time was around 1 h 30 min for J041757 and around 1 h for J042118.

^bPositions corresponding to the intensity peak.

^cPeak intensity and flux density derived by computing statistics within the aperture corresponding to the 3σ level of the source.

^dMasses derived assuming a dust temperature of 10–20 K, and a dust mass opacity coefficient from Ossenkopf & Henning (1994, see the main text). The uncertainty in the masses due to the opacity law is estimated to be a factor of 2.

extended and faint emission at 350 μm about 10 arcsec to the west. In Table 4, we give the position, size, flux density and estimated mass of the submillimetre source detected in J042118. The mass of the disc/envelope obtained for J042118 is in the range 0.3–3 M_{Jup} , assuming a distance of 140 pc, a dust temperature in the range 10–20 K, and a dust opacity law at 350 μm of 0.101 $\text{cm}^2 \text{g}^{-1}$ (Ossenkopf & Henning 1994). Such a mass is at least three orders of magnitude larger than the disc masses estimated from *Herschel* observations for a small sample of young BDs (Harvey et al. 2012).

3.2 J041757

3.2.1 CSO 350 μm and APEX 870 μm continuum emission

J041757, with a 2.6σ flux at 1.2 mm (Table 2), is well detected at 350 μm , and presents a complex emission, with one partially resolved source (~ 16 arcsec) associated with the selected IRAC source J041757, at the centre of the field of view, and one stronger and more extended source (~ 20 arcsec) to the north-east (hereafter, J041757-NE), which is elongated in the southwest–northeast direction (Fig. 2, bottom, and Table 4). The flux density of J041757 at 350 μm is 165 mJy, yielding a mass of the envelope of 1–10 M_{Jup} (for a dust temperature of 10–20 K). Assuming a mean molecular weight (of H_2 molecule) of 2.8 m_{H} and a size (diameter) of 2200 au, we obtain a density of $\lesssim 2.1 \times 10^5 \text{cm}^{-3}$, an H_2 column density of $\lesssim 4.6 \times 10^{21} \text{cm}^{-2}$ and $A_{\nu} \lesssim 4.9$ mag (following Frerking, Langer & Wilson 1982). Similarly, for J041757-NE, we estimate a mass of 2–20 M_{Jup} , a density of $\lesssim 2.4 \times 10^5 \text{cm}^{-3}$, an H_2 column density of $\lesssim 6.4 \times 10^{21} \text{cm}^{-2}$ and $A_{\nu} \lesssim 6.7$ mag. These A_{ν} values are in perfect agreement with the values derived by Schmalzl et al. (2010) for this region. It is interesting to note that J041757-NE is also recovered in the archived APEX/LABOCA 870 μm image (see Fig. 2, bottom) for which we measured a peak intensity of 100 mJy beam⁻¹ and a flux density (within the 2σ contour) of 420 mJy. This corresponds to a mass of 75–160 M_{Jup} (assuming a dust temperature of 10–15 K, and a dust opacity law at 870 μm of 0.018 $\text{cm}^2 \text{g}^{-1}$, Ossenkopf & Henning 1994). The mass of the J041757-NE source as estimated from the APEX/LABOCA image is about a factor of 8 higher than the mass estimated from the CSO because structures >30 arcsec were filtered out during the reduction process of the CSO data (Section 2.3). Thus, the total mass of J041757-NE is $\sim 100 M_{\text{Jup}}$, from which $\sim 10 M_{\text{Jup}}$ are found in a compact structure.

3.2.2 VLA 3.6 and 6 cm continuum emission

The VLA centimetre emission around J041757 reveals one single source at both 3.6 and 6 cm wavelengths at the position of J041757, with no significant emission at the position of J041757-NE.

Table 5. VLA radio continuum emission detected at 3.6 and 6 cm towards J041757.

λ (cm)	Position ^a		S_{ν}^b (mJy)	Size, P.A. ^a (arcsec ² , °)
	α (J2000)	δ (J2000)		
6	04 ^h 17 ^m 57 ^s .74	+27° 41′ 04″.2	0.20 ± 0.05	2.4 × 1.3, 45
3.6	04 ^h 17 ^m 57 ^s .75	+27° 41′ 04″.4	0.19 ± 0.11	2.9 × 0.8, 40

^aResults based on observations in the B configuration of the VLA (see Section 2.4) and derived from Gaussian fits. Uncertainties in sizes are around 0.5 arcsec and in P.A. are around 10°. Sizes and P.A. are deconvolved.

^bResults based on observations in the D configuration of the VLA. Uncertainties are 2σ .

Using D configuration only, the emission is unresolved at both frequencies, and it is detected at $\approx 7\sigma$ at 6 cm, and marginally detected at $\approx 4\sigma$ at 3.6 cm, with a flux density at both wavelengths of ~ 0.2 mJy (Table 5). The spectral index in the range of 6–3.6 cm is $\alpha_{6-3.6\text{cm}} = 0.0 \pm 0.5$ (1σ error; where $S_{\nu} \propto \nu^{\alpha}$), which is indicative of optically thin free–free emission (typically arising in shocks for the case of low-mass young stellar objects and VeLLOs, see Section 5.2.3). To extract the morphological information of the centimetre emission, we imaged the B configuration data only, and the resulting images are presented in Fig. 3. The figure shows first that the centimetre source is clearly associated with the NIR source ‘B’ detected by Barrado et al. (2009), while source ‘A’ shows no centimetre emission. Furthermore, the figure reveals at both 6 and 3.6 cm a faint extension in the southwest–northeast direction. A Gaussian fit to the source yields, at both wavelengths, a deconvolved size of approximately $\sim 2.5 \times 1.0 \text{arcsec}^2$, oriented at P.A. $\sim 40^\circ$ (Table 5).

3.2.3 IRAM 30 m molecular emission

The CO (1–0) spectra averaged over a region of 40 arcsec of diameter around J041757 (left) and J041757-NE (right) are shown in Fig. 4, top. The spectra reveal the three well-known velocity components associated with the Taurus molecular cloud, at around 5, 7 and 8 km s^{-1} (e.g. Heyer et al. 1987; Mizuno et al. 1995). Among the three main velocity components seen in Taurus, the velocity maps of fig. 6 of Mizuno et al. (1995) show that at the position of J041757 there are two intersecting filamentary structures with different velocities: one in the 4.5–6.0 km s^{-1} range and the other in the 6.0–7.5 km s^{-1} range. We made line area maps by integrating the CO emission in ranges of 1 km s^{-1} wide, from 4.5 to 8.5 km s^{-1} , and found that for this velocity range CO traces the large-scale structure of the B211 filament (located to the southwest of J041757, see Fig. 1), with strong intensity gradients and no clear features associated with J041757 (Fig. 5, top). This was expected




Figure 3. High angular resolution VLA radio continuum emission (red contours) towards J041757, obtained with the array in B configuration, superimposed on the Omega2000 *H*-band image (grey-scale) and the CSO 350 μm image (black contours). Top: 6 cm emission. Red contours are -4 , -3 (dashed), 3, 4, 6, 8, 10 and 12 (solid) times the rms of the map (rms is not given due to technical problems with the flux density scale during the B configuration observations, see Section 2.4). A synthesized beam is shown in the bottom-right corner and is $2.45 \times 1.77 \text{ arcsec}^2$, with P.A. = 61° . Bottom: 3.6 cm emission. Red contours are -4 , -3 (dashed), 3, 4, 5 and 6 (solid) times the rms of the map. A synthesized beam is shown in the bottom-right corner and is $1.74 \times 1.64 \text{ arcsec}^2$, with P.A. = -89° . In both panels, the grey-scale and black contours are the same as in Fig. 2, bottom.

because CO is optically thick and spread out over all the Taurus region (e.g. Goldsmith et al. 2008; Narayanan et al. 2008). However, we fitted with Gaussians the three velocity components seen in the CO spectrum and found, for the emission line at 5 km s^{-1} , an excess in the blueshifted emission with respect to the Gaussian fit.⁸ The line area map integrated in the range $2\text{--}4 \text{ km s}^{-1}$, where the blueshifted excess is seen, indicates that this emission is strongest around J041757, suggesting a possible relationship (Fig. 5, top).

Concerning $^{13}\text{CO}(1\text{--}0)$, well suited to trace the gas column density in the region because it is optically thinner than $^{12}\text{CO}(1\text{--}0)$, the averaged spectra also reveal at least three velocity components for both J041757 and J041757-NE. In Fig. 5, bottom, we show the line area maps of the ^{13}CO emission integrated between 4.5 and 8.5 km s^{-1} in steps of 1 km s^{-1} . Among the four mapped velocity intervals, the emission integrated between 6.5 and 7.5 km s^{-1} is more prominent at the position of J041757-NE, and has an extension to the south and another extension or stream to the southwest, passing through J041757. This could be a hint of the molecular gas

⁸ An excess with respect to a Gaussian fit was also found with the Kitt Peak Telescope in $^{12}\text{CO}(1\text{--}0)$ between 2 and 4 km s^{-1} (Morata, private communication), with the integrated emission also concentrated on J041757.




Figure 4. Left-hand column: spectra averaged in a region of $\pm 20 \text{ arcsec}$ with respect to the position of J041757. Right-hand column: idem for J041757-NE [offset of (37, 30) arcsec with respect to J041757]. Note the excess of blueshifted emission at velocities in the range $2\text{--}4 \text{ km s}^{-1}$ for CO(1-0). For the case of N_2H^+ , J041757 was fitted with one Gaussian, while for J041757-NE we did fit the hyperfine structure of N_2H^+ .

associated with J041757-NE and J041757 coming from the same physical region.

Finally, the $\text{N}_2\text{H}^+(1\text{--}0)$ averaged spectra for J041757 and J041757-NE reveal emission associated with J041757-NE (see Fig. 4, bottom). We fitted the hyperfine structure of the $\text{N}_2\text{H}^+(1\text{--}0)$ transition for J041757-NE using the package CLASS of the GILDAS software, and obtained the output parameter $A\tau = 0.059 \pm 0.006 \text{ K}$,⁹ $V_{\text{LSR}} = 7.02 \pm 0.02 \text{ km s}^{-1}$, $\Delta v = 0.42 \pm 0.17 \text{ km s}^{-1}$ and an opacity of the $F_1'F_1''\text{--}F_1F_1'$ transition of ~ 0.1 , which corresponds to an excitation temperature of 3.4 K and an N_2H^+ column density of $2.5 \times 10^{11} \text{ cm}^{-2}$. This corresponds to a total H_2 column density of $\sim 9 \times 10^{20} \text{ cm}^{-2}$ (assuming an N_2H^+ abundance of $(2\text{--}5) \times 10^{-10}$; Pirogov et al. 2003; Tafalla et al. 2004) and a mass of $\sim 3 M_{\text{Jup}}$ (for a size of $\sim 20 \text{ arcsec}$ in diameter), similar to the lower limit of the mass derived from the submillimetre continuum emission (Table 4). Towards J041757, the strongest hyperfine component of the $\text{N}_2\text{H}^+(1\text{--}0)$ transition is detected (at 5σ), and a Gaussian fit yields $7.04 \pm 0.08 \text{ km s}^{-1}$, very similar to the velocity of the J041757-NE source. This suggests that both objects are associated with dense material [as the critical density of $\text{N}_2\text{H}^+(1\text{--}0)$ is $\sim 10^5 \text{ cm}^{-3}$] possibly coming from the same region of Taurus, as also indicated by the morphology of the ^{13}CO emission integrated between 6.5 and 7.5 km s^{-1} (Fig. 5, bottom).

4 SPECTRAL ENERGY DISTRIBUTIONS

Both J042118 and J041757 are *Spitzer*/IRAC sources emitting at $350 \mu\text{m}$. In order to further constrain its nature, we checked the NASA Extragalactic Database (NED), the National VLA Sky Survey (NVSS) at 21 cm and JCMT/SCUBA catalogue at 450 and

⁹ In the CLASS package of the GILDAS software, the first output parameter of a fit to the hyperfine structure of a given transition is $A\tau = f(J_\nu(T_{\text{ex}}) - J_\nu(T_{\text{bg}}))\tau$, where f is the filling factor, assumed to be 1, $J_\nu(T) = h\nu/k/(e^{h\nu/kT} - 1)$, with k being the Boltzmann constant, T the temperature and ν the frequency; T_{ex} is the excitation temperature, T_{bg} the background temperature and τ is the opacity of the main hyperfine component, $F_1'F_1''\text{--}F_1F_1'$ for the case of $\text{N}_2\text{H}^+(1\text{--}0)$.




Figure 5. Top: $^{12}\text{CO}(1-0)$ channel maps in the J041757 region. The rms is 0.16 K (or 0.23 K for the panel showing the emission integrated from 2 to 4 km s $^{-1}$). Bottom: $^{13}\text{CO}(1-0)$ channel maps in the J041757 region. The rms is 0.05 K. For all panels, colour scale is in main beam brightness temperature (K), the beam is ~ 22 arcsec and stars mark the position of J041757 and J041757-NE.

850 μm (di Francesco et al. 2008), and found no data for SCUBA, no significant emission in NVSS above 1 mJy beam $^{-1}$ and no registered object in NED. To look for further archival data, we used the new capabilities of the Virtual Observatory Spectral energy distribution Analyzer (VOSA; Bayo et al. 2008; Bayo et al., in preparation). The tool queries virtual observatory compliant catalogues in the ultraviolet (*GALEX*, *International Ultraviolet Explorer* HPDP photometry), optical (Tycho-2, CMC-14; Hauck & Mermilliod 1998 Mermilliod & Mermilliod 1994, Sloan DSS (SDSS) release 7 and IPHAS), NIR, MIR and far-infrared (FIR) regimes (2MASS, *IRAS*, MSX6C, *AKARI* IRC and FIS, GLIMPSE, UKIDSS data release 7 (DR7), WISE preliminary release), and allows us to fit several grids of models with extinction as a free parameter. Apart from the data already presented in this work and in Barrado et al. (2009), we find UKIDSS counterparts for both sources but with unreliable photometry in the *H* band, and WISE counterparts at less than 1 arcsec from the Omega2000 position of J041757 (NIR component B) and the *Spitzer*/IRAC position of J042118. Since SDSS did not detect any source at the position of J041757 and J042118, we give only upper limits at the corresponding frequencies. In Appendix A, we list the flux density measurements and upper limits (Tables A1 and A2), and in Fig. 6 we present the spectral energy distributions (SEDs) for both J042118 and J041757.

5 DISCUSSION

5.1 On the nature of J042118

J042118 is undetected in the SDSS (DR 8) but detected in UKIDSS (DR 7, Fig. 2), WISE, *Spitzer*/IRAC+MIPS and the CSO (see Table A1). In order to test if the J042118 SED (Section 4) is easily reproducible by a very extinguished photospheric model (as expected if it was a background object seen through the molecular cloud), we used VOSA tool (Section 4) and found that even allowing extinction values up to $A_v \sim 15$ mag, it is not possible to get a reasonable fit (reduced χ^2 values of the order of 100) neither with Kurucz (Castelli, Gratton & Kurucz 1997) nor with Lyon models (Allard et al. 1997, 2001, 2003, 2007; Baraffe et al. 1997, 1998; Hauschildt, Allard &




Figure 6. SED of J042118 (black line) and J041757, NIR component B (brown line), after Barrado et al. (2009).

Baron 1999; Chabrier et al. 2000; Allard 2009), suggesting that the object belongs to Taurus.

With the aim of shedding light on the nature of J042118, we compared the CSO emission to the *Herschel* Archive images at 250 and 160 μm ,¹⁰ which are the wavelengths for which the *Herschel* beam (17 and 11 arcsec) is more similar to the CSO beam (10 arcsec). While at 160 μm there is no significant emission towards J042118, the *Herschel*/SPIRE 250 μm emission shows an extended large-scale structure elongated in the southwest–northeast direction (see Fig. 7), with a subcondensation matching well the

¹⁰ *Herschel* Archive data from projects within the Gould Belt Key Programme (see André et al. 2010) were downloaded (observing identification numbers 1342202090 and 1342202254). The *Herschel*/PACS(SPIRE) data at 160(250) μm were processed with HIPE(madMap) version 8 and SCANAMORPHOS version 14 (Roussel 2012). The *Herschel* data towards B211/L1495 will be published in a forthcoming paper by Palmeirim et al. as part of the ‘*Herschel* Gould Belt Survey’ project.




Figure 7. Submillimetre emission towards J042118. The grey-scale and black contours correspond to the *Herschel*/SPIRE 250 μm emission. Contours range from 0.06 to 0.12, increasing in steps of 0.02 Jy beam^{-1} (1σ is approximately 0.009 Jy beam^{-1}). The *Herschel* beam at 250 μm , shown in the bottom-left corner, is ~ 17 arcsec. White contours are the CSO 350 μm emission (same contours as in Fig. 2) and the CSO beam is ~ 10 arcsec.

position of J042118, within the telescope positional uncertainties. The location of J042118 at the tip of the elongated structure, which seems to be part of the extended structures of the Taurus complex, favours its membership to Taurus. In this case, the submillimetre source detected with the CSO would be tracing a disc and/or small (< 5 arcsec or 700 au of radius) envelope with a mass of $\sim 1 M_{\text{Jup}}$ (Table 4). The luminosity (upper limit) for J042118 estimated from the SED (Table A1) would be $\lesssim 0.0023 L_{\odot}$, and the SED shape is reminiscent of Class I(II) young stellar objects, with a flat slope between the NIR and the submillimetre range. This can be quantified by estimating the bolometric temperature, which is ~ 140 K,¹¹ typical of Class I young stellar objects (e.g. Chen et al. 1995). The fact that the submillimetre emission, as recovered by the CSO, is unresolved could be indicative of most of the envelope being accreted/dispersed, with the submillimetre emission arising only from a compact disc.

5.2 On the nature of J041757

The nature of J041757 is discussed in Barrado et al. (2009), who present series of colour–colour and colour–magnitude diagrams from *Spitzer* and NIR data to conclude that the most likely scenario for J041757 (for either component A and/or B) is that the object is galactic. Subsequent IR spectroscopic observations by Luhman & Mamajek (2010) indicate that component A is most likely an M2 background star, while no conclusion could be drawn for component B because no spectrum was shown for this component in that work. This, together with the fact that source B has a raising slope in the NIR SED matching well with the FIR and submillimetre emission from *Spitzer*/MIPS and CSO (Barrado et al. 2009), indicates that the most likely counterpart of the submillimetre source in J041757 is the NIR source B, which in turn is associated with centimetre emission

¹¹ This number must be regarded with caution because the SED for this source is better sampled in the IR than in the submillimetre and radio range and because we treated all flux densities which are upper limits as detections (c.f. Kauffmann et al. 2008).




Figure 8. J041757: in both panels, colour scale is the $^{13}\text{CO}(1-0)$ area integrated in the range 6.5–7.5 km s^{-1} (as in Fig. 5) and white contours are the CSO 350 μm emission (same contours as in Fig. 2). Top: Black contours correspond to the *Herschel*/SPIRE 250 μm emission. Contours range from 0.30 to 0.54, increasing in steps of 0.03 Jy beam^{-1} (1σ is approximately 0.014 Jy beam^{-1}). The *Herschel* beam at 250 μm , shown in the bottom-right corner, is ~ 17 arcsec (^{13}CO beam is ~ 22 arcsec). Bottom: Black contours correspond to the *Herschel*/PACS 160 μm emission. Contours range from 0.14 to 0.24, in steps of 0.014 Jy beam^{-1} (1σ is approximately 0.018 Jy beam^{-1}). The *Herschel* beam at 160 μm , shown in the bottom-right corner, is ~ 11 arcsec, similar to the CSO beam.

(Section 3.2.2). Hereafter, we will refer to source B (identified in the NIR and in the centimetre range) as J041757-B.

Similarly to the case of J042118, we compared the CSO emission of J041757 to the *Herschel* Archive images at 250 and 160 μm . See Fig. 8 to visualize such a comparison. The *Herschel*/SPIRE emission at 250 μm (black contours in Fig. 8, top) is strongest at the position of J041757-NE, matching well the 350 μm emission observed with the CSO (white contours). From J041757-NE, there are two streams of emission, extending for about ~ 1 arcmin, one towards the south and the other towards the southwest and passing through J041757-B. The 250 μm emission towards the southwestern side of the map is part of the B211 filament (Fig. 1). We note the remarkable similarity between the emission at 250 μm and the $^{13}\text{CO}(1-0)$ emission integrated between 6.5 and 7.5 km s^{-1} (colour scale in Fig. 8, top). Compared to *Herschel* data, the CSO is recovering the most compact (as we used the ‘compact’ option in the reduction process, see Section 2.3) and fainter emission (1σ noise of *Herschel*/SPIRE at 250 μm : ~ 14 mJy beam^{-1} ; 1σ noise of CSO




Figure 9. SED of J041757-B (black) compared to average SEDs for radio-loud (thick red) and radio-quiet (thick blue) AGNs from Shang et al. (2011). Thin blue lines are the extinguished radio-quiet SEDs with A_v of 5 and 10 mag. The spectrum of J041757-B in the visible range will be presented in a forthcoming paper (Huélamo et al., in preparation).

at $350\ \mu\text{m}$: $\sim 6\ \text{mJy beam}^{-1}$). Regarding the emission seen with *Herschel*/PACS at $160\ \mu\text{m}$, this is again strongest at J041757-NE, and the northeast–southwestern stream splits up into one slightly resolved source coinciding, within the position uncertainties of the telescopes, with the $350\ \mu\text{m}$ source at the position of J041757-B. This suggests that J041757-B is associated with the hottest dust of the stream linking the J041757-NE source and the B211 filament. Flux densities measured by *Herschel* at 160 and $250\ \mu\text{m}$ are consistent with the flux densities given here as measured with the CSO, but we refrain from presenting a detailed study of the *Herschel* data in this work as this will be the subject of a forthcoming paper.

5.2.1 J041757-B: an extragalactic object?

We consider first the possibility that J041757-B is an extragalactic object, which would be aligned by chance with the submillimetre stream linking the J041757-NE source and the B211 filament. In Fig. 9, we show a comparison of the SED for J041757-B to the recent compilation of average SEDs of radio-loud and radio-quiet active galactic nuclei (AGNs) of Shang et al. (2011, which are in good agreement with the classical SEDs of Elvis et al. 1994). The SEDs are normalized at $12\ \mu\text{m}$ (a good intrinsic indicator for AGNs; e.g. Gandhi et al. 2009) to the flux of J041757-B, and the error bars are the standard deviation of the fluxes for the Shang et al. sample, being representative of the scatter in the AGN fluxes at each wavelength. From the figure, it is seen that J041757-B SED is not compatible with the radio-loud AGN SEDs, mainly because of its important drop from MIR to radio wavelengths.¹² In the visible range, J041757-B SED is much more obscured than AGN SEDs. In order to simulate an AGN extinguished SED, we reddened the AGN average SED for radio quiet using the extinction law by Fitzpatrick (1999; improved by Indebetouw et al. 2005 in the IR and $R_v \sim 3.1$) and with A_v ranging from 1 to 10 (see figure, thin blue lines). The result is that the J041757 SED would be compatible with an AGN SED extinguished by $A_v \sim 5$ in the visible IR range. However, the

¹² We note however that if we estimate the radio loudness from the classical definition, $S_{6\text{cm}}/S_{4400\text{\AA}}$ (following Kellermann et al. 1989), we obtain ~ 700 , which would classify J041757-B as radio loud. As explained in the text, this would be still compatible with a radio-quiet AGN if the extinction for J041757-B is important.

J041757-B SED is not fully consistent with radio-quiet AGNs either, because of its important contribution of submillimetre (and radio) emission compared to MIR. This could be indicating that there is an excess of submillimetre emission in the direction of the galaxy, which could be explained by the Taurus stream joining the J041757-NE source and the B211 filament. Nevertheless, there would be an excess of submillimetre emission from the galaxy candidate itself, as revealed by the image of *Herschel*/PACS at $160\ \mu\text{m}$, which shows that in the stream there is only emission at $160\ \mu\text{m}$ at the position of J041757-B (similar case for the CSO $350\ \mu\text{m}$ emission, Fig. 8). In addition, flat spectral indices in the centimetre range in AGNs are not usual, as centimetre emission from AGNs is typically produced by synchrotron emission from relativistic electrons accelerated in jets (e.g. Giroletti & Panessa 2009; Ibar et al. 2009; Shang et al. 2011; Randall et al. 2012). Furthermore, a preliminary spectrum of J041757-B in the visible range taken with Osiris on the GTC (Huélamo et al., in preparation) shows no broad emission lines, while the extinguished SED in Fig. 9 indicates that even with $A_v \gtrsim 5$ still strong and broad lines are expected for an AGN. If J041757-B was a galaxy it should be detected in forbidden emission lines in the MIR range ([OIV], [NeV]; e.g. Goulding & Alexander 2009) and in hardest X-rays ($> 10\ \text{keV}$),¹³ which could be the final observational tests to definitively discard the nature of J041757-B as a background galaxy.

5.2.2 J041757-B: a background object behind Taurus?

Similarly to J041757-A (Barrado et al. 2009; Luhman & Mamajek 2010), J041757-B could be a background object behind the Taurus complex. However, if J041757-A is a background star, the IR source mainly contributing to the FIR and submillimetre emission must be J041757-B. Therefore, whatever the true nature of J041757-B is, the object is embedded in considerable amounts of dust, and hence should be associated with molecular gas as well. This molecular gas, if J041757-B does not belong to Taurus, should appear as a velocity component in the line of sight different from Taurus velocities, while in a $9000\ \text{km s}^{-1}$ band we did not detect any other line apart from those of Taurus. If the object is associated with dust and gas, it should be a young stellar object (most likely in the Class II/III phase) or an evolved star. However, the centimetre emission for Class II/III objects is typically due to non-thermal gyrosynchrotron emission, which is not resolved (e.g. André et al. 1992, Forbrich et al. 2007) and is polarized, while J041757-B has a flat (thermal) spectral index, is slightly elongated in the southwest–northeast direction and shows no polarized emission (we checked this in the data set of 2008 observations and had no detection of polarized emission). In addition, the lack of X-ray emission (no *ROSAT* detection) is also uncommon for Class II/III objects (e.g. Güdel et al. 2007).

The possibility that J041757 is an evolved stellar object, either a carbon star or a giant, is also considered. Luminous carbon stars are evolved stars with more carbon than oxygen in their atmosphere, characterized by red NIR colours and spectra with no H_2O (Liebert et al. 2000). The *JHK* colours for J041757-B match well within the *JHK* colour–colour diagrams for these stars (e.g. Quanz et al. 2010). However, we used the K_s magnitude of J041757-B (Barrado et al. 2009) and table 8 from Quanz et al. (2010) to estimate the

¹³ In X-rays, we could only set an upper limit for J041757-B, which is not detected in the *ROSAT* All-Sky Survey (Voges et al. 1999, covering the range $0.1\text{--}2.4\ \text{keV}$ with a typical limiting sensitivity of $10^{-13}\ \text{erg cm}^{-2}\ \text{s}^{-1}$; Anderson et al. 2003).

distance and find >160 kpc, larger than the currently known extent of our Galaxy. Since the emission in J041757 is resolved in the sub-millimetre range, such a large distance would imply an unrealistic large size for the envelope of a carbon star of ~ 9 pc. In addition, the carbon star SEDs have the peak at around $12 \mu\text{m}$, and longwards the flux decreases strongly (e.g. Liebert et al. 2000), which is clearly not the case of J041757-B. Thus, the properties of J041757-B are not fully consistent with being either a Class II/III young stellar object, or an evolved star. In fact, by making use of the data compiled to build the SED (Section 4) and using the VOSA tool as outlined in Section 5.1, we found that the J041757-B SED cannot be reproduced with a very extinguished photospheric model, providing further support against the galactic background object scenario for J041757-B.

5.2.3 J041757-B: a proto-BD candidate in Taurus?

Finally, J041757-B could be an embedded object belonging to Taurus. This is strongly suggested by the discovery of a narrow (~ 16 arcsec width) stream joining the J041757-NE source and the B211 filament and passing exactly through the position of J041757-B. Since both J041757-NE and the stream are detected at around 7 km s^{-1} (in N_2H^+ and/or ^{13}CO), both structures belong to Taurus.

If J041757-B belongs to Taurus, its low luminosity, of $\lesssim 0.005 L_{\odot}$ (using the values in Table A2), would imply a proto-BD nature (Barrado et al. 2009). This would naturally explain the centimetre emission with a flat spectral index, since flat/positive spectral indices, indicative of thermal free-free emission, are typically observed in young stellar objects due to shocks in the jet (e.g. Anglada et al. 1998; Avila, Rodríguez & Curiel 2001; Beltrán et al. 2001; Pech et al. 2010; Scaife et al. 2011; Ward-Thompson et al. 2011), and have also been found for VeLLOs (e.g. Shirley et al. 2007, Scaife et al. 2011). In these cases, the gas is ionized in shocks produced by the jet driven by the nascent star, and the centimetre emission is usually elongated along the outflow axis. This would also easily explain the elongation seen at both 6 and 3.6 cm for J041757-B. And interestingly, the $^{12}\text{CO}(1-0)$ excess detected between 2 and 4 km s^{-1} is reminiscent of wings observed in outflows driven by low-mass young stellar objects. In addition, if J041757-B is a proto-BD, its bolometric luminosity and centimetre luminosity match well the relation found for young stellar objects driving radio jets, proposed first by Anglada (1995). Following the recent correlations of Shirley et al. (2007) and Scaife et al. (2011) for the low-mass case, the expected centimetre luminosity, given the bolometric luminosity of J041757-B, is $0.0002-0.007 \text{ mJy kpc}^2$, consistent with the measured centimetre luminosity of $0.004 \pm 0.002 \text{ mJy kpc}^2$. Last, if J041757-B belongs to Taurus, the properties of the extended envelope where the NIR source is embedded, of about ~ 1000 au radius and about $\sim 5 M_{\text{Jup}}$, coincide with the extrapolation of the trend presented by Kauffmann et al. (2008) in a radius versus mass diagram for small c2d *Spitzer* cores (see fig. 4a of Kauffmann et al. 2008), and more recently by Sadavoy et al. (2010) and Schmalzl et al. (2010) for the Taurus region.

Thus, if J041757-B belongs to Taurus, its properties can *naturally* be explained simply as a scaled-down version of low-mass young stellar objects.

5.3 On the nature of J041757-NE: a pre-substellar core?

J041757-NE is the strongest submillimetre source detected in the field of J041757 with the CSO, and has been confirmed by APEX, IRAM 30 m and *Herschel* images. This source is clearly associ-

ated with Taurus, as shown by the N_2H^+ and ^{13}CO emission detected at 7 km s^{-1} (Figs 4 and 5). Since we estimated a total mass (compact+extended emission) for J041757-NE of $\sim 75 M_{\text{Jup}}$ (Section 3.2), and no point IR source (up to $24 \mu\text{m}$) has been found associated with the submillimetre peak, J041757-NE could be harbouring a true pre-substellar core, i.e. a starless core which will form a BD in the future. A first requisite for this to happen is that the submillimetre source should be gravitationally bound. Given the line width measured from the $\text{N}_2\text{H}^+(1-0)$ hyperfine fit (Section 3.2.3), and an average radius of 10 arcsec (Table 4), we estimate a gravitational mass of $\sim 140 M_{\text{Jup}}$ (following equation 4 of Pound & Blitz 1993), similar to the total mass measured for J041757-NE, suggesting that J041757-NE might indeed be a pre-substellar core. Interestingly, its total mass and radius follow the correlation found between mass and radius for protostellar/pre-stellar cores (e.g. Kauffmann et al. 2008; Sadavoy et al. 2010; Schmalzl et al. 2010). Also, the fact that this pre-substellar core candidate is found in B211, the portion of L1495 with smaller number of young stellar objects (e.g. Schmalzl et al. 2010) supports the idea suggested by Schmalzl et al. (2010) that this region is at the very beginning of the star (or BD) formation process.

The number of pre-substellar core candidates in the literature is very scarce. A pioneer study searching for pre-substellar cores and proto-BDs was carried out by Pound & Blitz (1993, 1995), who found about four cores with substellar masses in the Taurus and Ophiucus clouds. However, the cores were at the detection limit and all but one were not gravitationally bound. In addition, the authors indicate that the candidate objects should be re-observed deeply to rule out the possibility that the cores extend considerably, making them massive enough to overcome the substellar domain. More recent studies in Gould Belt clouds report samples of starless and pre-stellar cores, with some of them being substellar, but lack the kinematical information required to assess if they are gravitationally bound (e.g. Ward-Thompson et al. 2007; Davis et al. 2010; Sadavoy et al. 2010). Thus, J041757-NE is an excellent pre-substellar core candidate, and high angular resolution millimetre observations would be extremely useful to study possible infall motions within the core, which would definitely indicate that the object is on the way to form a BD.

6 CONCLUSIONS

We present the results of a multiwavelength search for pre-substellar cores and proto-BD candidates in the B213-L1495 clouds of Taurus. The studied sample of 12 candidates, selected from *Spitzer*/IRAC photometry of the region (Barrado et al. 2009), was observed at 1.2 mm with the IRAM 30 m/MAMBO-II bolometer. The two most promising objects, J042118 and J041757, were subsequently observed at $350 \mu\text{m}$ with the CSO. Additional observations were performed towards J041757, to further constrain its nature. Our main conclusions can be summarized as follows.

(i) None of the 12 *Spitzer*/IRAC selected sources observed with IRAM 30 m/MAMBO-II were detected at a level $>4\sigma$, and two sources, J042118 and J041757, had fluxes at the 2σ level. We set upper limits to the envelope masses of $\lesssim 3 M_{\text{Jup}}$.

(ii) CSO $350 \mu\text{m}$ observations towards J042118 reveal one compact submillimetre source associated with the selected *Spitzer*/IRAC source, with a size of <10 arcsec ~ 1400 au (at the Taurus distance) and a mass of $\sim 1 M_{\text{Jup}}$. A preliminary analysis of the SED indicates a luminosity of about $0.002 L_{\odot}$, and a bolometric temperature around 140 K, typical of Class I/II young stellar objects.

(iii) CSO 350 μm observations towards J041757 reveal one extended source associated with the selected *Spitzer*/IRAC source, of about ~ 16 arcsec or ~ 2000 au size, and a mass of $\sim 5 M_{\text{Jup}}$. Additionally, we detect one slightly resolved source at 3.6 and 6 cm exactly at the position of the IR counterpart ‘B’ of J041757, called J041757-B, with a flat spectral index. J041757-B is found at the centre of a stream of dust and gas, extending from the B211 main filament to the northeast, mainly detected with $^{13}\text{CO}(1-0)$ at a velocity of $\sim 7 \text{ km s}^{-1}$. $\text{N}_2\text{H}^+(1-0)$ emission is detected at the same velocity at 5σ towards J041757-B. All this seems to indicate that J041757-B is probably a proto-BD in the Class 0/I stage associated with a radio jet and an extended envelope, whose properties seem to be a scaled-down version of low-mass protostars.

(iv) We discovered a partially extended submillimetre source about 40 arcsec to the northeast of J041757-B, of 20 arcsec ~ 3000 au size and total mass of $\sim 100 M_{\text{Jup}}$, called J041757-NE, and with no centimetre emission neither optical/IR (up to 24 μm) emission associated. This indicates that J041757-NE seems to be starless. J041757-NE is well detected in ^{13}CO and N_2H^+ , and is also found at $\sim 7 \text{ km s}^{-1}$. With the measured line width of N_2H^+ , and its measured size and mass, we conclude that J041757-NE is likely gravitationally bound and thus is a good pre-substellar core candidate.

Overall, this work shows the power of submillimetre instruments to trace extremely low masses and thus to reveal substellar objects. We found evidence of two proto-BD candidates and one pre-substellar core candidate which have scaled-down properties of low-mass young stellar objects and which are embedded in large-scale structures, all this favouring the in situ scenario for BD formation in this portion of the Taurus complex.

ACKNOWLEDGMENTS

We thank the referee Thomas Henning for key comments and suggestions. AP is grateful to Attila Kovacs, Stephane Leon, Robert Zylka and Jean-François Lestrade for useful suggestions and help regarding the reduction of the CSO and IRAM 30m data and to Álvaro Hacar, Mario Tafalla, Benjamín Montesinos, Rosario López, María Rosa Zapatero-Osorio, Almudena Alonso, Giovanni Miniutti, Tommy Wiklund and Violeta González for insightful discussions. AP and NH are grateful to Catarina Alves de Oliveira, Florian Rodler, Markus Schmalzl and Darren Dowel for kindly providing data for comparison and DB is grateful to Manuel Güdel for checking that the targets of this work have not been observed with *Chandra* and *XMM*. We thank Calar Alto Observatory and IRAM 30m telescope for allocation of director’s discretionary time to this programme. AP and IdG-M are supported by the Spanish MICINN grant AYA2008-06189-C03 (co-funded with FEDER funds) and AP is also supported by a JAE-Doc CSIC fellowship co-funded with the European Social Fund under the programme ‘Junta para la Ampliación de Estudios’. This research has been partially funded by Spanish MICINN under the Consolider-CSD2006-00070, AYA2010-21161-C02-02 and PRICIT-S2009/ESP-1496 grants and by the Marie Curie Actions of the European Commission (FP7-COFUND). This publication makes use of VOSA, developed under the Spanish Virtual Observatory project supported from the Spanish MICINN through grant AyA2008-02156; of the SIMBAD data base, operated at CDS, Strasbourg, France; of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute

of Technology, funded by the National Aeronautics and Space Administration. Some of the data reported in this paper were obtained as part of the United Kingdom Infrared Telescope (UKIRT) Service Programme. The UKIRT is operated by the Joint Astronomy Centre on behalf of the Science and Technology Facilities Council of the UK Social Fund.

REFERENCES

- Allard F., 2009, *A&A*, 500, 93
 Allard F., Hauschildt P. H., Alexander D. R., Starrfield S., 1997, *ARA&A*, 35, 137
 Allard F., Guillot T., Ludwig H.-G., Hauschildt P. H., Schweitzer A., Alexander D. R., Ferguson J. W., 2003, in Martín E., ed., *Proc. IAU Symp.* 211, Brown Dwarfs. Astron. Soc. Pac., San Francisco, p. 325
 Allard F., Allard N. F., Homeier D., Kielkopf J., McCaughrean M. J., Spiegelman F., 2007, *A&A*, 474, L21
 Allen L. E. et al., 2004, *ApJS*, 154, 363
 Anderson S. F. et al., 2003, *AJ*, 126, 2209
 André P., Deeney B. D., Phillips R. B., Lestrade J.-F., 1992, *ApJ*, 401, 667
 André P., Ward-Thompson D., Barsony M., 1993, *ApJ*, 406, 122
 André P., Motte F., Bacmann A., 1999, *ApJ*, 513, L57
 André P. et al., 2010, *A&A*, 518, L102
 Anglada G., 1995, *Rev. Mex. Astron. Astrofis. Conf. Ser.*, 1, 67
 Anglada G., Villuendas E., Estalella R., Beltrán M. T., Rodríguez L. F., Torrelles J. M., Curiel S., 1998, *AJ*, 116, 2953
 Apai D., Tóth L. V., Henning T., Vavrek R., Kovács Z., Lemke D., 2005, *A&A*, 433, L33
 Avila R., Rodríguez L. F., Curiel S., 2001, *Rev. Mex. Astron. Astrofis.*, 37, 201
 Baraffe I., Chabrier G., Allard F., Hauschildt P. H., 1997, *A&A*, 327, 1054
 Baraffe I., Chabrier G., Allard F., Hauschildt P., 1998, *VizieR Online Data Catalog*, 333, 70403
 Barrado y Navascués D., Martín E. L., 2003, *AJ*, 126, 2997
 Barrado y Navascués D., Mohanty S., Jayawardhana R., 2004, *ApJ*, 604, 284
 Barrado y Navascués D. et al., 2007, *ApJ*, 664, 481
 Barrado D. et al., 2009, *A&A*, 508, 859
 Basu S., Vorobyov E. I., 2012, preprint (astro-ph/1203.0274)
 Bayo A., Rodrigo C., Barrado y Navascués D., Solano E., Gutiérrez R., Morales-Calderón M., Allard F., 2008, *A&A*, 492, 277
 Beltrán M. T., Estalella R., Anglada G., Rodríguez L. F., Torrelles J. M., 2001, *AJ*, 121, 1556
 Bourke T. L., Crapsi A., Myers P. C., Evans, N. J., II, Wilner D. J., Huard T. L., Jørgensen J. K., Young C. H., 2005, *ApJ*, 633, L129
 Bouy H. et al., 2009, *A&A*, 504, 199
 Cambresy L., Epchtein N., Copet E., de Batz B., Kimeswenger S., Le Bertre T., Rouan D., Tiphene D., 1997, *A&A*, 324, 5
 Carter M. et al., 2012, *A&A*, 538, 89
 Castelli F., Gratton R. G., Kurucz R. L., 1997, *A&A*, 318, 841
 Chabrier G., Baraffe I., Allard F., Hauschildt P., 2000, *ApJ*, 542, 464
 Chen H., Myers P. C., Ladd E. F., Wood D. O. S., 1995, *ApJ*, 445, 377
 Davis C. J. et al., 2010, *MNRAS*, 405, 759
 di Francesco J., Evans N. J., II, Caselli P., Myers P. C., Shirley Y., Aikawa Y., Tafalla M., 2007, in Reipurth B., Jewitt D., Keil K., eds, *Protostars and Planets V*. Univ. Arizona Press, Tucson, p. 17
 di Francesco J., Johnstone D., Kirk H. M., MacKenzie T., Ledwosinska E., 2008, *ApJS*, 175, 277
 Dunham M. M., Crapsi A., Evans N. J., II, Bourke T. L., Huard T. L., Myers P. C., Kauffmann J., 2008, *ApJS*, 179, 249
 Dunham M. M., Evans N. J., III, Bourke T. L., Myers P. C., Huard T. L., Stutz A. M., 2010, *ApJ*, 721, 995
 Elvis M. et al., 1994, *ApJS*, 95, 1
 Fernández M., Comerón F., 2001, *A&A*, 380, 264
 Fitzpatrick E. L., 1999, *PASP*, 111, 63
 Forbrich J., Massi M., Ros E., Brunthaler A., Menten K. M., 2007, *A&A*, 469, 985

- Frerking M. A., Langer W. D., Wilson R. W., 1982, *ApJ*, 262, 590
- Froebrich D., 2005, *ApJS*, 156, 169
- Gandhi P., Horst H., Smette A., Hšnig S., Comastri A., Gilli R., Vignali C., Duschl W., 2009, *A&A*, 502, 457
- Giroletti M., Panessa F., 2009, *ApJ*, 706, L260
- Goldsmith P. F., Heyer M., Narayanan G., Snell R., Li D., Brunt C., 2008, *ApJ*, 680, 428
- Goulding A. D., Alexander D. M., 2009, *MNRAS*, 398, 1165
- Greaves J. S., Holland W. S., Pound M. W., 2003, *MNRAS*, 346, 441
- Güdel M. et al., 2007, *A&A*, 468, 353
- Harvey P. M. et al., 2006, *ApJ*, 644, 307
- Harvey P. M., Jaffe D. T., Allers K., Liu M., 2010, *ApJ*, 720, 1374
- Harvey P. M. et al., 2012, *ApJ*, 744, L1
- Hauck B., Mermilliod M., 1998, *A&AS*, 129, 431
- Hauschildt P. H., Allard F., Baron E., 1999, *ApJ*, 512, 377
- Hewett P. C., Warren S. J., Leggett S. K., Hodgkin S. T., 2006, *MNRAS*, 367, 454
- Heyer M. H., Snell R. L., Goldsmith P. F., Myers P. C., 1987, *ApJ*, 321, 370
- Ibar E., Ivion R. J., Biggs A. D., Lal D. V., Best P. N., Green D. A., 2009, *MNRAS*, 397, 281
- Indebetouw R. et al., 2005, *ApJ*, 619, 931
- Jayawardhana R., Ivanov V. D., 2006, *ApJ*, 647, L167
- Jayawardhana R., Coffey J., Scholz A., Brandeker A., van Kerkwijk M. H., 2006, *ApJ*, 648, 1206
- Joergens V., Pohl A., Sicilia-Aguilar A., Henning Th., 2012, *A&A*, preprint (arXiv:1205.3166)
- Jørgensen J. K. et al., 2006, *ApJ*, 645, 1246
- Kauffmann J., Bertoldi F., Evans N. J., II, the C2D Collaboration, 2005, *Astron. Nachrichten*, 326, 878
- Kauffmann J., Bertoldi F., Bourke T. L., Evans N. J., II, Lee C. W., 2008, *A&A*, 487, 993
- Kauffmann J., Bertoldi F., Bourke T. L., Myers P. C., Lee C. W., Huard T. L., 2011, *MNRAS*, 416, 2341
- Kellermann K. I., Sramek R., Schmidt M., Shaffer D. B., Green R., 1989, *AJ*, 98, 1195
- Kenyon S. J., Hartmann L., 1995, *ApJS*, 101, 117
- Klein R., Apai D., Pascucci I., Henning T., Waters L. B. F. M., 2003, *ApJ*, 593, L57
- Kovács A., 2008, *Proc. SPIE*, 7020, 45
- Lee C. W. et al., 2009, *ApJ*, 693, 1290
- Lee J.-E., 2007, *J. Korean Astron. Soc.*, 40, 83
- Liebert J., Cutri R. M., Nelson B., Kirkpatrick J. D., Gizis J. E., Reid I. N., 2000, *PASP*, 112, 1315
- Lucas P. W. et al., 2008, *MNRAS*, 391, 136
- Luhman K. L., Mamajek E. E., 2010, *ApJ*, 716, L120
- Luhman K. L., Liebert J., Rieke G. H., 1997, *ApJ*, 489, L165
- Luhman K. L., Adame L., D'Alessio P., Calvet N., Hartmann L., Megeath S. T., Fazio G. G., 2005, *ApJ*, 635, L93
- Luhman K. L. et al., 2007, *ApJ*, 666, 1219
- Luhman K. L., Hernández J., Downes J. J., Hartmann L., Briceño C., 2008, *ApJ*, 688, 362
- Luhman K. L., Allen P. R., Espaillat C., Hartmann L., Calvet N., 2010, *ApJS*, 186, 111
- Mermilliod J.-C., Mermilliod M., 1994, *Catalogue of Mean UBV Data on Stars*, VI. Springer-Verlag, Berlin, p. 1387
- Mizuno A., Onishi T., Yonekura Y., Nagahama T., Ogawa H., Fukui Y., 1995, *ApJ*, 445, L161
- Mohanty S., Jayawardhana R., Basri G., 2005, *ApJ*, 626, 498
- Monin J.-L. et al., 2010, *A&A*, 515, A91
- Narayanan G., Heyer M. H., Brunt C., Goldsmith P. F., Snell R., Li D., 2008, *ApJS*, 177, 341
- Natta A., Testi L., Muzerolle J., Randich S., Comerón F., Persi P., 2004, *A&A*, 424, 603
- Ossenkopf V., Henning T., 1994, *A&A*, 291, 943
- Padoan P., Nordlund Å., 2004, *ApJ*, 617, 559
- Pascucci I., Apai D., Henning T., Dullemond C. P., 2003, *ApJ*, 590, L111
- Pech G., Loinard L., Chandler C. J., Rodríguez L. F., D'Alessio P., Brogan C. L., Wiener D. J., Ho P. T. P., 2010, *ApJ*, 712, 1403
- Phan-Bao N. et al., 2008, *ApJ*, 689, L141
- Pirogov L., Zinchenko I., Caselli P., Johansson L. E. B., Myers P. C., 2003, *A&A*, 405, 639
- Pound M. W., Blitz L., 1993, *ApJ*, 418, 328
- Pound M. W., Blitz L., 1995, *ApJ*, 444, 270
- Quanz S. P., Goldman B., Henning T., Brandner W., Burrows A., Hofstetter L. W., 2010, *ApJ*, 708, 770
- Randall K., Hopkins A., Norris R., Zinn P.-C., Middelberg E., Mao M., Sharp R., 2012, *MNRAS*, 421, 1644
- Reipurth B., Clarke C., 2001, *AJ*, 122, 432
- Riaz B., Lodato G., Stamatellos D., Gizis J. E., 2012a, *MNRAS*, 422, L6
- Riaz B., Lodieu N., Goodwin S., Stamatellos D., Thompson M., 2012b, *MNRAS*, 420, 2497
- Rice W. K. M., Armitage P. J., Bonnell I. A., Bate M. R., Jeffers S. V., Vine S. G., 2003, *MNRAS*, 346, L36
- Roussel H., 2012, preprint (arXiv:1205.2576)
- Sadavoy S. I. et al., 2010, *ApJ*, 710, 1247
- Scaife A. M. M. et al., 2011, *MNRAS*, 415, 893
- Schmalzl M. et al., 2010, *ApJ*, 725, 1327
- Scholz A., Jayawardhana R., Wood K., 2006, *ApJ*, 645, 1498
- Scholz A., Jayawardhana R., Wood K., Lafrenière D., Schreyer K., Doyon R., 2008, *ApJ*, 681, L29
- Shang Z. et al., 2011, *ApJS*, 196, 2
- Shirley Y. L., Claussen M. J., Bourke T. L., Young C. H., Blake G. A., 2007, *ApJ*, 667, 329
- Skrutskie M. F. et al., 2006, *AJ*, 131, 1163
- Stamatellos D., Whitworth A. P., 2009, *MNRAS*, 392, 413
- Tafalla M., Myers P. C., Caselli P., Walmsley C. M., 2004, *A&A*, 416, 191
- Umbreit S., Burkert A., Henning T., Mikkola S., Spurzem R., 2005, *ApJ*, 623, 940
- Voges W. et al., 1999, *A&A*, 349, 389
- Ward-Thompson D. et al., 2007, *PASP*, 119, 855
- Ward-Thompson D., Kirk J. M., Greaves J. S., André P., 2011, *MNRAS*, 415, 2812
- White R. J., Ghez A. M., Reid I. N., Schultz G., 1999, *ApJ*, 520, 811
- Whitworth A. P., Zinnecker H., 2004, *A&A*, 427, 299
- Young C. H. et al., 2004, *ApJS*, 154, 396

APPENDIX A: PHOTOMETRY FOR J042118 AND J041757-B

In this appendix, we give the photometry for J042118 and J041757-B, from visible wavelengths up to the upper limit set at 1.2 μm with MAMBO-II. To compile the photometry, we used the VOSA (Bayo et al. 2008). See Hewett et al. (2006) and Lucas et al. (2008)

Table A1. Photometry for J042118.

λ (μm)	S_V (mJy)	σ (mJy)	Instrument
0.89	<0.0142	–	SDSS-z
1.03	<0.0130	–	UKIDSS-Y
1.25	0.0165	0.0003	UKIDSS-J
2.20	0.0453	0.0009	UKIDSS-K
3.4	0.196	0.010	WISE
3.6	0.249	0.004	<i>Spitzer</i> /IRAC
4.5	0.439	0.006	<i>Spitzer</i> /IRAC
4.6	0.435	0.023	WISE
5.8	0.674	0.010	<i>Spitzer</i> /IRAC
8.0	1.093	0.016	<i>Spitzer</i> /IRAC
12	1.13	0.33	WISE
24	2.73	0.07	<i>Spitzer</i> /MIPS
70	<33.6	<4 σ	<i>Spitzer</i> /MIPS
350	46	9	CSO
1200	<4.0	<4 σ	IRAM 30 m

Table A2. Photometry for J041757-B.

λ (μm)	S_{ν} (mJy)	σ (mJy)	Instrument
0.75	0.0034	0.0001	CFHT- <i>i</i>
0.90	0.0077	0.0002	CFHT- <i>z</i>
1.03	<0.0130	–	UKIDSS- <i>Y</i>
1.25	0.0360	0.0006	CAHA- <i>J</i>
1.65	0.0680	0.0012	CAHA- <i>H</i>
2.17	0.127	0.002	CAHA- <i>K</i>
3.6	0.295	0.008	<i>Spitzer</i> /IRAC
4.5	0.441	0.009	<i>Spitzer</i> /IRAC
5.8	0.491	0.010	<i>Spitzer</i> /IRAC
8.0	0.736	0.011	<i>Spitzer</i> /IRAC
12	1.79	0.15	WISE

Table A2 – continued

λ (μm)	S_{ν} (mJy)	σ (mJy)	Instrument
22	6.5	1.0	WISE
24	7.5	1.0	<i>Spitzer</i> /MIPS
70	<33.6	<4 σ	<i>Spitzer</i> /MIPS
350	165	9	CSO
870	<80	<4 σ	APEX
1200	<2.9	<4 σ	IRAM 30 m
36 000	0.19	0.11	VLA
60 000	0.20	0.05	VLA

for a review of the UKIDSS. For the case of J041757-B, we again measured the flux at 24 μm to take into account extended emission (neglected in our previous measurement given in Barrado et al. 2009).

This paper has been typeset from a \TeX/L\AA\TeX file prepared by the author.