



EUROPEAN
HEMATOLOGY
ASSOCIATION


Journal of the European Hematology Association

Chronic lymphocytic leukemia cells in lymph nodes show frequent NOTCH1 activation

by Arantza Onaindia, Sagrario Gómez, Miguel Piris-Villaespesa, Carolina Martínez-Laperch, Laura Cereceda, Santiago Montes-Moreno, Ana Batlle, Sonia González de Villambrosia, Marina Pollán, Paloma Martín-Acosta, Julia González-Rincón, Javier Menarguez, Javier Alvés, Socorro M. Rodríguez-Pinilla, Juan F. García, Manuela Mollejo, Máximo Fraga, José A. García-Marco, Miguel A. Piris, and Margarita Sanchez-Beato

Haematologica 2014 [Epub ahead of print]

*Citation: Onaindia A, Gómez S, Piris-Villaespesa M, Martínez-Laperch C, Cereceda L, Montes-Moreno S, Batlle A, de Villambrosia SG, Pollán M, Martín-Acosta P, González-Rincón J, Menarguez J, Alvés J, Rodríguez-Pinilla SM, García JF, Mollejo M, Fraga M, García-Marco JA, Piris MA, and Sanchez-Beato M. Chronic lymphocytic leukemia cells in lymph nodes show frequent NOTCH1 activation. Haematologica. 2014; 99:xxx
doi:10.3324/haematol.2014.117705*

Publisher's Disclaimer.0

E-publishing ahead of print is increasingly important for the rapid dissemination of science. Haematologica is, therefore, E-publishing PDF files of an early version of manuscripts that have completed a regular peer review and have been accepted for publication. E-publishing of this PDF file has been approved by the authors. After having E-published Ahead of Print, manuscripts will then undergo technical and English editing, typesetting, proof correction and be presented for the authors' final approval; the final version of the manuscript will then appear in print on a regular issue of the journal. All legal disclaimers that apply to the journal also pertain to this production process.

Chronic lymphocytic leukemia cells in lymph nodes show frequent NOTCH1 activation

Arantza Onaindia¹, Sagrario Gómez², Miguel Piris-Villaespesa³, Carolina Martínez-Laperche⁴, Laura Cereceda¹, Santiago Montes-Moreno¹, Ana Batlle¹, Sonia González de Villambrosia¹, Marina Pollán^{5, 6}, Paloma Martín-Acosta⁷, Julia González-Rincón², Javier Menarguez⁴, Javier Alvés⁸, Socorro M. Rodríguez-Pinilla⁹, Juan F. García¹⁰, Manuela Mollejo¹¹, Máximo Fraga¹², José A. García-Marco³, Miguel A. Piris¹ and Margarita Sánchez-Beato^{2,#}

¹Pathology Department, Hospital U. Marqués de Valdecilla-Santander; ²Group of Research in Lymphomas (Medical Oncology Department), Oncohematology Area, Instituto Investigación Puerta de Hierro-Majadahonda (IDIPHIM), Madrid; ³Hematology Department, Hospital U. Puerta de Hierro-Majadahonda, Madrid; ⁴Pathology Department, Hospital U. Gregorio Marañón, Madrid; ⁵Cancer Epidemiology Unit, National Center for Epidemiology, Instituto de Salud Carlos III, Madrid; ⁶Cancer Epidemiology Research Group, Oncohematology Area, Instituto Investigación Puerta de Hierro-Majadahonda (IDIPHIM), Madrid; ⁷Group of Research in Molecular Pathology of Cancer (Pathology Department), Oncohematology Area, Instituto Investigación Puerta de Hierro-Majadahonda (IDIPHIM), Madrid; ⁸Pathology Department, Hospital U. La Paz, Madrid; ⁹Pathology Department, Fundación Jiménez Díaz, Madrid; ¹⁰Pathology Department, MD Anderson Cancer Center, Madrid; ¹¹Pathology Department, Hospital Virgen de la Salud, Toledo; ¹²Pathology Department, Hospital U. Santiago de Compostela, La Coruña. SPAIN.

Running heads: NOTCH1 activation in CLL

Acknowledgments: We are indebted to the patients who contributed to this study and to the hospitals who supplied the samples. We acknowledge the Biobanks of the CNIO (RD09/0076/00113), IDIVAL-HUMV (RD09/0076/00076), HU 12 de Octubre (RD09/0076/00118) and HGU Gregorio Marañón for their help in collecting the samples. This work was supported by grants from the Ministerio de Economía y Competitividad (MINECO) (SAF2013-47416-R) Instituto de Salud Carlos III (ISCIII)- FEDER – MINECO- AES (CP11/00018, PI10/00621, RD012/0036/0060), and Asociación Española contra el Cáncer (AECC). MS-B is supported by a Miguel Servet contract from ISCIII-FEDER (CP11/00018). Salary support to SG is provided by CP11/00018, from ISCIII-FEDER. JG-R is supported by a predoctoral grant from the Fundación Investigación Puerta de Hierro.

Correspondence

Dr. Margarita Sanchez-Beato
IIS Puerta de Hierro (IDIPHIM)
Calle Joaquín Rodrigo 2,
Majadahonda-Madrid. ES-28222, Spain
E-mail: msbeato@idiphim.org

Letter

Chronic lymphocytic leukemia (CLL) is the most common adult leukemia in the western world. Pathogenic mechanisms involve multiple external events (such as microenvironmental and antigenic stimuli) and internal events (genetic and epigenetic alterations) that are associated with the transformation, progression and evolution of CLL. CLL is characterized by an accumulation of mature B-cells in peripheral blood, bone marrow and lymphoid tissues. Extracellular stimuli play an important role in the development and maintenance of neoplastic cells. B-CLL cells proliferate and activate pathogenic signaling pathways in anatomical structures known as proliferation centers, which are usually more conspicuous in involved lymph nodes (1). Its clinical course is quite heterogeneous, whereby some patients progress rapidly and have short survival, whereas others have a more stable clinical course that may not need treatment for years.

Several clinical and biological prognostic factors for CLL have been identified, such as the Rai and Binet clinical staging systems, specific cytogenetic alterations, mutational status of immunoglobulin (IgHv) genes, TP53 mutations and the expression levels of CD38 and ZAP70. More recently, massive sequencing data have revealed new genetic alterations, some of which have a significant impact on the clinical course. Of these, mutations in the *NOTCH1* and *SF3B1* genes, which are associated with an adverse clinical outcome, are emerging as the most frequent mutations in CLL (2).

The *NOTCH1* gene codes for a trans-membrane receptor that regulates critical cellular functions such as proliferation, apoptosis and differentiation, and is associated with tumorigenesis. NOTCH1 signaling is initiated when a ligand from the Jagged or Delta families binds the receptor and induces a cascade of proteolytic processes that results in the release and nuclear translocation of the NOTCH1 intracellular domain (NICD). This, in conjunction with the transcription factor CBF1/RBP-Jk, leads to the activation or repression of its target genes (3, 4).

Although a considerable number of studies have addressed the consequences of *NOTCH1* and *SF3B1* mutations in CLL (2), few have so far analyzed the frequency and biological impact of these mutations in lymph nodes (LN) affected by CLL (InCLL). There

is a growing interest in therapeutic targeting of NOTCH1 activation signaling in tumors with agonists such as γ -secretase inhibitors (5). It has also been proposed that mutation-independent NOCH1-activation contributes to tumor cell growth and survival (6, 7).

Therefore, we have analyzed the frequency of *NOTCH1* and *SF3B1* mutations and the functional status of the NOTCH1 pathway through the expression analysis of NOTCH1-induced targets in a series of samples of InCLL.

The first step was the analysis of *NOTCH1* and *SF3B1* mutations in the LN infiltrated with CLL. The series included 155 LN samples from 147 patients, who were biopsied at diagnosis or during the course of the disease (Supplemental Materials and Methods). The characteristics of the series are summarized in **Table 1**. *NOTCH1* and *SF3B1* mutations were successfully analyzed in 140 and 142 samples by qPCR respectively, finding that 32/140 (22.8%) were positive for the p.P2515fs*4 *NOTCH1* mutation and 14/142 (9.8%) for the p.K700E *SF3B1* mutation. As this *SF3B1* mutation constitutes only around 50-60% of all *SF3B1* mutations identified in CLL, we also study exons 14 and 16, in which most of the other mutations reside. Although capillary sequencing is a less sensitive method than qPCR, we were still able to identify four additional *SF3B1* mutations in exon 14 and none in exon 16. Therefore, 18/142 (12.7%) *SF3B1* gene mutations were detected. Three out of 132 samples showed a mutation in both genes (**Table 2**). When comparing the rate of mutations in both genes in samples taken at the time of diagnosis or progression, no significant differences were found (**Table 2**). Samples at diagnosis and during progression were available for eight patients. *NOTCH1* status did not change in any of the patients (2 mutated and 6 non-mutated), while one acquired an *SF3B1* mutation during progression.

We found that although the percentage of cases with mutated *SF3B1* at the time of diagnosis (11.7%) was similar to that of other series analyzed in peripheral blood (PB) lymphocytes (2), the *NOTCH1* mutation rate we found in InCLL (24.5% at diagnosis) seems to be higher than in other studies performed in PB samples, in which the percentages of mutations range from 8% to 31% (8, 9) depending on the series and its characteristics. The lowest values (less than 15%) were obtained from the analysis

of samples taken at diagnosis and the highest were found in patients transformed to Richter syndrome (8) or who showed refractoriness to treatment (10). It is also higher than the frequency found in a different series of CLL PB taken at the time of diagnosis analyzed in our laboratory using the same method in which we found 12% of cases with p.P2515fs*4 NOTCH1 mutation (data not shown).

We only analyzed the p.P2515fs*4 mutation, which accounts for around 80% of all *NOTCH1* mutations reported in CLL (from 66% to 98% (11)) and therefore the real percentage of mutated cases in our InCLL might be even higher. This could be due to the greater aggressiveness of InCLL and/or to the high sensitivity of the qPCR technique used for mutation detection.

We next analyzed the association of *NOTCH1* and *SF3B1* mutations in samples taken at diagnosis with patient, clinical, biological and cytogenetic characteristics. We found no significant associations between the presence of *NOTCH1* or *SF3B1* mutations and any biological and clinical characteristics, or with overall survival and time to treatment (**Table S1** in the Supplementary Information).

The expression of NOTCH1 itself together with NOTCH1 targets, including NFATc1, p52, p50, C-MYC, MUM1, XBP1s, LEF and HES1 (3, 4, 12-14), and Ki67, were analyzed by immunohistochemistry in the 155 CLL paraffin samples and in six samples of reactive lymphoid tissue included in the Tissue Micro Arrays (TMA) (Supplemental Materials and Methods).

Lymphocytes in reactive LN exhibited mainly cytoplasmic expression of NOTCH1, NFAT, p52 and p50. MUM1 expression was restricted to plasma cells and scattered cells within the germinal centers, XBP1 only recognized plasma cells, while MYC, LEF1 and HES1 expression were negative in normal B-cells.

NOTCH1 nuclear accumulation was analyzed with a rabbit-mAb recognizing the NICD1 epitope (15), a surrogate of NOTCH1 activation, and was detected in both 23/29 (79.3%) of *NOTCH1*-mutated case sand in 82/91 (90%) of *NOTCH1*-wild-type cases. This means that in a large proportion of samples (105/120, 87.5%), NOTCH1 could be detected in the nuclear compartment of the tumoral B-cells (mainly in the proliferation

centers) (**Table S2**), indicating that the WNT-NOCTH1 pathway is frequently activated in InCLL, irrespective of the presence of the NOTCH1 mutation, as has been previously reported by others (15).

The *NOTCH1* mutation was associated with nuclear expression of NFAT ($\chi^2 = 8.081$, $p = 0.005$), NF-kB p52 subunit ($\chi^2 = 5.841$, $p = 0.016$) and CMYC ($\chi^2 = 4.077$, $p = 0.043$), which was selectively expressed in the proliferation center B-cells (**Table S2** and **Figure 1**). The intensity and selectiveness of the expression of CMYC by neoplastic B-cells within proliferation centers were unexpected, being similar to those found in tumoral cells in Burkitt lymphoma samples, although they were restricted to the proliferation centers in our study (Figure 1). Other NOTCH1 downstream targets (such as HES1, LEF1 and XBP1) were also diffusely expressed by neoplastic B-cells, and were detected in mutated and non-mutated samples, further supporting the hypothesis that the WNT-NOCTH1 pathway is frequently activated in InCLL, irrespective of the presence of the *NOTCH1* mutation.

However, NOTCH1 nuclear expression was not significantly associated with the detection of any of the markers analyzed (**Table S3**), their expression being found in samples with or without nuclear NOTCH1 expression.

The NOTCH1 pathway is activated when ligands of the JAG or Delta families bind to their receptors, leading to the release and nuclear translocation of NOTCH1. These ligands are known to be constitutively expressed in B-CLL cells (7). Therefore, we decided to analyze JAG1 expression in these samples. We found that 31/94 (31.9%) expressed the protein and that its detection was associated with the expression of some NOTCH1 targets such as LEF1 ($\chi^2 = 7.095$, $p = 0.008$), MYC ($\chi^2 = 5.158$, $p = 0.023$) and MUM1 ($\chi^2 = 5.369$, $p = 0.021$), although not with NOTCH1 expression itself (**Table S4**) as NOTCH1 could be activated by mutation independently of JAG1 presence. JAG1 was expressed in the B-CLL cells in the proliferative centers of wild-type and mutated cases in dots in the Golgi apparatus.

These results reflect the complexity of the NOTCH1 pathway in CLL as it could be constitutively activated by different mechanisms including gene mutation, ligand

activation or, even, by the participation of other members such as NOTCH2; as has been also previously reported (7).

In conclusion, we have found that the mutational frequency of *NOTCH1*, but not of *SF3B1*, is higher in LN with CLL cell involvement than in other series analyzing CLL PB samples.

Another major finding of this study is that NOTCH1 pathway is frequently activated in InCLL, independently of *NOTCH1* gene mutational status, thus suggesting the relevance of NOTCH1 pathway in the survival of CLL cells, and the existence of previously reported additional mechanisms of NOTCH1 activation (7). These include the stronger expression of NOTCH1 ligands, which suggests the existence of autocrine/paracrine mechanisms that could lead to NOTCH1 signaling activation in addition to *NOTCH1* gene mutation.

REFERENCES

1. Herreros B, Rodriguez-Pinilla SM, Pajares R, et al. Proliferation centers in chronic lymphocytic leukemia: the niche where NF-kappaB activation takes place. *Leukemia*. 2010;24(4):872-876.
2. Villamor N, Lopez-Guillermo A, Lopez-Otin C, Campo E. Next-generation sequencing in chronic lymphocytic leukemia. *Semin Hematol*. 2013 Oct;50(4):286-295.
3. Vilimas T, Mascarenhas J, Palomero T, et al. Targeting the NF-kappaB signaling pathway in Notch1-induced T-cell leukemia. *Nat Med*. 2007;13(1):70-77.
4. Guiu J, Shimizu R, D'Altri T, et al. Hes repressors are essential regulators of hematopoietic stem cell development downstream of Notch signaling. *J Exp Med*. 2012;210(1):71-84.
5. Lopez-Guerra M, Xargay-Torrent S, Rosich L, et al. The gamma-secretase inhibitor PF-03084014 combined with fludarabine antagonizes migration, invasion and angiogenesis in NOTCH1-mutated CLL cells. *Leukemia*. 2014 Apr 30;28(4):Apr 30. doi: 10.1038/leu.2014.2143.
6. Arruga F, Gizdic B, Serra S, et al. Functional impact of NOTCH1 mutations in chronic lymphocytic leukemia. *Leukemia*. 2014;28(5):1060-1070.
7. Rosati E, Sabatini R, Rampino G, et al. Constitutively activated Notch signaling is involved in survival and apoptosis resistance of B-CLL cells. *Blood*. 2009;113(4):856-865.
8. Fabbri G, Khiabanian H, Holmes AB, et al. Genetic lesions associated with chronic lymphocytic leukemia transformation to Richter syndrome. *J Exp Med*. 2013;210(11):2273-2288.
9. Rossi D, Rasi S, Fabbri G, et al. Mutations of NOTCH1 are an independent predictor of survival in chronic lymphocytic leukemia. *Blood*. 2011;119(2):521-529.
10. Schnaiter A, Paschka P, Rossi M, et al. NOTCH1, SF3B1, and TP53 mutations in fludarabine-refractory CLL patients treated with alemtuzumab: results from the CLL2H trial of the GCLLSG. *Blood*. 2013;122(7):1266-1270.
11. Jeromin S, Weissmann S, Haferlach C, et al. SF3B1 mutations correlated to cytogenetics and mutations in NOTCH1, FBXW7, MYD88, XPO1 and TP53 in 1160 untreated CLL patients. *Leukemia*. 2014;28(1):108-117.
12. Mammucari C, Tommasi di Vignano A, et al. Integration of Notch 1 and calcineurin/NFAT signaling pathways in keratinocyte growth and differentiation control. *Dev Cell*. 2005 May;8(5):665-76.
13. Palomero T, Lim WK, Odom DT, et al. NOTCH1 directly regulates c-MYC and activates a feed-forward-loop transcriptional network promoting leukemic cell growth. *Proc Natl Acad Sci U S A*. 2006;103(48):18261-18266.
14. Petropoulos K, Arseni N, Schessl C, et al. A novel role for Lef-1, a central transcription mediator of Wnt signaling, in leukemogenesis. *J Exp Med*. 2008;205(3):515-522.
15. Kluk MJ, Ashworth T, Wang H, et al. Gauging NOTCH1 Activation in Cancer Using Immunohistochemistry. *PLoS One*. 2013;8(6):e67306.

TABLES**Table 1.-** Summary of the main clinical and biological characteristics of patients with samples at diagnosis.

Lymph node samples obtained at diagnosis Summary of clinical and biological characteristics	
Median age at diagnosis, years (range)	69.8 (39-89)
Male, n (%)	65/100 (65.0%)
Binet B-C, n (%)	36/72 (82.0%)
Unmutated IgH, n (%)	40/65 (61.5%)
13q del, n (%)	6/38 (15.8%)
Trisomy 12, n (%)	9/39 (23.1%)
11q del, n (%)	6/60 (10.0%)
17p del, n (%)	4/63 (6.3%)
Lymphocytes x10 ⁹ /L, median (range)	7 (1-130)
CD38-positive, n (%)	43/65 (66.2%)
ZAP70-positive, n (%)	22/45 (49.0%)
Treatment, n (%)	67/134 (50.0%)
Median follow-up, mo (range)	28 (0-152)
Death by CLL	47/100 (47.0%)

Table 2.- Summary of samples with mutations in NOTCH1 and/or SF3B1.

	DX (%)	PR (%)	TOTAL (%)
NOTCH1 mut	28/114 (24.5)	4/22 (18.2)	32/140 (22.8)
SF3B1 mut	14/119 (11.7)	3/19 (15.8)	18/142 (12.7)
Any mutation	39/112 (34.8)	8/19 (42.1)	44/133 (33.1)

Mut: mutated; DX: diagnostic; PR: progression.

FIGURE LEGENDS

Figure 1. (A) Negative NOTCH1 expression in a wild-type case. (B) Mild NOTCH1 expression in a mutated case. (C) Intense NOTCH1 expression in a wild-type case. (D) NFAT cytoplasmic expression in a negative case. (E) NFAT nuclear expression in a positive case. (F) p50 nuclear expression in a negative case. (G) p50 nuclear expression in a positive case. (H) p52 cytoplasmic expression in a negative case. (I) p52 nuclear expression in a positive case. (J) myc expression in the proliferation center of a wild-type case. (K) JAG-negative stain in a mutated case (L). JAG expression in dots in the Golgi apparatus of a wild-type case.

