

ORIGINAL ARTICLE

Child hospital admissions associated with influenza virus infection in 6 Spanish cities (2014–2016)☆



Javier Arístegui Fernández^{a,b,*}, Eduardo González Pérez-Yarza^{b,c,d},
María José Mellado Peña^e, Carlos Rodrigo Gonzalo de Liria^{f,g},
Teresa Hernández Sampelayo^h, Juan José García Garcíaⁱ, Jesús Ruiz Contreras^j,
David Moreno Pérez^k, Elisa Garrote Llanos^{a,b}, José Tomás Ramos Amador^l,
Carlos Gustavo Cilla Eguiluz^{d,m}, María Méndez Hernández^{g,n},
en representación del grupo de estudio HOSPIGRIP◊

^a Sección de Infectología Pediátrica, Hospital Universitario Basurto, Bilbao, Vizcaya, Spain

^b Departamento de Pediatría, Universidad del País Vasco, UPV/EHU, Bilbao, Vizcaya, Spain

^c Servicio de Pediatría, Hospital Universitario Donostia, San Sebastián, Guipúzcoa, Spain

^d Centro de Investigación Biomédica en Red, Enfermedades Respiratorias (CIBERES), San Sebastián, Guipúzcoa, Spain

^e Servicio de Pediatría Hospitalaria, Enfermedades Infecciosas y Tropicales, Hospital Universitario La Paz, Madrid, Spain

^f Servicio de Pediatría, Hospital Universitario Vall d'Hebron, Barcelona, Spain

^g Facultad de Medicina, Unidad Docente Germans Trias i Pujol, Universidad Autónoma de Barcelona, Barcelona, Spain

^h Servicio de Pediatría, Hospital General Universitario Gregorio Marañón, Madrid, Spain

ⁱ Servicio de Pediatría, Hospital Sant Joan de Déu, Universitat de Barcelona, Barcelona, Spain

^j Servicio de Pediatría, Hospital Universitario 12 de Octubre, Madrid, Spain

^k Servicio de Pediatría, Hospital Regional Universitario de Málaga, Málaga, Spain

^l Departamento de Salud Pública y Materno-infantil, Universidad Complutense, Hospital Clínico San Carlos, Instituto de Investigación Sanitaria del Hospital Clínico San Carlos, Madrid, Spain

^m Servicio de Microbiología, Hospital Universitario Donostia, San Sebastián, Guipúzcoa, Spain

ⁿ Servicio de Pediatría, Hospital Universitario Germans Trias i Pujol, Badalona, Barcelona, Spain

Received 25 May 2018; accepted 22 June 2018

Available online 21 December 2018

KEYWORDS

Influenza virus;
Hospital admissions;
Children

Abstract

Introduction: There are only a limited number of studies on the impact of influenza in the Spanish child population. The present work intends to increase this knowledge by studying some key aspects, such as the incidence of hospital admissions, clinic variables, comorbidities, and the vaccination status in the hospitalised children.

☆ Please cite this article as: Arístegui Fernández J, González Pérez-Yarza E, Mellado Peña MJ, Rodrigo Gonzalo de Liria C, Hernández Sampelayo T, García García JJ, et al. Hospitalizaciones infantiles asociadas a infección por virus de la gripe en 6 ciudades de España (2014–2016). An Pediatr (Barc). 2019;90:86–93.

* Corresponding author at:

E-mail address: javier.aristegui@ehu.eus (J. Arístegui Fernández).

◊ Appendix A lists all the authors of the article.

Methods: A retrospective, observational study was conducted by reviewing the medical records of children under 15 years and hospitalised due to community acquired influenza confirmed microbiologically, during 2 flu seasons (2014–2015 and 2015–2016). The study was carried out in 10 hospitals of 6 cities, which represent approximately 12% of the Spanish child population.

Results: A total of 907 children were admitted to hospital with main diagnosis of influenza infection (447 <2 years), estimating an average annual rate of hospitalization incidence of 0.51 cases/1000 children (95% CI, 0.48–0.55). Just under half (45%) of the cases had an underlying disease considered a risk factor for severe influenza, and most (74%) had not been vaccinated. The percentage of children with underlying diseases increased with age, from 26% in children <6 months to 74% in children >10 years. Admission to the PICU was required in 10% (92) of the cases, mainly due to acute respiratory failure.

Conclusion: Influenza continues to be an important cause of hospitalization in the Spanish child population. Children <6 months of age and children with underlying diseases make up the majority (>50%) of the cases. Many of the severe forms of childhood influenza that occur today could be avoided if current vaccination guidelines were met.

© 2018 Asociación Española de Pediatría. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Virus de la gripe;
Hospitalizaciones;
Niños

Hospitalizaciones infantiles asociadas a infección por virus de la gripe en 6 ciudades de España (2014–2016)

Resumen

Introducción: La información existente sobre el impacto de la gripe en la población infantil española es escasa. El presente trabajo pretende aumentar este conocimiento estudiando aspectos clave como la incidencia de hospitalización, clínica, comorbilidades y el estado vacunal en los niños hospitalizados.

Métodos: Estudio retrospectivo, observacional, por revisión de historias clínicas, en menores de 15 años hospitalizados por gripe adquirida en la comunidad, confirmada microbiológicamente, durante 2 temporadas gripales (2014-2015 y 2015-2016). El estudio se realizó en 10 hospitales de 6 ciudades, que atienden aproximadamente al 12% de la población infantil española.

Resultados: Fueron hospitalizados 907 niños con diagnóstico principal de gripe (447 <2 años), con una tasa media anual de incidencia de hospitalización de 0,51 casos/1.000 niños (IC del 95% 0,48-0,55). El 45% presentó enfermedades subyacentes consideradas factores de riesgo para gripe grave, y la mayor parte de ellos (74%) no habían sido vacunados. El porcentaje con enfermedades subyacentes aumentó con la edad, desde el 26% en menores de 6 meses al 74% en mayores de 10 años. El 10% de los casos (n = 92) precisaron cuidados intensivos pediátricos por fallo respiratorio agudo.

Conclusión: La gripe es causa importante de hospitalización en la población infantil española. Los menores de 6 meses de edad y los niños con enfermedades subyacentes constituyen una parte mayoritaria (> 50%) de los casos. Una gran parte de las formas graves de gripe en población infantil podrían ser evitada si se cumplieran las indicaciones actuales de vacunación.

© 2018 Asociación Española de Pediatría. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Viral respiratory infections are the most frequent infectious diseases in infants and children.¹ They encompass a broad clinical spectrum, from mild conditions like the common cold to severe diseases that often require inpatient evaluation and treatment, such as infections by respiratory syncytial virus and influenza viruses. The latter constitute a significant health care burden in the paediatric population and entail a substantial use of resources at both the outpatient and the inpatient care level.²⁻⁴ At the international

level, there is wide variability in the rates of hospitalization reported in the literature, which depend on the country where the study is conducted, the ages of the individuals under study and the methodology used. Most hospital admissions occur in healthy children,⁵ but there are also well-known risk factors for admission, such as young age (age <6–12 months), membership in specific ethnic groups and underlying chronic disease.⁶

Few population-based studies have spanned more than 1 flu season. Furthermore, nationwide data for the impact of influenza on hospital admissions in the Spanish paediatric

population are not available, although there is partial evidence from regional or small sample-size case series.^{1,5,7–10} The primary objective of our study was to establish the rate of hospitalization due to influenza in children aged less than 15 years in Spain based on data from 10 tertiary care hospitals. The secondary objectives included assessing the need for paediatric intensive care, the frequency of comorbidities and the history of vaccination against influenza in hospitalised children.

Methods

We conducted an epidemiological, multicentre, retrospective observational study through the review of health records of children aged less than 15 years admitted with infection by influenza virus in 2 consecutive epidemic seasons (September 15, 2014 to April 15, 2015 and September 15, 2015 to April 15, 2016). We conducted the study in 4 Spanish regions that were geographically distant, including 10 university hospitals): Basque Country (Hospital Universitario [HU] Basurto and HU Donostia), Madrid (HU 12 de Octubre, HU Clínico San Carlos, Hospital General Universitario [HGU] Gregorio Marañón, HU La Paz), Catalonia (HU Germans Trias i Pujol, HU Sant Joan de Déu, HU Vall d'Hebron) and Andalusia (Hospital Regional Universitario [HRU] de Málaga). We collected data on the catchment population of these hospitals and the number of admissions made (excluding surgical and neonatal admissions) in the 2 periods under study.

We included children aged less than 15 years hospitalised for more than 24 h with a primary clinical diagnosis of influenza virus infection with microbiological confirmation. We excluded children who had onset 48 h or later following admission, as these could be cases of nosocomial infection.

We obtained the lists of patients from the departments of clinical records, preventive medicine and epidemiology or microbiology of participating hospitals. We reviewed all the health records in which the discharge report included as the primary diagnosis (first listed diagnosis) codes 487.0–487.8 of the International Classification of Diseases (ICD) 9th revision, or J09–J18 of the ICD 10th revision, in the databases of participating hospitals, collecting data for several demographic and clinical variables. We entered the previously anonymised data in an online database designed for the purpose that was shared by all participating hospitals, including information on demographic, clinical and treatment variables.

The methods used for detection of influenza A and B viruses in upper respiratory tract secretion samples were commercially available rapid immunochromatographic assays for influenza virus antigens, multiplex real-time PCR (preceded by reverse RNA transcription) or viral culture in MDCK, A549 and LLC-MK2 cell lines using the shell-vial technique.

An external consultant performed the statistical analysis using the R software, without input from any of the researchers.¹¹ We calculated the incidence of hospital admission using the catchment population of the 10 referral hospitals as the denominator, which amounted to 878 808 children aged less than 15 years (12.5% of the Spanish population in this age range). Since we were unable to determine

the age distribution in every hospital, we calculated the rate of hospitalization for certain age groups (<6 months, 2 years, <5 years) using as denominator a reference population estimated based on the age distribution of the entire Spanish population in 2017.¹²

The study protocol was approved by the ethics boards of all participating hospitals and submitted to the Agencia Española de Medicamentos y Productos Sanitarios (Spanish Agency of Medicines and Medical Devices, AEMPS). We conducted the study in adherence with the international ethical guidelines for research and clinical studies in humans of the Declaration of Helsinki, the principles of good clinical practice and the guidelines for clinical research of the AEMPS. All collected data were kept confidential and handled in adherence with Decree SAS/3470/2009 and the Law on the Protection of Personal Data.

Results

We identified 984 cases of influenza, of which we excluded 77 (8.5%) that we suspected to be nosocomial infections. Thus, the final sample comprised 907 cases, of which 389 (42.9%) occurred in the first flu season and 518 (57.1%) in the second. Of the total cases, 815 (89.9%) corresponded to patients hospitalised in the paediatric ward, and the other 92 (10.1%) to patients that required admission to the paediatric intensive care unit (PICU) at some point. The mean estimated annual rate of hospitalization due to influenza per 1000 children in the reference population for the 2 seasons under study was 0.51 (95% CI, 0.48–0.55), and it varied between regions, going from 0.31 (95% CI, 0.25–0.40) in Andalusia (HRU de Málaga) to 0.36 (95% CI, 0.31–0.43) in Catalonia hospitals, 0.46 (95% CI, 0.35–0.62) in hospitals in the Basque Country and 0.83 (95% CI, 0.73–0.94) in hospitals in Madrid (Table 1). Overall, infection by influenza virus caused 1.60% (95% CI, 1.51–1.71%) of all paediatric admissions during the study period (mean of the 2 seasons), excluding surgical and neonatal admissions.

The sex distribution was 508 male (56%) and 399 female (44%), and the mean age was 3.2 years (median, 2 years), with 49.3% aged less than 2 years and 77.6% aged less than 5 (Table 2). The annual rate of hospitalization decreased with age, with an estimated rate of 3.48/1000 children aged less than 6 months (95% CI, 2.82–4.26), of 2.14 in children aged less than 24 months (95% CI, 1.88–2.44), 1.31 in children aged less than 60 months (95% CI, 1.18–1.45) and 0.17 in children aged 5–15 years (95% CI, 0.14–0.20). Hospital admissions were most frequent between January and March (90%) (Fig. 1).

Most of the patients were admitted due to respiratory symptoms (wheezing, bronchiolitis, pneumonia/bronchopneumonia), documented in 492 cases (54.2%) and fever without a focus, documented in 256 (28.2%). The working diagnosis was suspected sepsis in 27 cases (3.3%) and encephalitis in 10 (1.1%) (Table 3). Of all patients, 409 (45.1%) had comorbidities or underlying disease, with a single such problem in 288 cases and more than 1 in 121. The most frequent comorbidities were moderate or severe asthma and associated conditions, documented in 140 patients (15.4%), followed by immunosuppression in 88 cases (9.7%), preterm birth in 84 (9.3%) and congenital

Table 1 Rate of hospitalization due to influenza per 1000 children in the catchment population. Distribution by geographical area.

Geographical area	Population (N) ^a	Hospital admissions due to influenza (n) ^b	Rate per 1000 children (95% CI) ^c	Relative frequency (%) ^d
Madrid	281 380	464	0.83 (0.73–0.94)	51.2
Basque Country	97 685	90	0.46 (0.35–0.62)	9.9
Barcelona	400 311	290	0.36 (0.31–0.43)	32.0
Malaga	99 432	63	0.31 (0.25–0.40)	6.9
Total	878 808	907	0.51 (0.48–0.55)	100

CI, confidence interval (lower bound-upper bound).

^a Catchment populations of university hospitals used as reference (age < 15 years): Madrid (Gregorio Marañón, San Carlos, La Paz and 12 de Octubre); Basque Country (Donostia [San Sebastian] and Basurto [Bilbao]); Barcelona (San Joan de Déu, German Trías i Pujol and Vall d'Hebron); Malaga (Carlos Haya).

^b Total number of cases in the 2 seasons.

^c Mean annual rate of the 2 seasons.

^d Percentage of total cases diagnosed in each geographical area.

Table 2 Patients admitted with community-acquired influenza. Estimated age distribution and incidence by age group.

Age	Absolute frequency	Percentage (95% CI)	Reference population ^a	Mean annual incidence (95% CI)
<6 months	178	19.6 (17.2–22.3)		3.48 (2.82–4.26)
6–24 months	269	29.7 (26.8–32.7)	78 840	1.71 (1.45–2.03)
>24 months to <60 months	257	28.3 (25.5–31.4)	165 249	0.78 (0.65–0.92)
5 years to <10 years	135	14.9 (12.7–17.4)	308 437	0.22 (0.17–0.28)
10 to <15 years	68	7.5 (6.0–9.4)	300 570	0.11 (0.08–0.16)
<24 months	447	49.3 (46.0–52.5)	104 551	2.14 (1.88–2.44)
<60 months	704	77.6 (74.8–80.2)	269 800	1.31 (1.18–1.45)
Total	907	100	878 808	0.51 (0.48–0.55)

CI, confidence interval (lower bound-upper bound).

^a Estimate of the age distribution of the reference population based on the distribution of the Spanish population (National Institute of Statistics 2017 census) (cases/1000 children).


**Figure 1** Distribution by month (number of cases) of hospital admissions due to community-acquired influenza infection (2014–2015 and 2015–2016 flu seasons).

Table 3 Reason for admission in the 907 children hospitalised due to community-acquired influenza.

	Absolute frequency	Percentage (95% CI)
Fever and dyspnoea	295	32.5 (29.6–35.6)
Fever without a focus	256	28.2 (25.4–31.2)
Fever and vomiting/diarrhoea	93	10.3 (8.4–12.4)
Pneumonia/bronchopneumonia	109	12.0 (10.1–14.3)
Acute respiratory disease	125	13.8 (11.7–16.2)
Bronchiolitis with wheezing	51	5.6 (4.3–7.3)
Bronchiolitis	44	4.8 (3.6–6.5)
Bronchitis	39	4.3 (3.2–5.8)
Laryngitis/tracheitis	28	3.1 (2.1–4.4)
Suspected sepsis	27	3.0 (2.0–4.3)
Seizure	65	7.2 (5.7–9.0)
Encephalitis	10	1.1 (0.6–2.0)
Miscellaneous	14	1.5 (0.9–2.6)

CI, confidence interval (lower bound-upper bound).

Main reason for admission. There were cases with more than 1 diagnosis at admission.

heart disease in 81 (8.9%). Other frequent comorbidities were chronic encephalopathy (61 cases; 6.7%), neuromuscular disorders (29 cases; 3.2%), chromosome disorders (21 cases, 2.3%), upper gastrointestinal tract malformation (21 cases, 2.3%) and bronchopulmonary dysplasia (20 cases, 2.2%). Of the 716 patients, 36 (5%) were overweight, with a weight below the 95th percentile. The mean weight of admitted children was 14.7 ± 11.34 kg (range, 2.5–98), corresponding to a mean percentile of 37 ± 30 .

We were able to obtain information on the vaccination history of 301 patients aged more than 6 months (41.3%). Of the 143 children in the group that had one or more chronic diseases in whom vaccination was indicated (due to chronic respiratory disease [44.3%], immunosuppression [24.5%], neuromuscular disease [19.8%], preterm birth before 37 weeks' gestation [18.9%], cardiovascular disease or congenital heart defects [12.6%]), only 37 (25.9%) had been vaccinated (**Table 4** details the most frequent comorbidities in the 106 children that had not been vaccinated). On the other hand, 145 children had received antibiotic therapy prior to admission, and 18 antiviral medication, of who 8 had a relevant history of disease: immunosuppression ($n=5$), preterm birth ($n=3$), bronchopulmonary dysplasia ($n=1$), bronchiolitis obliterans ($n=1$) and severe asthma ($n=1$).

In 626 patients (68.0%), the causative agent was influenza A virus, which predominated in both seasons, while influenza B virus was involved in another 278 (30.7%), while the type

Table 4 Most frequent comorbidities in children aged more than 6 months that were not vaccinated against influenza ($n=106$).

Disease	Absolute frequency	Percentage (95% CI)
Moderate/severe asthma	40	37.7 (29.1–47.3)
Immunosuppression	26	24.5 (17.3–33.5)
Preterm birth	20	18.9 (12.6–27.3)
Encephalopathy	16	15.1 (9.5–23.1)
Cardiovascular	13	12.2 (7.3–19.9)

Other: bronchopulmonary dysplasia (4), sickle-cell anaemia (3), cystic fibrosis (2), upper gastrointestinal tract malformation (2), diabetes mellitus type 1 (1) and chromosome disorder (1).

was unknown in 3 cases (**Table 5**). In 634 children, sterile samples were collected for bacterial culture (blood, cerebrospinal fluid, etc.), resulting in isolation of pathogens in 15 cases (2.4%): *Streptococcus pneumoniae* in 9 cases, *Streptococcus pyogenes* in 4 and *Staphylococcus aureus* in 2.

Ninety-two children (10.1%) required admission to the PICU, and the most frequent reasons for it were: acute respiratory failure or apnoea in 73 (79.3%); seizures in 9 (9.8%); heart failure in 6 (6.5%), and decreased level of consciousness in 4 (4.3%). Of the patients admitted to the PICU, 59% had some type of chronic disease, and patients with chronic disease were more likely to require intensive supportive care (OR = 1.84 [95% CI, 1.19–2.85]; $P=.008$). In the hospitalised group, the proportion of children with underlying disease increased with age, ranging from 26.4% in infants aged less than 6 months to 73.5% in children aged more than 10 years ($\chi^2 = 45.7$; $P < .001$) (**Table 6**). The percentage of children with influenza that needed supportive care in the PICU ranged between 9.7% and 14.7% depending on the age group, except for children aged 5 to less than 10 years, in whom it was 3.7% ($\chi^2 = 7.26$; $P = .007$).

The mean length of stay was 6.76 days, with the cumulative stay amounting to a total of 7010 inpatient days, of which 6129 days corresponded to the general paediatrics ward and 881 days to the PICU. During their hospitalization, 524 children were treated with oseltamivir (57.8%) and 488 with antibiotics (53.8%). In addition, 311 received bronchodilators (34.3%), 225 corticosteroids (24.8%) and 412 oxygen therapy (45.4%). There were no deaths among the children included in the study, and none exhibited sequelae associated with the influenza infection at the time of discharge.

Discussion

We conducted this study in 10 tertiary care hospitals in 6 cities in 4 different geographical areas that were distant from each other, which together served a total population of nearly 900 000 children aged less than 15 years, slightly more than 10% of the Spanish population in this age range. For all the above reasons, we expect that our findings may be generalizable to the entire paediatric population of Spain.

Overall, the mean annual rate of hospitalization due to community-acquired influenza in the 2014–2015 and

Table 5 Detection of influenza viruses for each season under study.

Virus	Season 1 n (%)	Season 2 n (%)	Total
Influenza A	282 (72.5) ^a	344 (66.4) ^a	626
Influenza B	104 (26.7)	174 (33.6)	278
Undetermined type	3 (0.7)	0 (0.0)	3
Total	389	518	907

^a Subtype analyses were performed for 90 samples in the first season (31.9%), of which 57 corresponded to subtype H3N2 (63.3%) and 33 to the 2009 pandemic H1N1 subtype (36.7%). Subtype analyses were performed in 100 samples in the second season (29.1%), of which 11 corresponded to subtype H3N2 (11%) and 89 to the 2009 pandemic H1N1 subtype (89%).

Table 6 Association between age of hospitalised children, admission to PICU and presence of chronic disease.

Age group	Hospitalised, n	Chronic disease, n (%)	Admitted to PICU, n (%)	Chronic disease, n (%)
<6 months	178	47 (26.4)	23 (12.9)	15 (65.2)
6 to <24 months	269	105 (39.0)	29 (11.8)	15 (51.7)
2 to <5 years	257	132 (51.4)	25 (9.7)	15 (60.0)
5 to <10 years	135	75 (55.5)	5 (3.7)	4 (80.0)
≥10 years	68	50 (73.5)	10 (14.7)	5 (50.0)
Total	907	409 (45.1)	92 (10.1)	54 (58.7)

2015–2016 seasons was high (0.51/1000 children <15 years; 95% CI, 0.48–0.55), and consistent with the findings of longitudinal studies conducted in developed countries in children aged 0–15 or 17 years: 0.70/1000 in the United States (Philadelphia, 2000–2004)¹³; 0.67/1000 in Germany (Kiel, 1996–2001)¹⁴; 0.36/1000 in a study with a 16-year follow-up in Finland (Turku, 1988–2004)¹⁵; 0.22–0.24/1000 in another long-term study in Sweden (Stockholm, 1998–2014)¹⁶ and the rate estimated in a nationwide study conducted in Canada (2003–2014), of 0.15/1000.¹⁷ Nearly half of the hospitalised patients in our study were aged less than 2 years. In fact, the rate of hospitalization decreased drastically with age, with the rate in infants aged less than 6 months being twice that in children aged 6–24 months, 4-fold that in children aged 2–5 years, 20-fold that in children aged 5–10 years and 32-fold that in children aged 10–15 years, which was consistent with the previous literature.^{5,13,17–19} It is well known that in infants and young children, fever without a focus and sepsis-like presentations are frequent manifestations of influenza that prompt hospitalization for the main purpose of ruling out bacterial infection.^{5,20–22} The rate in infants aged less than 6 months (3.48/1000 [95% CI, 2.82–4.26]) was similar to the rates found in Finland (2.76/1000 [95% CI, 2.20–3.36]) (1988–2004),¹⁵ Sweden (1.44/1000 [95% CI, 0.69–3.31]) (1998–2014)¹⁶ or the United States in the pre-pandemic period (2000–2009), with annual rates between 1.6 and 9.1/1000 (studies conducted in 3 cities).^{19,23,24}

In children aged less than 5 years, the rate found in our study (1.31/1000 [95% CI, 1.18–1.45]) was somewhat higher compared to those reported by other authors of studies conducted in European countries such as Spain, Germany or Finland (0.9–1.23/1000),^{7,14,15} or in the United States (0.9/1000 in 2000–2004 and 0.58/1000 in 2004–2009).^{19,23} Regardless of the variability found in the literature, the studies we reviewed evince a substantial burden of influenza in paediatric services at different levels of care.

A constant feature in all case series was the identification among children hospitalised due to influenza of a subset with underlying chronic diseases that are considered risk factors for severe or complicated influenza and who accounted for an important proportion of the total cases (between 25% and 49%),^{8,13,21,25,26} which was 45% in our study. The comorbidities detected in our study were those expected based on past evidence,^{8,13,27} chiefly asthma and chronic respiratory diseases, followed by immunosuppression, neurologic or neuromuscular diseases, congenital heart defects and preterm birth.

We also ought to mention that we excluded 77 cases where we believed the infection had been contracted in the hospital, a finding that reinforces the need for yearly vaccination of all health care staff to reduce the incidence of these events.

An aspect that has not been studied frequently is the need for paediatric intensive care in children hospitalised with influenza. In our study, 10% of admitted children required this type of care, most frequently due to respiratory failure, and this proportion fell approximately in the middle of the range of the proportions reported in studies conducted in other countries (5–19%),^{8,13,15,16,21,23} which shows that the course of disease can be severe. In Spain, during the 2009 pandemic and in the first season that followed, Hernández-Bou et al.²⁸ and Launes et al.²⁶ found proportions of PICU admission of 17.5% ($n=54$) in patients aged less than 14 years and 14% ($n=20$) in patients aged 6 months to 18 years, respectively. As was the case in our study, the previous literature has also described a greater need for intensive care in children with underlying chronic diseases, which are considered risk factors for severe or complicated influenza.^{8,13,21} In fact, 59% of the children admitted to the PICU in our study had underlying chronic diseases, and this percentage was even higher, of 77%, in a recent study conducted in 24 German PICUs (2010–2013).²⁷

Despite the significant impact of influenza in the paediatric population in terms of hospitalization, we ought to note that of the 301 patients aged more than 6 months for whom we could obtain data regarding vaccination (the flu vaccine can be administered starting at age 6 months) a significant percentage (35.2%, $n=106$) corresponded to children with risk factors for severe disease that had not been vaccinated and in whom vaccination was indicated. This finding, which corroborates previous results in Spain,^{8,10,26} suggests that the impact of influenza on the Spanish paediatric population could be reduced significantly by adhering to current vaccination recommendations. Infants aged less than 6 months also amounted to a substantial proportion of the total admissions (20%), which highlights the need to recommend universal vaccination during pregnancy, a measure that would prevent at least a portion of these infections²⁹ and has been specifically recommended by the World Health Organization in 2012, in addition to advising parents to adopt additional precautions to prevent infection in infants, such as vaccination of household members or avoiding exposure of the infant to individuals with respiratory infections or fever.²¹

There are some limitations to this study, so its results should be interpreted with caution. First of all, it has a retrospective design, so we could not explore differences in clinical practice (such as situations where children were not hospitalised) or in the approach to diagnosis (such as the use of virologic testing) between different hospitals. Secondly, the catchment populations of participating hospitals may not be representative of the entire Spanish population, as most of the sample was drawn from predominantly urban areas and from 6 cities in particular. In addition, influenza outbreaks can vary in intensity based on the year, geographical area and age group, among other factors.^{8,19} In this regard, this study spanned 2 flu seasons (2014–2015 and 2015–2016) with a low-to-moderate incidence of influenza,³⁰ and only covered the usual months of the season, so we may have missed cases occurring outside these intervals. Thus, the incidence results should be considered minimum values.

In conclusion, influenza continues to cause a significant number of paediatric hospitalizations in Spain every year; a substantial proportion of cases (more than 50%) occur in infants aged less than 6 months and in children with underlying disease, in whom the administration of the vaccine against influenza is clearly indicated (in case of infants aged less than 6 months, by vaccination of the mother during pregnancy),^{31,32} and who clearly must be targeted more intensely by interventions aiming at improving vaccine coverage.

Funding

The study was funded by the Instituto de Investigación Sanitaria Biodonostia (San Sebastian, Spain) as sponsor of the project. Protocol code: HOSPIGRIP16.

Conflicts of interest

EGPY, GCE, JAF, EGLL, THSP, MJMP, JTRA, JRC, MMH, JJGG, CRGL, DMP have no conflicts of interest to declare.

Acknowledgments

The authors of the article want to thank all of the following:

J. Arrizabalaga, director of the Instituto de Investigación Sanitaria Biodonostia (sponsor of the HOSPIGRIP study), and L. Codesido, for the management of the HOSPIGRIP research project.

J. Arístegui, the scientific director, and E.G. Pérez-Yarza, the national coordinator of the HOSPIGRIP study.

All the researchers in the HOSPIGRIP research group.

L.M. Molinero (Alce Ingeniería, Madrid, Spain) for IT support and for conducting the statistical analysis.

Appendix A. Authors of the HOSPIGRIP research group

J. Aristegui, E. Garrote, A. Larrauri, Hospitals and researchers: Hospital Universitario (HU) Basurto, Bilbao, E.G. Pérez-Yarza, G. Cilla, M. Unsain, HU Donostia, San Sebastian, J. Ruiz Contreras, E. García-Ochoa, J.C. Gordillo, HU 12 de Octubre, Madrid, T. Hernández Sampelayo, R. Rodríguez, F. González, Hospital General Universitario Gregorio Marañón, Madrid, M.J. Mellado, C. Calvo, A. Méndez, J. Bustamante, D. Salas, C. Lacasta, HU La Paz, Madrid, J.T. Ramos, M. Illán, HU San Carlos, Madrid, M. Mendez, M. Barjuan, HU Germans Trias i Pujol, Badalona, J.J. García, S. Urraca, M. Caballero, C. Launes, HU Sant Joan de Déu, Barcelona, C. Rodrigo, A. Fàbregas, R. Esmel, A. Antón, HU Vall d'Hebron, Barcelona, D. Moreno, A.I. Valdivielso, P. Piñero, B. Carazo, Hospital Regional Universitario de Málaga, Malaga

References

1. Navarro-Marí JM, Palacios del Valle E, Pérez-Ruiz M, De la Rosa M. The impact of influenza viruses on hospitalizations in infants younger than 2 years old during epidemics of respiratory syncytial virus infection. *Clin Microbiol Infect.* 2003;9:959–63.
2. Antonova EN, Rycroft CE, Ambrose CS, Heikkinen T, Principi N. Burden of paediatric influenza in Western Europe: a systematic review. *BMC Public Health.* 2012;12:968.
3. Lafond KE, Nair H, Rasooly MH, Valente F, Booy R, Rahman M, et al., Global Respiratory Hospitalizations—Influenza Proportion Positive (GRIPP) Working Group. Global role and burden of influenza in pediatric respiratory hospitalizations, 1982–2012: a systematic analysis. *PLoS Med.* 2016;13:e1001977.
4. Ruf BR, Knuf M. The burden of seasonal and pandemic influenza in infants and children. *Eur J Pediatr.* 2014;173:265–76.
5. Arostegui-Kareaga N, Montes M, Pérez-Yarza EG, Sardón O, Vicente D, Cilla G. Clinical characteristics of children hospitalized for influenza virus infection. *An Pediatr (Barc).* 2005;62:5–12.
6. Muñoz FM. Influenza virus infection in infancy and early childhood. *Paediatr Respir Rev.* 2003;4:99–104.
7. Montes M, Vicente D, Pérez-Yarza EG, Cilla G, Pérez-Trallero E. Influenza-related hospitalizations among children aged less than 5 years old in the Basque Country Spain: a 3-year study (July 2001–June 2004). *Vaccine.* 2005;23:4302–6.
8. Rojo JC, Ruiz-Contreras J, Fernández MB, Marín MA, Folgueira L. Influenza-related hospitalizations in children younger than 3 years of age. *Pediatr Infect Dis J.* 2006;25:596–601.
9. Oñate-Vergara E. Características clínicas y epidemiológicas de los niños hospitalizados por infección por virus influenza durante

- la pandemia del año 2009 en Gipuzkoa. Comparación con la época prepandémica y pospandémica (2004–2012) en Gipuzkoa [tesis doctoral]. Universidad del País Vasco (UPV/EHU). Donostia, julio del 2013.
10. Ortiz-Lana N, Garrote E, Arístegui J, Rementeria J, García-Martínez JA, McCoig C, et al. Estudio prospectivo para estimar la carga de hospitalización y visitas a urgencias de la gripe en población pediátrica en Bilbao (2010–2011). *An Pediatr (Barc)*. 2017;87:299–362.
 11. R Core Team. R Foundation for Statistical Computing. A language and environment for statistical computing Vienna, Austria; 2017 [accessed 14 April 2018]. Available in: <https://www.r-project.org/>
 12. Instituto Nacional de Estadística. Datos demográficos. [accessed 14 April 2018]. Available in: http://www.ine.es/dyngs/INEbase/es/categoría.htm?c=Estadística_P&cid=1254735572981
 13. Coffin SE, Zaoutis TE, Rosenquist AB, Heydon K, Herrera G, Bridges CB, et al. Incidence, complications, and risk factors for prolonged stay in children hospitalized with community-acquired influenza. *Pediatrics*. 2007;119:740–8.
 14. Weigl JA, Puppe W, Schmitt HJ. The incidence of influenza-associated hospitalizations in children in Germany. *Epidemiol Infect*. 2002;129:525–33.
 15. Silvennoinen H, Peltola V, Vinionpää R, Ruuskanen O, Heikkinen T. Incidence of influenza-related hospitalizations in different age groups of children in Finland: a 16-year study. *Pediatr Infect Dis J*. 2011;30:e24–8.
 16. Bennet R, Hamrin J, Wirgart BZ, Östlund MR, Örtqvist Å, Eriksson M. Influenza epidemiology among hospitalized children in Stockholm Sweden 1998–2014. *Vaccine*. 2016;34:3298–302.
 17. Schanzer DL, Sabou M, Lee L, Nwosu A, Bancej C. Burden of influenza, respiratory syncytial virus, and other respiratory viruses and the completeness of respiratory viral identification among respiratory inpatients, Canada, 2003–2014. *Influenza Other Respir Viruses*. 2018;12:113–21.
 18. Iwane MK, Edwards KM, Szilagyi PG, Walker FJ, Griffin MR, Weinberg GA, et al. New vaccine surveillance network population-based surveillance for hospitalizations associated with respiratory syncytial virus, influenza virus, and parainfluenza viruses among young children. *Pediatrics*. 2004;113:1758–64.
 19. Poehling KA, Edwards KM, Griffin MR, Szylagy PG, Staat MA, Iwane MK, et al. The burden of influenza in young children, 2004–2009. *Pediatrics*. 2013;131:207–16.
 20. Bender JM, Ampofo K, Gesteland P, Sheng X, Korgenski K, Raines B, et al. Influenza virus infection in infants less than 3 months of age. *Pediatr Infect Dis J*. 2010;29:6–9.
 21. Chaves SS, Perez A, Farley MM, Miller L, Schaffner W, Lindgren ML, et al., Influenza Hospitalization Surveillance Network. The burden of influenza hospitalizations in infants from 2003 to 2012, United States. *Pediatr Infect Dis J*. 2014;33:912–9.
 22. Dagan R, Hall CB. Influenza A virus infection imitating bacterial sepsis in early infancy. *Pediatr Infect Dis J*. 1984;3:218–21.
 23. Phoepling KA, Edwards KM, Weinberg GA, Szilagyi P, Staat MA, Iwane MK, et al. New Vaccine Surveillance Network. The under recognized burden of influenza in young children. *N Engl J Med*. 2006;355:31–40.
 24. Fell DB, Johnson J, Mor Z, Katz MA, Skidmore B, Neuzil KM, et al. Incidence of laboratory-confirmed influenza disease among infants under 6 months of age: a systematic review. *BMJ Open*. 2017;7:e016526.
 25. Ampofo K, Gesteland PH, Bender J, Mills M, Daly J, Samore M, et al. Epidemiology, complications, and cost of hospitalization in children with laboratory-confirmed influenza infection. *Pediatrics*. 2006;118:2409–17.
 26. Launes C, García-García JJ, Martínez-Planas A, Moraga F, Soldevila N, Astigarraga I, et al., CIBERESP Cases and Controls in Pandemic Influenza Working Group. Clinical features of influenza disease in admitted children during the first post-pandemic season and risk factors for hospitalization: a multicentre Spanish experience. *Clin Microbiol Infect*. 2013;19:E157–62.
 27. Streng A, Prifert C, Weissbrich B, Liese JG, Bavarian PICU Study Group on Influenza and Other Viral ARI. Continued high incidence of children with severe influenza A(H1N1)pdm09 admitted to paediatric intensive care units in Germany during the first 3 post-pandemic influenza seasons, 2010/11–2012/13. *BMC Infect Dis*. 2015;15:573.
 28. Hernández-Bou S, Novell CB, Alins JG, García-García JJ, Infectious Diseases Working Group, Spanish Society of Pediatric Emergencies. Hospitalized children with influenza A H1N1 (2009) infection: a Spanish multicenter study. *Pediatr Emerg Care*. 2013;29:49–52.
 29. Madhi SA, Cutland CL, Kuwanda L, Weinberg A, Hugo A, Jones S, et al., Maternal Flu Trial (Matflu) Team. Influenza vaccination of pregnant women and protection of their infants. *N Engl J Med*. 2014;371:918–31.
 30. Instituto de Salud Carlos III. Informe de Vigilancia de la Gripe en España. Temporada 2015–2016 (desde la semana 40/2015 hasta la semana 20/2016). Sistema de Vigilancia de la Gripe en España [accessed 14 April 2018]. Available in: <http://www.isciii.es/ISCIII/es/contenidos/fd-servicios-cientifico-tecnicos/fd-vigilancias-alertas/fd-enfermedades/gripe.shtml>
 31. Committee on Infectious Diseases. Recommendations for prevention and control of influenza in children, 2017–2018. *Pediatrics*. 2017;140, pii: e20172550.
 32. Comité Asesor de Vacunas de la AEP. Vacunación frente a la gripe estacional en la infancia y la adolescencia. Recomendaciones 2017–2018. AEP [Internet]. October; 2017 [accessed 14 April 2018]. Available in: http://vacunasaep.org/sites/vacunasaep.org/files/recomendaciones_cav-aep_vacunacion_gripe_2017-18.pdf