

Deconstructing the narrow-line region of the nearest obscured quasar

M. Villar Martín^{1,2}, E. Bellocchi¹, J. Stern³, C. Ramos Almeida^{4,5}, C. Tadhunter⁶

R. González Delgado⁷

¹*Centro de Astrobiología (INTA-CSIC), Carretera de Ajalvir, km 4, E-28850 Torrejón de Ardoz, Madrid, Spain. villarmm@cab.inta-csic.es*

²*Astro-UAM, UAM, Unidad Asociada CSIC, Facultad de Ciencias, Campus de Cantoblanco, E-28049, Madrid, Spain*

³*Max Planck Institute für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany*

⁴*Instituto de Astrofísica de Canarias, C/ Vía Láctea, s/n, E38205 La Laguna, Tenerife, Spain*

⁵*Departamento de Astrofísica, Universidad de La Laguna, E-38206, La Laguna, Tenerife, Spain*

⁶*Department of Physics and Astronomy, University of Sheffield, Sheffield S3 7RH, UK*

⁷*Instituto de Astrofísica de Andalucía (CSIC), Glorieta de la Astronomía, s/n, 18008 Granada, Spain*

Accepted 2015 August 11. Received 2015 July 14; in original form 2015 January 18

ABSTRACT

We study the physical and kinematic properties of the narrow line region (NLR) of the nearest obscured quasar MRK 477 ($z=0.037$), using optical and near-infrared (NIR) spectroscopy. About 100 emission lines are identified in the optical+NIR spectrum (90 in the optical), including several narrow *optical* Fe⁺ lines. To our knowledge, this is the first type 2 active galactic nucleus (AGN) with such a detection. The Fe⁺ lines can be explained as the natural emission from the NLR photoionized by the AGN. Coronal line emission can only be confirmed in the NIR spectrum.

As in many other AGN, a significant correlation is found between the lines full width at half-maximum and the critical density $\log(n_{\text{crit}})$. We propose that it is caused by the outflow. This could be the case in other AGNs.

The nuclear jet-induced ionized outflow has been kinematically isolated in many emission lines covering a broad range of ionization potentials and critical densities. It is concentrated within $R \sim \text{few} \times 100$ pc from the central engine. The outflowing gas is denser ($n \gtrsim 8\,000\text{ cm}^{-3}$) than the ambient non-perturbed gas ($n \sim 400\text{--}630\text{ cm}^{-3}$). This could be due to the compression effect of the jet induced shocks. Alternatively, we propose that the outflow has been triggered by the jet at $R \lesssim 220$ pc (possibly at $\lesssim 30$ pc) and we trace how the impact weakens as it propagates outwards following the radiation-pressure dominated density gradient.

The different kinematic behaviour of [FeII] $\lambda 1.644\text{ }\mu\text{m}$ suggests that its emission is enhanced by shocks induced by the nuclear outflow/jet and is preferentially emitted at a different, less reddened spatial location.

Key words:

galaxies: active quasars: emission lines - quasars: general - quasars: individual: MRK 477

1 INTRODUCTION

MRK 477 (SDSS J144038.1+533016, $z=0.037$ and luminosity distance $D_L=161$ Mpc) is a type 2 luminous active galactic nucleus (AGN). Also known as I Zw 92, it was first identified by Zwicky (1966) as a compact galaxy. It is usually referred to as the most luminous Seyfert 2 in the local Universe. As pointed out by Heckman et al. (1997), it has the highest [O III] $\lambda 5007$ luminosity ($L_{[\text{OIII}]}=3.3 \times 10^{42}\text{ erg s}^{-1}$) of any of the 140 Seyfert nuclei (type 1 or 2) com-

piled by Whittle (1992) and the fifth highest radio power ($P_{1.4\text{GHz}}=2.0 \times 10^{30}\text{ erg s}^{-1}\text{ Hz}^{-1}$ at 1.4 GHz). Indeed, its high $L_{[\text{OIII}]}$ places it in the regime of optically selected obscured quasars (QSO2), according to the selection criteria defined by Zakamska et al. (2003), $L_{[\text{OIII}]} > 1.2 \times 10^{42}\text{ erg s}^{-1}$. The [OIII] luminosity implies a bolometric luminosity $L_{\text{bol}}=8.6 \times 10^{45}\text{ erg s}^{-1}$ (Stern & Laor 2012; see also Lamas-tra et al. 2009), which is in the quasar range (Shen et al. 2011).

The quasar host galaxy is interacting with an emission line companion, possibly a LINER, located 50 arcsec (~ 36 kpc) to the north (De Robertis et al. 1987). Hubble Space Telescope (HST) images are shown in Fig. 1¹. Both galaxies are connected by a faint bridge and prominent tidal tails are apparent. The ratio of stellar masses between the quasar host and the companion is $M_1/M_2=1.6$ (Koss et al. 2012).

The quasar host shows a compact blue central source, peculiar among type 2 AGNs, but no evidence for a broad line region (BLR), prompting some authors to suggest that the quasar lacks one. However, polarization observations have revealed a hidden BLR (e.g. Tran et al. 1992; Tran 1995; Shu et al. 2007). The nuclear continuum polarization is $\lesssim 1.5\%$, significantly lower than that of the broad lines, implying that the majority of the optical continuum cannot be due to scattered light from the hidden type 1 nucleus. The blue central source is instead associated with a compact dusty starburst (effective radius < 0.2 kpc) which occurred ~ 6 Myr ago in a brief period of time and whose light dominates from the ultraviolet to the near-infrared (NIR; Heckman et al. 1997, González Delgado et al. 1998). This is therefore a starburst-AGN hybrid system. MRK 477 has a star-forming rate $SFR=24 M_\odot \text{ yr}^{-1}$ inferred from the infrared (IR) luminosity (assuming that it is starburst-dominated) $\log \frac{L_{\text{IR}}}{L_\odot}=11.14$ (Krug et al. 2010). This places it in the regime of luminous IR galaxies (LIRGs: $11 \leq \log \frac{L_{\text{IR}}}{L_\odot} < 12$). The detection of the Wolf Rayet (WR) blue bump around the He II $\lambda 4686$ line by Heckman et al. (1997) led these authors to propose that MRK 477 is a luminous member of the class of WR galaxies.

The QSO2 host harbors a supermassive black hole with a mass in the range $\log(M_{\text{BH}}) \sim 7.18$ (as derived from the M_{BH} versus σ_* correlation) to 8.84 (as inferred from the polarimetric H β line; Zhang, Bian & Huang 2008).

Although radio quiet according to the $L_{[\text{OIII}]}$ versus L_{rad} , radio loud versus radio quiet, classification criteria (explained in Villar-Martín et al. 2014), MRK 477 shows a clear excess of radio emission compared to that expected from the stellar contribution. The 8.4 GHz Very Large Array (VLA) radio continuum map (0.26 arcsec resolution) shows a triple radio source (~ 1.2 arcsec total extension) whose morphology correlates with that of the narrow line region (NLR). The size and overall northeast-southwest axis of the radio structures are shared by the [OIII] emission, which extends up to a similar distance to the north-east (Heckman et al. 1997).

Shuder & Osterbrock (1981) reported for the first time that the emission lines have a large blue-ward asymmetry (i.e., excess) at low intensity level relative to the continuum. Villar Martín et al. (2014) proposed that the ionized nuclear outflow responsible for such an asymmetry has been triggered by the interaction between the radio source and the NLR. These authors proposed that negative feedback can be triggered by the radio structures in a significant fraction of radio quiet quasars. Thanks to its high intrinsic luminosity

and closeness, MRK 477 is an excellent test object to gain further insight into this mechanism.

We present here a detailed study of the optical and near-infrared spectra of MRK 477. We explore a diversity of aspects that provide a more complete understanding of the nature of this object, example of a type 2 quasar in the nearby universe, as well as a starburst-AGN hybrid system. The paper is organized as follows. The spectra are described in Sect. 2, together with the spectral fitting procedure. We present results in Sect. 3 regarding line identification, correlations between the gas kinematics and a diversity of parameters (ionization potential, critical density), the spatial extension of the ionized gas, the ionized outflow, the presence of narrow Fe⁺ emission lines, the coronal line spectrum and the detection of WR features. The results are discussed in Sect. 4 and the conclusions are presented in Sect. 5.

We assume $\Omega_\Lambda=0.7$, $\Omega_M=0.3$, $H_0=71 \text{ km s}^{-1} \text{ Mpc}^{-1}$. At $z=0.037$, 1 arcsec corresponds 0.725 kpc (Wright 2006).

2 DATA SET

2.1 Optical spectrum

The optical spectrum was obtained as part of the Sloan Digital Sky Survey (SDSS, York et al. 2000). It spans the rest frame range $\sim 3660\text{--}8880 \text{ \AA}$ range (Figs. 2 and 3). It corresponds to an aperture defined by the 3 arcsec diameter SDSS fibre (~ 2.2 kpc at $z=0.037$) centred at the galaxy nucleus. For comparison, the triple radio source identified by Heckman et al. (1997) has a total extension of ~ 1.2 arcsec and thus it is well within the fibre area.

The spectral resolution is $\sim 180 \pm 20 \text{ km s}^{-1}$. All values of emission line full width at half-maximum (FWHM) have been corrected for instrumental broadening.

The main emission lines have large equivalent widths and the stellar features are comparatively weak. Underlying stellar absorption of the Balmer lines is expected to be negligible and therefore subtracting the stellar continuum is not necessary for our purposes.

2.2 NIR spectrum: observations and data reduction

NIR H+K long-slit spectra were obtained with the NIR camera/spectrometer LIRIS (Long-slit Intermediate Resolution Infrared Spectrograph; Acosta-Pulido et al. 2003; Manchado et al. 2004), attached to the Cassegrain focus of the 4.2 m William Herschel Telescope (WHT). LIRIS is equipped with a Rockwell Hawaii 1024 \times 1024 HgCdTe array detector, whose spatial scale is 0.25 arcsec pixel⁻¹.

The NIR observations were performed as part of the Isaac Newton Group service programme (programme SW2014b22) in two different nights, 28 March 2015 and 02 May 2015. The H+K grism was used. It covers the 1.388–2.419 μm spectral range and provides a dispersion of 9.7 \AA pix⁻¹. The seeing size during the March observations was FWHM $\sim 2''$ (as measured from the standard star). A 1'' wide slit was used oriented at position angle PA=43° East of North. This PA is aligned with the radio axis and the NLR axis as seeing in the HST image presented by Heckman et al. (1997). The resulting spectral resolution was $35.7 \pm 2.2 \text{ \AA}$ (or

¹ The HST images were retrieved from the HST Science Legacy Archive. They are part of the programmes 9379 with principal investigator PI H. Schmitt (ACS/HRC image), 8597 with PI M. Michael Regan (WFPC image) and 7330 with PI J. Mulchaey (NICMOS image).

Mrk 477




Figure 1. Hubble Space Telescope (HST) images of MRK 477. They were obtained with Wide-Field Planetary Camera 2 (WFPC2, F606W filter, left), the Advanced Camera for Surveys (ACS/HRC, F330W filter, middle) and the near-infrared Camera and Multi-Object Spectrometer (NICMOS, F160W filter, right) respectively. The flux intensity maps are represented in logarithmic scale such that $F [\text{erg s}^{-2} \text{cm}^{-2} \text{\AA}^{-1} \text{arcsec}^{-2}] = 10^{A+b}$, where A is a constant factor (i.e., -18.3, -14.5 and -16.4, respectively, for WFPC2, ACS and NICMOS images) and b is indicated in the coloured bar. The white box in the WFPC2 image identifies the FoV covered by the ACS and NICMOS images. The horizontal line at the bottom left of these images corresponds to a scale of 10 kpc for the WFPC2 image and to a scale of 2 kpc for the ACS and NICMOS images. The spatial scale for this galaxy is 0.725 kpc/ arcsec . North is at the top and East to the left in all the panels.

$550 \pm 34 \text{ km s}^{-1}$ at the observed wavelength of $\text{Pa}\alpha$). In Fig. 4 we show the NIR spectrum extracted from a $3.5''$ aperture, centred at the position of the nucleus. The total exposure time on source was 6400 s (16×400 s).

The May data were obtained with seeing $\text{FWHM} = 0.65 \pm 0.05''$ (measured from the standard star). A $0.75''$ slit was used, also with $\text{PA} = 43^\circ$. The resulting spectral resolution was $23.2 \pm 1.2 \text{ \AA}$ (or $318 \pm 18 \text{ km s}^{-1}$). The observations were performed through clouds and the signal to noise ratio of the final spectrum was low. In spite of this, the good seeing conditions allowed to obtain more accurate information on the spatial extension of the strongest emission line $\text{Pa}\alpha$. The total exposure time on source was 4000 s (10×400 s). All FWHM values of the NIR spectra have been corrected for instrumental broadening.

The data were reduced following standard procedures for near-IR spectroscopy, using the LIRISDR dedicated software within the IRAF² environment. For a detailed description of the reduction process see Ramos Almeida et al. (2009). Consecutive pairs of AB two-dimensional spectra were subtracted to remove the sky background. The resulting frames were then wavelength calibrated and flat-fielded before registering and co-adding all frames to provide the final spectra.

The absolute flux calibration of the March spectrum is intended to be an approximation since the spectra of the comparison star are likely subjected to slit losses due to centring and tracking errors. We note that in this work we refer to flux ratios only, and therefore the uncertainty in the

absolute flux calibration of the spectra does not affect our results.

2.3 Spectral fitting

In order to study the kinematic and physical properties of the nuclear ionized outflow we have fitted the spectral profiles of numerous emission lines (Sect. 3). For this we used the STARLINK package DIPSO. This software is based on the optimization of fit coefficients, in the sense of minimizing the sum of the squares of the deviations of the fit from the spectrum data. The output from a completed fit consists of the optimized parameters (Gaussian central λ , FWHM, intensity peak, flux) and their errors (calculated in the linear approximation, from the error matrix).

Two methods were attempted. In method I, we assumed that all emission lines have the same kinematic substructure as $[\text{OIII}]\lambda\lambda 4959, 5007$ (same number of kinematic components with identical FWHM in km s^{-1} and relative velocities, as explained in Villar Martín et al. 2014). In physical terms, this method corresponds to a situation where all gaseous regions emit all lines, although with different relative fluxes.

On the other hand, this is not necessarily the case and some lines might come from different regions, possibly resulting in variations of the kinematic substructure from line to line. As an example, low critical density lines ($\sim \text{few} \times 10^3 \text{ cm}^{-3}$) are quenched in high density gas with $n > 10^6 \text{ cm}^{-3}$. To account for this possibility, whenever possible we have also attempted to fit the lines without applying prior kinematic restrictions (method II).

Also, all fits that produced unphysical results were rejected even if mathematically valid (e.g. a fit producing $[\text{NII}] \frac{\lambda 6583}{\lambda 6548}$ very different from the theoretical value 3.0).

As we will see the lines are complex with multiple kine-

² IRAF is distributed by the National Optical Astronomy Observatory, which is operated by the Association of Universities for the Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation (<http://iraf.noao.edu/>).




Figure 2. SDSS spectrum of MRK 477. Flux is given in units of $\times 10^{-13}$ erg s $^{-1}$ cm $^{-2}$ Å $^{-1}$ and the rest-frame wavelength is in Å. Some of the strongest emission lines are indicated.




Figure 3. Four zoomed spectral windows covering the full spectral range are shown to highlight some of the weakest lines. Unless otherwise specified, fluxes in these and other figures are given in units of $\times 10^{-14}$ erg s $^{-1}$ cm $^{-2}$ Å $^{-1}$. Some relevant lines are indicated.

| (1) Species | (2) λ_{air} (Å) | (3) λ_{obs} (Å) | (4) $\frac{Flux}{Flux(H\beta)}$ | (5) n_{crit} (cm ⁻³) |
|---------------------------------|--------------------------------------|--------------------------------------|------------------------------------|---|
| [OII] | 3727.0+3728.8 | 3726.9 | 2.26 ± 0.18 | (1.3/4.5) × 10 ³ |
| H12 | 3750.2 | 3750.1 | 0.014 ± 0.003 | |
| [FeVII] | 3758.9 | 3759.3 | 0.046 ± 0.005 | 3.7 × 10 ^{7a} |
| H11 | 3770.6 | 3770.4 | 0.023 ± 0.002 | |
| H10 | 3797.9 | 3797.5 | 0.026 ± 0.003 | |
| H9 | 3835.4 | 3835.5 | 0.045 ± 0.003 | |
| [NeIII] | 3868.8 | 3868.6 | 1.04 ± 0.03 | 1.3 × 10 ⁷ |
| HeI+H8 | 3888.7+3889.05 | 3888.8 | 0.16 ± 0.02 | |
| [NeIII]+H7 | 3967.5+3970.1 | 3968.2 | 0.43 ± 0.04 | 1.3 × 10 ⁷ |
| HeI | 4026.2 | 4026.1 | 0.013 ± 0.002 | |
| [SII] | 4068.6 | 4068.6 | 0.19 ± 0.01 | 3.0 × 10 ⁶ |
| [SII] | 4076.4 | 4076.1 | 0.065 ± 0.03 | 1.5 × 10 ⁶ |
| Hδ | 4101.7 | 4101.5 | 0.24 ± 0.02 | |
| [FeV] | 4180.9 | 4180.7 | 0.008 ± 0.002 | |
| FeII];[FeV] | 4227.2;4228.0 | 4228.2±0.1 | 0.017 ± 0.002 | |
| [FeII] | 4244.0 | 4244.5 | 0.019 ± 0.002 | |
| [FeII] | 4276.8 | 4278.3±0.1 | 0.013 ± 0.002 | |
| [FeII] | 4287.4 | 4289.1±0.4 | 0.014 ± 0.002 | |
| Hγ | 4340.5 | 4340.31 | 0.45 ± 0.05 | |
| [OIII] | 4363.2 | 4363.2 | 0.23 ± 0.02 | 3.4 × 10 ⁷ |
| [FeII] | 4416.3 | 4416.4 | 0.021 ± 0.002 | |
| HeI | 4471.4 | 4472.2 | 0.035 ± 0.002 | |
| [FeII] | 4570.0 | 4570.9 | 0.018 ± 0.005 | |
| [FeIII];CIV | 4658.1 | 4658.7 | 0.034 ± 0.001 | |
| HeII | 4685.7 | 4685.7 | 0.138 ± 0.008 | |
| [FeIII] | 4701.5 | 4700.2 | 0.010 ± 0.008 | |
| [ArIV] | 4711.4 | 4711.5 | 0.011 ± 0.002 | 1.7 × 10 ⁴ |
| [NeIV] | 4725.5;4726.8 | 4725.1 | 0.008 ± 0.003 | (6.2/2.5) × 10 ⁷ |
| [ArIV] | 4740.2 | 4740.2 | 0.016 ± 0.003 | 1.6 × 10 ⁵ |
| [FeIII] | 4754.7 | 4754.0±0.5 | 0.007 ± 0.002 | |
| [FeIII] | 4769.5 | 4769.6 | 0.006 ± 0.002 | |
| [FeIII] | 4777.7 | 4777.5 | 0.007 ± 0.002 | |
| [Fe II] | 4814.5 | 4816.3± | 0.016 ± 0.002 | |
| Hβ | 4861.3 | 4861.3 | 1.00 | |
| [FeVII] | 4893.4 | 4893.9 | 0.008 ± 0.002 | |
| [OIII] | 4958.9 | 4958.9 | 3.7 ± 0.3 | 8.6 × 10 ⁵ |
| [OIII] | 5006.8 | 5006.8 | 10.5 ± 0.3 | 8.6 × 10 ⁵ |
| [FeVI] | 5145.8 | 5145.9 | 0.007 ± 0.002 | 2.7 × 10 ^{7a} |
| [FeVII] | 5158.9 | 5159.4 | 0.042 ± 0.005 | 3.3 × 10 ^{6a} |
| [FeVI] | 5176.4 | 5177.0 | 0.010 ± 0.001 | 3.3 × 10 ^{7a} |
| [NI] | 5197.9+5200.4 | 5199.2 | 0.05 ± 0.01 | (1.6/0.51) × 10 ³ |
| [Fe III];[FeII];[FeII];[Fe VII] | 5270.4;5273.4;5276.0;5276.4 | 5272.1 | 0.058 ± 0.004 | |
| [CaV] | 5309.1 | 5309.7 | 0.032 ± 0.002 | 5.1 × 10 ⁷ |
| [FeVI] | 5335.2 | 5335.5 | 0.017 ± 0.002 | |
| [FeII] | 5379.0 | 5379.7 | 0.011 ± 0.002 | |
| HeII];[FeII] | 5411.5;5412.7 | 5412.0 | 0.015 ± 0.005 | |
| FeII];[FeVI] | 5425.3;5425.7 | 5424.1 | 0.018 ± 0.006 | |
| [FeVI] | 5485.0 | 5484.5 | 0.005 ± 0.001 | |
| [Cl III] | 5517.7 | 5517.5 | 0.006 ± 0.002 | 8.5 × 10 ³ |
| [ArX];[FeII];FeII] | 5533.0;5527.6;5534.8 | 5531.39 | 0.013 ± 0.003 | |
| [Cl III] | 5537.9 | 5537.9 | 0.009 ± 0.002 | 2.9 × 10 ⁴ |
| [OI] | 5577.34 | 5577.9 | 0.014 ± 0.001 | 9.0 × 10 ⁷ |
| [FeVI] | 5631.1 | 5630.9 | 0.008 ± 0.002 | |
| [FeVI] | 5677.0 | 5677.1 | 0.005 ± 0.001 | |
| [FeVII] | 5720.9 | 5721.8±0.2 | 0.047 ± 0.002 | 3.6 × 10 ^{7a} |
| [NII] | 5754.6 | 5754.7 | 0.040 ± 0.002 | 1.8 × 10 ⁷ |
| HeI | 5875.6 | 5875.7 | 0.131 ± 0.007 | |
| [FeVII] | 6086.9 | 6087.7 | 0.12 ± 0.01 | 3.6 × 10 ^{7a} |
| [OI] | 6300.3 | 6300.5 | 0.81 ± 0.02 | 1.1 × 10 ⁶ |
| [SIII] | 6312.1 | 6311.8 | 0.016 ± 0.003 | 1.6 × 10 ⁷ |
| [OI] | 6363.8 | 6365.0 | 0.26 ± 0.02 | 1.1 × 10 ⁶ |
| [ArV] | 6435.1 | 6435.7 | 0.010 ± 0.002 | 1.4 × 10 ⁷ |
| [NII] | 6548.1 | 6548.1 | 0.53 ± 0.05 | 1.2 × 10 ⁵ |
| Hα | 6562.8 | 6562.8 | 3.9 ± 0.2 | |
| [NII] | 6583.5 | 6583.4 | 1.5 ± 0.1 | 1.2 × 10 ⁵ |
| HeI | 6678.2 | 6678.7 | 0.035 ± 0.005 | |
| [SII] | 6716.4 | 6716.5 | 0.60 ± 0.03 | 1.5 × 10 ³ |
| [SII] | 6730.8 | 6730.9 | 0.75 ± 0.03 | 4.0 × 10 ³ |
| [ArV] | 7005.8 | 7005.7 | 0.014 ± 0.003 | 1.3 × 10 ⁷ |
| HeI | 7065.7 | 7065.3 | 0.07 ± 0.005 | |
| [ArIII] | 7135.8 | 7135.8 | 0.20 ± 0.01 | 6.4 × 10 ⁶ |
| [FeII] | 7155.2 | 7156.8 | 0.06 ± 0.01 | |
| [FeII] | 7172.0 | 7173.8 | 0.014 ± 0.004 | |
| [CaII] | 7291.5 | 7292.8 | 0.02 ± 0.01 | |
| [OII] | 7318.9 | 7321.0 | 0.23 ± 0.04 | 6.5 × 10 ⁶ |
| [OII] | 7339.7 | 7329.2 | 0.32 ± 0.04 | 6.5 × 10 ⁶ |
| [NII] | 7377.8 | 7379.1 | 0.040 ± 0.008 | |
| [FeII] | 7452.6 | 7453.9±0.5 | 0.016 ± 0.002 | |
| FeI];[FeII] | 7634.0;7637.5 | 7639±2 | 0.007 ± 0.002 | |
| [ArIII] | 7751.1 | 7750.9 | 0.062 ± 0.003 | 6.4 × 10 ⁶ |
| MgII]; [NiIII]; [FeXI]; MgII | 7877.0; 7889.9; 7891.8;7896.0 | 7888.8 | 0.023 ± 0.003 | |
| MgII | 8232.0 | 8232.7±0.5 | 0.020 ± 0.005 | |
| OI | 8446.0 | 8446.6 | 0.029 ± 0.002 | |
| HI Pa 14 | 8598.5 | 8595.4 | 0.010 ± 0.003 | |
| [FeII] | 8605.5 | 8604.5 | 0.033 ± 0.004 | |
| [FeII] | 8617.0 | 8617.8 | 0.05 ± 0.02 | |
| HI Pa 12 | 8750.6 | 8750.2 | 0.013 ± 0.002 | |

Table 1. Emission lines identified in the SDSS spectrum of MRK 477. The observed line wavelengths λ^{obs} (column 3) have been determined by fitting a single Gaussian to the emission lines. Errors are quoted when they provide relevant information regarding possible multiple identifications or there is a significant shift between the observed and the air wavelengths (column 2). The line ratios (column 4) are given relative to Hβ, with $F(H\beta)=(1.01\pm0.04)\times10^{-13}$ erg s⁻¹ cm⁻². They are not corrected for reddening. The critical densities are quoted when available. Superscript ^a in column (5) indicates that n_{crit} has been taken from De Robertis & Osterbrock (1984). The other n_{crit} values have been computed with the PYNEB software (see the text).

| (1) Species | (2) λ_{air} (Å) | (3) λ_{obs} (Å) | (4) $\frac{\text{Flux}}{\text{Flux}(\text{Br}\gamma)}$ | (5) n_{crit} (cm^{-3}) |
|------------------------|--------------------------------------|--------------------------------------|---|--|
| [Si X] | 1.4305 | 1.4308 | 0.61 ± 0.10 | $6.3 \times 10^8^{(a)}$ |
| [Fe II] | 1.5339 | 1.5337 | 0.79 ± 0.10 | $4.6 \times 10^4^{(b)}$ |
| [Fe II] | 1.6440 | 1.6435 | 4.5 ± 0.5 | $5.6 \times 10^4^{(b)}$ |
| Pa α | 1.8756 | 1.8746 | 12.6 ± 1.4 | |
| H ₂ 1-0S(3) | 1.9576 | 1.9564 | 0.84 ± 0.10 | |
| [Si VI] | 1.9629 | 1.9626 | 0.97 ± 0.12 | $6.3 \times 10^8^{(a)}$ |
| H ₂ 1-0S(2) | 2.0338 | 2.0329 | 0.20 ± 0.02 | |
| HeI | 2.0587 | 2.0575 | 0.29 ± 0.03 | |
| H ₂ 1-0S(1) | 2.1218 | 2.1206 | 0.63 ± 0.17 | |
| Br γ | 2.1661 | 2.1650 | 1.00 ± 0.15 | |

Table 2. Emission lines identified in the NIR spectrum of of MRK 477 extracted from a $\sim 1 \text{ arcsec} \times 3.5 \text{ arcsec}$ aperture centered on the nucleus. $^{(a)}n_{\text{crit}}$ from Rodríguez Ardila et al. (2011). $^{(b)}n_{\text{crit}}$ from Drougados et al. (2010). Fluxes are quoted relative to Br γ , for which $F(\text{Br}\gamma) = (3.0 \pm 0.3) \times 10^{-15} \text{ erg s}^{-1} \text{ cm}^{-2}$, although the absolute flux calibration is uncertain (see the text).




Figure 5. The FWHM (top panels) and the velocity shift V_s (bottom panels) of a subsample of ~ 30 forbidden lines are plotted against the critical density ($\log(n_{\text{crit}})$), the lower and higher ionization potentials of the species involved. The measurements from De Robertis & Osterbrock (1986, dR & Ost 86) are also shown in the top-left panel as open circles. The outlier at $\log(n_{\text{crit}}) \sim 3.1$ and $\text{FWHM} \sim 500 \text{ km s}^{-1}$ corresponds to [OII] $\lambda 3727$, which appears artificially broader due to blended doublet components. The NIR lines are marked with red triangles. [FeII] $\lambda 1.644$ is a clear outlier in the top panels. It is too broad compared to what is expected from the general trend defined by the rest of the lines.

| Comp. | FWHM (km s ⁻¹) | V _S (km s ⁻¹) | $\frac{F_i}{F(\text{H}\beta)}$ |
|-------------|-------------------------------|---|--------------------------------|
| [SII]λ4068 | | | |
| Narrow | 94±37 | -2±8 | 0.07±0.01 |
| Interm. | 505±35 | 17±10 | 0.26±0.02 |
| Broad | 1534±97 | -376±85 | 0.25±0.05 |
| [SII]λ4076 | | | |
| Narrow | 94±37 | -2±8 | 0.010±0.006 |
| Interm. | 505±35 | 17±10 | 0.12±0.02 |
| Broad | 1534±97 | -376±85 | 0.08±0.07 |
| Hδ | | | |
| Narrow | 94±37 | -2±8 | 0.23±0.04 |
| Interm. | 505±35 | 17±10 | 0.23±0.06 |
| Broad | 1534±97 | -376±85 | 0.26±0.03 |
| Hγ | | | |
| Narrow | 157±28 | -7±7 | 0.47±0.03 |
| Interm. | 529±40 | 36±10 | 0.38±0.04 |
| Broad | 1830±133 | -224±51 | 0.57±0.06 |
| [OIII]λ4363 | | | |
| Narrow | 157±28 | -7±7 | 0.11±0.02 |
| Interm. | 529±40 | 36±10 | 0.43±0.03 |
| Broad | 1830±133 | -224±51 | 0.27±0.08 |
| Hβ | | | |
| Narrow | 137±22 | -2±6 | $(4.1\pm0.4)\times10^{-14}$ |
| Interm. | 471±22 | 7±7 | $(3.7\pm0.1)\times10^{-14}$ |
| Broad | 1527±75 | -188±24 | $(2.3\pm0.1)\times10^{-14}$ |
| [OIII]λ5007 | | | |
| Narrow | 177±15 | 5±9 | 8.5±0.9 |
| Interm. | 545±15 | 22±11 | 13.1±0.5 |
| Broad | 1839±53 | -227±23 | 9.6±0.7 |
| [OI]λ6300 | | | |
| Narrow | 211±11 | -4±5 | 0.60±0.07 |
| Interm. | 563±29 | 17±6 | 0.91±0.08 |
| Broad | 1745±40 | -257±24 | 1.08±0.07 |
| Hα | | | |
| Narrow | As [OIII] | As [OIII] | 3.1±0.3 |
| Interm. | As [OIII] | As [OIII] | 4.9±0.2 |
| Broad | As [OIII] | As [OIII] | 4.2±0.3 |
| [NII]λ6583 | | | |
| Narrow | As [OIII] | As [OIII] | 1.0±0.1 |
| Interm. | As [OIII] | As [OIII] | 1.70±0.07 |
| Broad | As [OIII] | As [OIII] | 0.9±0.2 |
| [SII]λ6716 | | | |
| Broad | 186±9 | -3±5 | 0.67±0.07 |
| Interm. | 501±17 | -4±6 | 0.68±0.04 |
| Broad | 1397±76 | -493±89 | 0.30±0.08 |
| [SII]λ6731 | | | |
| Narrow | 186±9 | -3±5 | 0.66±0.07 |
| Interm. | 501±17 | -4±6 | 0.80±0.03 |
| Broad | 1397±76 | -493±89 | 0.77±0.08 |

Table 3. Results of the spectral decomposition of the main lines in the spectrum of MRK 477. Three kinematic components are isolated in all cases, including a broad blueshifted component emitted by the ionized outflow. No values are quoted for Hα and [NII] because successful fits could only be obtained with full constraints from the [OIII] lines (see Sect. 2.1). The flux of each component relative to Hβ is given in the last column. The exception is Hβ, for which the actual fluxes are quoted.

velocity, high density and high ionization regions closer to the central engine.

We have done n estimations using the density-diagnostic line ratios [SII]λ6716/λ6731, [Cl III]λ5517/λ5537 and [ArIV]λ4711/λ4740, which are not affected by extinction or T_e variations. They are sensitive to different n regimes. The [SII] ratio ($n_{crit}=1.5\times10^3$ and 4.0×10^3 cm⁻³, respectively) is sensitive to n variations \sim several $\times10^3$ cm⁻³. These lines are suppressed in high density re-




Figure 6. Spatial profiles (normalized fluxes) of the NIR continuum and Paα along PA 43° N to E, compared with the seeing disk (star). Both the continuum and Paα are spatially extended.

gions by collisional de-excitation. Other line pairs that can be overcome this limitation are [Cl III]λ5517/λ5537 ($n_{crit}=8.5\times10^3$ and 2.9×10^4 cm⁻³), sensitive at $n \sim 10^4$ cm⁻³ and [ArIV]λ4711/λ4740 ($n_{crit}=1.7\times10^4$ and 1.6×10^5 cm⁻³) at $n \sim 10^4$ - 10^5 cm⁻³.

For MRK 477, [SII]λ6716/λ6731=0.70±0.06, [ClIII]λ5517/λ5537=0.74±0.22 and [ArIV]λ4711/λ4740=0.67±0.09, implying densities $n=1945^{+670}_{-460}$ cm⁻³ (from [SII]), $7250^{+13,300}_{-5100}$ (from [Cl III]) and $n=14,700^{+6150}_{-3750}$ (from [ArIV]). So, the [SII] doublet implies the existence of gas with $n \sim 2000$ cm⁻³, the [Cl III] doublet, in spite of the larger uncertainties suggests higher densities and the [ArIV] doublet confirms the existence of gas with $n > 10^4$ cm⁻³ in the NLR of MRK 477.

3.3 Spatial extension of the ionized gas

The only available information about the spatial distribution of the ionized gas in MRK 477 is provided by the [OIII] image presented by Heckman et al. (1997). It shows a bright knot or ridge of emission at ~ 0.4 arcsec to the northeast of the central source and aligned with the radio axis. This knot is not resolved from the nucleus in our NIR spectrum, for which the slit was roughly aligned along the same axis. We find, on the other hand, that Paα is extended at both sides of the central source. We show in Fig. 6 the spatial profiles of (1) a star observed during the May run (seeing size FWHM=0.65±0.05), (2) the MRK 477 continuum and (3) the Paα flux, with the underlying continuum subtracted. Although more compact than the continuum, Paα is extended, as demonstrated by the clear excess above the seeing wings, up to a maximum radial distance of ~ 2 arcsec or ~ 1.5 kpc to the NE.

3.4 The ionized outflow

Villar Martín et al. (2014) performed a kinematic and ionization characterization of the nuclear ionized outflow in MRK 477. Its signature is a broad blueshifted kinematic component in the [OIII]λλ4959,5007 lines, with




Figure 7. Left: data (black), fits (cyan) and residuals (dashed-red) for a diversity of emission lines. Right: individual kinematic components isolated in each line. Different line styles are used for different kinematic components. The same red colour is used for components with the same redshift: long-dashed red (narrow component), dotted-red (intermediate component), dot-dashed blue (broad component; i.e. the outflow emission). From top to bottom: [SII] $\lambda\lambda 4068, 4076$ and H δ ; H γ and [OIII] $\lambda 4363$; [NII] $\lambda\lambda 6548, 6583$ and H α ; [SII] $\lambda\lambda 6716, 6731$.




Figure 8. $\epsilon = \frac{F_{\text{broad}}}{F_{\text{narrow}}}$ parametrizes the relative contribution between the outflow flux emission and the rest of the line flux. The correlation with the critical density (panel A) implies that the outflow emission is relatively stronger in higher density gas. [OI] ‘a’ and ‘b’ correspond to the $\lambda 6300$ and $\lambda 6312$ lines respectively; [NeIII] ‘a’ and ‘b’ to $\lambda 3869$ and $\lambda 3967$; [SII] ‘b’ and ‘r’ refer to the blue and red [SII] doublets at $\lambda \lambda 6048, 6076$ and $\lambda \lambda 6716, 6731$ respectively. [FeII] $\lambda 1.644 \mu\text{m}$ (red triangle) is again an outlier.

FWHM ~ 1850 km s $^{-1}$ and $V_S \sim 225$ km s $^{-1}$. Based on the high turbulence of this gaseous component, the line ratios consistent with AGN (rather than stellar) excitation processes and the relatively high contribution of the outflowing gas to the total line fluxes, the authors proposed that it has been triggered by the 1.2 arcsec scale radio source. This is supported by the correlation between the radio and [OIII] morphologies (Heckman et al. 1997), which demonstrates that the radio source is interacting with the NLR.

3.4.1 Isolating the outflow emission in a diversity of emission lines

By means of the kinematic decomposition of the spectral profiles, we have isolated the emission from the outflowing gas in numerous emission lines (Sect. 2.1). Our goal is to characterize its physical properties and to constrain more accurately its spatial location.

We show in Table 3 the results of the fits for several optical emission lines (see also Fig. 7). FWHM and V_S correspond to those fits where no prior constraints from [OIII] were applied (method II, see Sect 2.3). When both methods could be applied, the flux values and errors account for the dispersion allowed by them.

Generally, we find rather coherent results for all strong lines for which multiple component fitting procedures could be applied. By “coherent” it is meant that all lines consist of three kinematic components (Table 3): two of them have similar z and are relatively narrow (FWHM \sim [95,210] and [470,560] km s $^{-1}$ respectively). A third broad blueshifted component is moreover isolated in all lines, which is emitted by the outflowing gas. It has FWHM \sim [1400,1840 km s $^{-1}$ and $V_S \sim$ [-490,-190] km s $^{-1}$. All the three kinematic components have line ratios consistent with type 2 AGN, as already pointed out by Villar Martín et al. (2014).

Other lines show clear evidence of a broad underlying component, although the triple Gaussian fit is not possible due to the low signal-to-noise and/or the complex blend with

neighbour lines. We have calculated the ratio $\epsilon = \frac{F_{\text{broad}}}{F_{\text{narrow}}}$ between the flux of the broad (outflowing) component and the rest of the line flux (which for simplicity we will name F_{narrow} , although it contains the narrow and intermediate components) for as many lines as possible. This gives a measurement of the relative contribution between the most turbulent outflowing gas and the more quiescent ambient gas.

The ratio ϵ is plotted against $\log(n_{crit})$, FWHM, IP_{low} and IP_{high} in Fig. 8. A significant correlation is found with the critical density ($r_s=0.73$ and $p=0.003$; panel A). No significant trend is found with the ionization potentials ($r_s=0.40$ and $p=0.15$ for ϵ versus IP_{low} and $r_s=0.35$ and $p=0.23$ for ϵ versus IP_{high}).

An interesting case is the high ionization line [FeVII] λ 6087, for which a very broad component is isolated with $\text{FWHM}=2460\pm340 \text{ km s}^{-1}$ (Fig. 9) and $V_S=-150\pm60 \text{ km s}^{-1}$. Thus, the line which shows simultaneously very high critical density and ionization potential, also shows the most extreme kinematics and the largest $\epsilon = 0.96\pm0.10$ of all optical lines. At the other end, the lowest n_{crit} (also low ionization) lines [NI] λ 5200 and [SII] $\lambda\lambda$ 6716,6731, are the narrowest ($\text{FWHM}\sim300 \text{ km s}^{-1}$) and have the smallest $\epsilon \sim 0.1\text{--}0.2$ (Fig. 8).

The correlation between ϵ and the FWHM suggests that the increasing broadening of the lines is due to the increasing outflow influence. The correlation with n_{crit} shows that the outflow emission is relatively stronger in higher density gas. Ultimately, these results suggest that the FWHM versus n_{crit} correlation is produced by the outflow in MRK 477.

[FeII] λ 1.644 is the only forbidden NIR line for which the spectral decomposition could be applied. It consists of a narrow, spectrally unresolved component with FWHM $\lesssim 340$ km s $^{-1}$ and a very broad and prominent underling component with FWHM $= 4770 \pm 830$ km s $^{-1}$ (Fig. 10) shifted by 315 ± 231 km s $^{-1}$ relative to the narrow core. It is the only line for which the broad component is not blueshifted. For comparison, Pa α consists of two components with FWHM $\lesssim 260$ km s $^{-1}$ and 2240 ± 230 km s $^{-1}$ respectively




Figure 9. Fit of the [FeVII]λ6087 spectral profile. The presence of a very broad underlying wing (FWHM~2500 km s⁻¹) shows that the high ionization Fe⁺6 region also participates in the outflow. Flux in units of 10⁻¹⁴ erg s⁻¹ cm⁻² Å⁻¹.




Figure 10. Fit of the NIR [FeII]λ1.644μm (top) and Paα (bottom) spectral profiles. Flux in units of 10⁻¹⁶ (top) and 10⁻¹⁵ (bottom) erg s⁻¹ cm⁻² Å⁻¹.

with the broad one blueshifted by -570 ± 119 km s⁻¹. As in Fig. 5, [FeII]λ1.64μm is a clear outlier in all ϵ diagrams, with $\epsilon = 1.3 \pm 0.2$, the highest of all lines. The line, therefore, has a dominant contribution of emission from the outflowing gas. The different behavior of this line will be discussed later (see below).

3.4.2 Reddening and electron density

(i) *Reddening correction.* Reddening correction (c and $E(B-V)$) has been estimated using the Balmer ratios $\frac{H\alpha}{H\beta}$, $\frac{H\gamma}{H\beta}$ and $\frac{H\delta}{H\beta}$, for which we assume case B values 2.8, 0.47 and 0.26 respectively, appropriate for gas densities $n \sim 10^2$ - 10^6 cm⁻³ and electron temperatures $T_e \sim 10,000$ - $20,000$ K (Osterbrock 1989).

We show in Table 4 the $E(B-V)$ and c values derived for the three kinematic components and the total line fluxes using the expressions:

$$E(B-V) = 1.99 \times \log \left[\frac{(\frac{H\alpha}{H\beta})_{\text{obs}}}{2.86} \right] \quad [\text{eq.1}]$$

and

$$\left[\frac{F}{H\beta} \right]_{\text{int}} = \left[\frac{F}{H\beta} \right]_{\text{obs}} \times 10^{c[f(\lambda) - f(H\beta)]} \quad [\text{eq.2}]$$

where “int” and “obs” denote the intrinsic and observed line ratios and $f(\lambda) - f(H\beta)$ is given by the standard interstellar extinction curve (Osterbrock 1989). The final values and errors take into account that negative c values are not allowed⁴. Once c is determined, other line ratios are corrected for reddening using the equation (2).

In spite of the complexity of the line profiles, the reddening values derived from the three ratios are in reasonable good agreement. From Table 3 we conclude that the narrowest component shows little or no reddening while the intermediate component shows the highest reddening. For the broad component, the results are less clear, since the Balmer decrement $\frac{H\alpha}{H\beta}$ suggests higher reddening ($E(B-V) = 0.32 \pm 0.06$) than the $H\delta$ and $H\gamma$ ratios, which imply little or no reddening. We cannot discard problems with the $H\alpha$ fit, which is severely blended with the [NII] doublet. However, it might be a real effect. As we will see next, the broad component is expected to have high densities $n \gtrsim 8000$ cm⁻³, maybe up to $\gtrsim 10^6$ cm⁻³, given the detection of very high critical density strong lines. Under these circumstances, the Balmer decrement can be enhanced due to collisional excitation of $H\alpha$. $n > 5 \times 10^5$ cm⁻³ will produce $\frac{H\alpha}{H\beta} > 4$ (e.g. Binette et al. 1993). Thus, the inconsistency between the reddening values inferred for the broad component might actually indicate the existence of high densities in the outflowing gas.

We measure $\text{Pa}\alpha/\text{Br}\gamma = 12.6 \pm 1.4$ from the NIR spectrum. Taking errors into account, this implies no reddening or, at most, $E(B-V) \leq 0.32$. This is consistent with the range of values allowed by the optical decrements.

⁴ This is the reason why in Table 3 the same line ratio with different errors (e.g. $\frac{H\gamma}{H\beta} = 0.23 \pm 0.04$ and 0.23 ± 0.06) can be associated with slightly different c and $E(B-V)$ values.




Figure 11. Constraints on the electron density n for the three kinematic components and for the integrated lines (adapted from Fig. 2 and 3 in Keenan et al. 1996)). A trend is found such that the broader the lines, the higher the density. Thus, the outflowing gas has the highest density.

| (1) Comp. | (2) $\frac{H\alpha}{H\beta}$ | (3) $c_{H\alpha}$ | (4) $E(B-V)_{H\alpha}$ | (5) $\frac{H\gamma}{H\beta}$ | (6) $c_{H\gamma}$ | (7) $E(B-V)_{H\gamma}$ | (8) $\frac{H\delta}{H\beta}$ | (9) $c_{H\delta}$ | (10) $E(B-V)_{H\delta}$ |
|--------------|---------------------------------|----------------------|---------------------------|---------------------------------|----------------------|---------------------------|---------------------------------|----------------------|----------------------------|
| Narrow | 3.1 ± 0.3 | 0.14 ± 0.12 | 0.08 ± 0.08 | 0.47 ± 0.03 | 0.09 ± 0.09 | 0.04 ± 0.03 | 0.23 ± 0.04 | 0.37 ± 0.37 | 0.24 ± 0.11 |
| Intermediate | 4.9 ± 0.2 | 0.70 ± 0.04 | 0.47 ± 0.03 | 0.38 ± 0.06 | 0.62 ± 0.46 | 0.42 ± 0.19 | 0.23 ± 0.06 | 0.40 ± 0.40 | 0.26 ± 0.12 |
| Broad | 4.2 ± 0.3 | 0.49 ± 0.08 | 0.32 ± 0.06 | 0.57 ± 0.09 | 0.00 | 0.00 | 0.26 ± 0.03 | 0.15 ± 0.15 | 0.09 ± 0.05 |
| Total | 4.0 ± 0.2 | 0.45 ± 0.06 | 0.29 ± 0.05 | 0.45 ± 0.05 | 0.21 ± 0.21 | 0.13 ± 0.06 | 0.24 ± 0.02 | 0.27 ± 0.23 | 0.17 ± 0.08 |

Table 4. Extinction correction (c and $E(B-V)$) determined from the Balmer ratios (columns 2, 5 and 8) for the three kinematic components and the integrated lines.

| (1) Comp. | (2) $r_1^{\text{obs}} = r_1^{\text{int}}$ | (3) r_2^{obs} | (4) r_2^{int} | (5) r_3^{obs} | (6) r_3^{int} | (7) $n \text{ (cm}^{-3}\text{)}$ |
|--------------|--|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------------------|
| Narrow | 1.03 ± 0.03 | 0.05 ± 0.01 | 0.07 ± 0.02 | 0.007 ± 0.005 | 0.010 ± 0.007 | $\sim 400\text{-}630$ |
| Intermediate | 0.73 ± 0.05 | 0.17 ± 0.02 | 0.37 ± 0.09 | 0.08 ± 0.01 | 0.17 ± 0.04 | $\sim 2000\text{-}4000$ |
| Broad | 0.41 ± 0.12 | 0.24 ± 0.05 | 0.34 ± 0.14 | 0.08 ± 0.07 | 0.25 ± 0.11 | $\gtrsim 8000$ |
| Total | 0.76 ± 0.06 | 0.13 ± 0.01 | 0.22 ± 0.03 | 0.05 ± 0.01 | 0.08 ± 0.02 | $\sim 1600\text{-}3200$ |

Table 5. Density sensitive line ratios observed ($^{\text{obs}}$) and corrected ($^{\text{int}}$) for reddening. $r_1 = \frac{[\text{SII}]\lambda 6716}{[\text{SII}]\lambda 6731}$, $r_2 = \frac{[\text{SII}]\lambda 4068}{[\text{SII}]\lambda 6716 + 6731}$, $r_3 = \frac{[\text{SII}]\lambda 4076}{[\text{SII}]\lambda 6716 + 6731}$. The range of densities n has been calculated using r_1 . A trend is found such that the broader the component, the higher the density.

(ii) *Electron density n .* We have constrained n with the ratio $r_1 = \frac{[\text{SII}]\lambda 6716}{[\text{SII}]\lambda 6731}$ (Osterbrock 1989). The results are shown in Table 5. The apparently non perturbed ambient gas (the narrow component) has $n \sim 400\text{-}630 \text{ cm}^{-3}$. The intermediate component, which shows intermediate kinematic and physical properties, has $n \sim 2,000\text{-}4,000 \text{ cm}^{-3}$. Finally, the most extreme kinematic component, which is associated with the outflowing gas has the highest density $n \gtrsim 8,000 \text{ cm}^{-3}$.

We have further checked these results using $r_2 =$

$\frac{[\text{SII}]\lambda 4068}{[\text{SII}]\lambda 6716 + 6731}$ and $r_3 = \frac{[\text{SII}]\lambda 4076}{[\text{SII}]\lambda 6716 + 6731}$ (Keenan et al. 1996). Unlike r_1 , these ratios have a strong dependence on the electron temperature T_e and reddening. They are therefore less efficient at constraining n , but they provide a useful test. The dependence of both ratios with n and T_e can be seen in Fig. 11.

To correct for reddening (eq. [2]), for each component we adopt c as the average of the maximum and minimum values obtained from the three Balmer ratios (Table 4). The uncertainties are calculated using half the difference between




Figure 12. Several narrow Fe^+ emission lines are detected in the spectrum of MRK 477. $[\text{CaII}]\lambda 7291$ is also tentatively detected (third panel).




Figure 13. Emission lines are detected from other refractory elements apart from Ca and Fe (Fig. 9), such as possibly Ni (top) and Mg (top and bottom).




Figure 14. Spectral fit, residuals and kinematic components of $[\text{OI}]\lambda 6300, 6364$. $[\text{SII}]\lambda 6312$ (black small Gaussian) is marginally detected as a small excess on the red side of the $[\text{OI}]\lambda 6300$. The detection of $[\text{FeX}]\lambda 6375$ cannot be confirmed. $[\text{OI}]\lambda 6364$ is successfully fitted without the contribution of such a line. Color and line style code as in Fig. 5.




Figure 15. Top: “blue bump” near HeII 4686 due to WR stars in the SDSS spectrum of MRK 477. Middle: the “red bump” (~ 5800 Å) is also detected. Bottom: The optical spectrum of the WR galaxy SBS 1222+614 is shown for comparison (taken from Guseva, Izotov & Thuan 2000). Flux for this spectrum in units of 10^{-16} erg s $^{-1}$ cm $^{-1}$ Å $^{-1}$.

these two values. The results are shown in Table 5 and Fig. 11. In this figure we show the n sensitive diagnostic diagrams $\log(r1)$ versus $\log(r2)$ and $\log(r1)$ versus $\log(r3)$ (Keenan et al. 1996). The coloured squares mark the areas covered by the allowed range of extinction corrected line ratios for the different kinematic components and the integrated line fluxes, taking into account all the uncertainties. The figure is consistent with the results from $r1$. They support a trend for increasing density from the narrow to the broad component.

In spite of the uncertainties inherent to fitting the mul-

tipole kinematic components in all the lines involved in the determination of n and reddening, a coherent picture can be drawn: the ionized gas within a radius of 1.5 arcsec or ~ 1.1 kpc (set by the size of the SDSS fibre) shows a gradient in physical and kinematic properties, which is apparent in the three kinematic components. The apparently non-perturbed ambient gas (the narrow component) has $n \sim 400$ -630 cm $^{-3}$. The intermediate component, which shows intermediate kinematic and physical properties, has $n \sim 2000$ -4000 cm $^{-3}$. Finally, the broadest component, which is associated with the outflowing gas has the highest density $\gtrsim 8000$ cm $^{-3}$. As mentioned above, the large $\frac{H\alpha}{H\beta}$ decrement might suggest densities as high as $n > 5 \times 10^5$ cm $^{-3}$ for the outflowing gas.

3.5 Narrow optical Fe $^{+}$ emission

Broad Fe $^{+}$ multiplet emission are prominent features in the optical spectra of most type 1 AGNs. Dong et al. (2010) have demonstrated statistically that *narrow optical* Fe $^{+}$ lines, either permitted or forbidden, are prevalent in type 1 AGNs, but are completely absent in type 2 AGNs across a wide luminosity range, from Seyfert 2 galaxies to QSO2.

MRK 477 is an exception. Heckman et al. (1997) detected [FeII] $\lambda 8617$ blended with some neighbour lines. We confirm the detection of more than 10 Fe $^{+}$ emission lines (see Table 1 and Fig. 12), besides several other features that might be blended or misidentified. The features for which the width could be measured show FWHM ~ 400 -650 km s $^{-1}$, suggesting similar kinematics as the NLR (Fig. 5, top panels). Other emission lines associated with highly refractory elements are also detected (Mg, Ni and, tentatively, Ca; Figs. 12 and 13).

3.6 Coronal lines

Coronal lines are collisionally excited forbidden transitions emitted within low-lying levels of highly ionized species ($IP_{low} > 100$ eV; Rodríguez Ardila et al. 2006). According to that exact definition, we cannot confirm the detection of any coronal feature in the *optical* spectrum of MRK 477, since the highest ionization lines we identify unambiguously are those produced by Fe $^{+6}$ ($IP_{low} = 99$ eV). On the other hand, coronal emission is confirmed in the NIR spectrum with the detection of [SiVI] $\lambda 1.963$ ($IP_{low} = 167$ eV) and [SiX] $\lambda 1.430$ ($IP_{low} = 351$ eV) (Fig. 4; Table 2).

Although [FeX] $\lambda 6375$ ($IP_{low} = 235$ eV) was reported as detected by De Robertis (1987) and Veilleux (1988), we cannot confirm this. Both groups used the same spectra in their analysis, one obtained in 1980 and another in 1985. The 1980 spectrum shows a strong emission feature at the right λ (see their Fig. 6, left), but this was discarded by Veilleux (1988) as an artifact. Although the 1985 spectrum shows a small excess near the [FeX] wavelength which the authors interpret as the detection of this line, our analysis shows that this is consistent with being due to the complex kinematic sub-structure of the [OI] $\lambda 6374$ line profile (Fig. 14).

A broad faint and noisy feature is detected with central $\lambda = 7888.8$ Å. Several lines are possibly contributing, including MgII $\lambda 7877$, [NiIII] $\lambda 7889$; [FeXI] $\lambda 7892$ and MgII $\lambda 7896$ (Fig. 13, top). Thus, the detection of [FeXI] emission ($IP_{low} = 262$ eV), which has been identified in some type

2 AGN (Penston et al. 1984, Rose et al. 2013) cannot be confirmed either.

The highest optical ionization lines unambiguously identified are [NeV] $\lambda\lambda$ 3346,3426 ($IP_{\text{low}}=97$ eV, out of the SDSS spectral range; reported by De Robertis 1987) and the [FeVII] lines ($IP_{\text{low}}=99$ eV) at λ 5159, 6087 Å (also reported by De Robertis 1987), and lines at 3759, 4893, 5721 Å. Also possibly [FeVII] at 5276 Å, [ArX] at 5533 Å, although other identifications cannot be discarded (Table 1).

With somewhat lower ionization level, [CaV] at 5309 Å ($IP_{\text{low}}=67$ eV) and several [FeVI] ($IP_{\text{low}}=75$ eV) lines at 5146, 5176, 5335, 5485, 5631, 5677 Å are also detected and possibly [FeV] λ 5726 ($IP_{\text{low}}=55$ eV) and [FeVI] at 5426 Å.

[FeVII] λ 6086 is fainter relative to low ionization lines (e.g. [OI] λ 6300) than usually found in active galaxies with strong coronal emission. $\frac{[\text{FeVII}]\lambda 6086}{[\text{OI}]\lambda 6300}=0.15\pm 0.01$ for MRK 477. For comparison, the Seyfert 1 and 2 galaxies studied by Rodríguez Ardila et al. (2006) with detected [FeX] and [FeXI] lines show in general (4 out of 5 objects) $\frac{[\text{FeVII}]\lambda 6086}{[\text{OI}]\lambda 6300}$ in the range ~ 0.6 -5.7.

3.7 The WR bumps

Heckman et al. (1997) detected a broad emission complex around He II λ 4686, which is clearly appreciated in the SDSS spectrum. They fitted this so called “blue bump” satisfactorily as a blend of the He II line together with other lines which are identified in Fig. 15 (top). This unresolved bump is a blend of lines emitted mainly by late WN and early WC WR stars, although some contribution of early WN stars might be present (Schaerer & Vacca 1998). Heckman et al. (1997) propose that it may be produced by an ensemble of about 30,000 WR stars (WN subtype), in which case MRK 477 would be a luminous (but not extraordinarily so) member of the class of WR galaxies.

Another feature often identified in WR galaxies due to WR stars is the so called “red bump” due to broad CIV λ 5808 emission, emitted mainly by WCE stars. This is usually much weaker than the “blue bump” and was not detected by Heckman et al. (1997).

We show in Fig. 15 (middle panel) the MRK 477 spectrum near the “red bump” and the optical spectrum of the WR galaxy SBS 1222+614 (Guseva, Izotov & Thuan 2000) for comparison (bottom panel). The appearance of the “red bump” in this and other WR galaxies is very similar to that in MRK 477. We thus believe that the “red bump” is detected in the SDSS spectrum of this QSO2. Its very large width (FWHM ~ 80 Å or ~ 4100 km s $^{-1}$) rules out as its origin the broad wings due to the ionized outflow identified in many other emission lines.

4 DISCUSSION

We have analysed the optical and NIR spectra of MRK 477, the nearest obscured quasar. WR features first identified by Heckman et al. (1997) and confirmed here by the new detection of the “red bump” at ~ 5800 Å (Sect. 3.7) show that it has undergone very recent star formation.

The optical+NIR spectrum of MRK 477 is very rich, with ~ 100 detected emission lines (~ 90 in the optical). In spite of the lack of spatial information of the SDSS spectrum,

the spectral decomposition of numerous lines has allowed us to characterize, at least partially, the spatial structure and the gradients in the physical and kinematic properties of the gas.

4.1 The origin of the FWHM versus n_{crit} correlation

As in many other type 1 and type 2 AGN, a significant correlation has been found between the FWHM of the lines and the critical density n_{crit} (in log). Based on the analysis of eight type 1 AGN spectra, Stern, Laor & Baskin (2014; S14 hereafter) showed that this relation is consistent with the velocity field within the black hole gravitational sphere of influence (radius R_g), assuming the $n \propto R^{-2}$ relation implied by the equilibrium between the NLR clouds and the radiation pressure and that the emission of each forbidden line is dominated by gas with $n \sim n_{\text{crit}}$. This explanation implies that beyond some threshold n_{crit} , n_{crit}^0 , which depends on the luminosity in Eddington units \dot{m} , the line emission will be dominated by gas within R_g , and therefore is expected to show larger velocities than gas which kinematics is dominated by the host galaxy. For MRK 477, the stellar velocity dispersion $\sigma_*=117$ km s $^{-1}$ (Zhang, Bian & Wang 2008), L_{bol} and the values of M_{BH} quoted in Sect. 1 imply $R_g \sim 1$ and 40 for $\log(M_{\text{BH}})=7.19$ and 8.84 respectively (eq. 26 in S14). This implies a threshold $n_{\text{crit}}^0=10^{8.6}$ cm $^{-3}$ (eq. 6 in S14) for $\log(M_{\text{BH}})=7.19$. This value is much larger than the $n_{\text{crit}} \sim 10^{4.5}$ - $10^{5.5}$ cm $^{-3}$ where the FWHM starts to increase in MRK 477 (top-left panel in Fig. 5), which makes the gravitational interpretation unlikely. For $\log(M_{\text{BH}})=8.84$, $n_{\text{crit}}^0=10^{5.4}$ cm $^{-3}$ is indeed consistent with the data. The success of this prediction supports the gravitational interpretation for the FWHM versus n_{crit} correlation, provided MRK 477 harbors such a massive black hole. On the other hand, the kinematic substructure of the lines raises doubts about whether the emission is truly emitted mostly by gas at $n \sim n_{\text{crit}}$. As an example, the [OIII] line flux is dominated by gas with $n \lesssim 4000$ cm $^{-3}$ (see Sect. 3.4).

Alternatively, we propose that the outflow produces this correlation in MRK 477. This is supported by the correlations between $\epsilon = \frac{F_{\text{broad}}}{F_{\text{narrow}}}$ and n_{crit} (Sect. 3.3.1), which imply that the outflow is relatively stronger and is responsible for the increasing line broadening at increasing densities. This correlation does not differ in any statistical sense from the usual FWHM versus n_{crit} one. The difference lies in the physical interpretation we propose.

It is possible that the effects of ionized outflows explain the FWHM versus n_{crit} correlation in other AGN.

Three kinematic components have been isolated in the main optical emission lines. The narrowest and intermediate components, have similar z ($V_S \sim 0$ km s $^{-1}$) and are relatively narrow (FWHM \sim [95-210] and [470-560] km s $^{-1}$ respectively). A third broad blueshifted component is moreover isolated in all lines with FWHM in the range \sim [1400-1840] km s $^{-1}$ and $V_S \sim$ [-490-(-190)] km s $^{-1}$. The narrow component is likely to trace the NLR ambient, non perturbed gas, while the broad component is emitted by the most turbulent, outflowing gas. The intermediate component traces gas of intermediate properties. All three components have line ratios consistent with type 2 active galaxies, as already pointed out by Villar Martín et al. (2014),

implying that they are spatially located within the quasar ionization cones.

The difference in kinematics is associated with a difference in physical properties. The broader the component, the higher the density. The sequence we find is $n \sim (400-630) \text{ cm}^{-3}$ for the ambient gas, $n \sim (2000-4000) \text{ cm}^{-3}$ for the intermediate component and $n \gtrsim 8000 \text{ cm}^{-3}$ for the outflowing gas (see §3.3.2). Higher n of the outflowing gas is also suggested by the correlation between $\epsilon = \frac{F_{\text{broad}}}{F_{\text{narrow}}}$ and n_{crit} for different forbidden lines (see above). Density enhancement of the outflowing gas has been found in some radio galaxies and QSO2 (e.g. Holt et al. 2011; Villar Martín et al. 2014).

4.2 The spatial location of nuclear ionized outflow

The isolation of the broad component in numerous emission lines demonstrates that the outflow involves gas covering a large range of ionization potentials and critical densities (at least $\text{IP}_{\text{high}} \sim 8$ to 125 eV , $\log(n_{\text{crit}}) \sim 3.3$ to 7.5). We try to constrain next the spatial location where it was triggered and how far its effects extend.

Bennert et al. (2006a,b) found that the observed gas density of the NLR gas decreases with increasing distance to the AGN in Seyfert 1 and 2 galaxies (see also S14). A similar behavior is expected in MRK 477. The difference in kinematic and physical properties between the broad, intermediate and narrow components can be naturally explained if they are located at increasing distance from the AGN, with the outflow emission dominating at smaller distances, in the inner part of the NLR or even closer (see also Villar-Martín et al. 2014). The outflow becomes weaker as it propagates outwards and reaches less dense regions, where lines of lower critical densities are preferentially emitted. Because the outflow has lost some power, it drags less and less mass and its emission becomes weaker relative to the ambient gas, which dominates the line fluxes in the most distant, lower density regions. This can also explain the correlation of ϵ with n_{crit} .

If the radiation pressure is responsible for the NLR density gradient (S14), the estimated n for the three kinematic components will help to constrain the distance from the central engine R at which each one is preferentially emitted. The expected behaviour of n with the ionizing luminosity L_{ion} and R is predicted to be (S14):

$$n = 7 \times 10^4 L_{\text{ion},45} R_{50}^{-2}$$

where $L_{\text{ion},45}$ is the ionizing luminosity in units of $10^{45} \text{ erg s}^{-1}$ and R_{50} is the distance in units of 50 pc.

Typical $\frac{L_{\text{bol}}}{L_{\text{ion}}}$ ratios for luminous Sy1 and QSO1 with L_{bol} in the range $\sim 10^{45-46} \text{ erg s}^{-1}$ have median value of 3.89 and standard deviation 4.36. Thus, for a ratio of 3.89, then $L_{\text{ion},45} \sim 2.2$ for MRK 477. Considering the range of n inferred for each gaseous component, the broad (outflowing), intermediate and narrow kinematic components would be located at $\lesssim 220$ (the upper limit is a consequence of the lower limit on $n \gtrsim 8000 \text{ cm}^{-3}$), 375 ± 65 and $880 \pm 100 \text{ pc}$ respectively. This sets an upper limit on the distance at which the outflow has been originated of $R \lesssim 220 \text{ pc}$. If densities as high as $\sim 5 \times 10^5 \text{ cm}^{-3}$ exist in the outflowing gas (Sect. 3.4.2), then $R \lesssim 30 \text{ pc}$. For comparison, the radio jet extends at $\sim 0.6 \text{ arcsec}$ or 435 pc from the central engine. With all

uncertainties involved, it is remarkable that these calculations place the intermediate and narrow components near and beyond the edge of the radio jet respectively. This adds further support to the idea that the radio jet has originated the outflow. In this scenario, our results suggest that the outflow in MRK 477 is concentrated in the nuclear region and does not reach distances beyond $\sim \text{few} \times 100 \text{ pc}$.

Alternatively, the higher density of the outflowing gas might be explained by the compression exerted on the outflowing gas by the radio-jet induced shocks (Villar Martín et al. 1999; Holt et al. 2011). We note that this scenario is qualitatively different from the scenario where the NLR clouds are compressed by radiation pressure, as assumed in the models used above. The higher density would instead be located at or near the radio jet and decrease moving away from it. In this case a correlation would be expected between the morphology of the radio jet and the ionized gas (whose emission is enhanced due to shock excitation and/or density enhancement). This is actually the case in MRK 477 (Heckman et al. 1997). Both the radio and [O III] images show a bright knot or ridge of emission about 0.4 arcsec ($\sim 290 \text{ pc}$) to the northeast of the central source. This clearly shows that the radio jet is interacting with the NLR and enhancing the emission at this location. Thus, an alternate scenario is that this detached knot is responsible for the bulk of emission of the outflowing gas. If so, it must emit a wide range of lines, including [FeVII] $\lambda 6087$. High spatial resolution spatially extended optical spectroscopy would help discriminate between both scenarios.

4.3 The origin of the NIR narrow [FeII] $\lambda 1.644$ emission

We have found that this line is much broader than expected for its critical density and ionization potentials (Sect. 3.2). The underlying broad component has $\text{FWHM} \sim 4770 \pm 830 \text{ km s}^{-1}$, larger than any other line and contributes more than half of the total line flux ($\sim 57\%$). It also shows the largest $\epsilon = 1.3 \pm 0.2$ and is the only line for which the broad component is not blueshifted. All these could be naturally explained in terms of reddening being much weaker in the NIR, so that both the approaching and receding parts of the expanding outflow are observed (unlike the optical lines). However, this is an unlikely explanation since $\text{Pa}\alpha$ shows that the broad component is blueshifted by $-570 \pm 119 \text{ km s}^{-1}$.

Alternatively, it is possible that [FeII] $\lambda 1.644$ is relatively enhanced by the shocks induced by the outflow, more than other lines and at a less obscured spatial location (hence the non-blueshift). Outflow induced shocks have been often proposed as a relevant [FeII] $\lambda 1.644$ excitation mechanism in active galaxies (e.g. Ramos Almeida et al. 2009, Contini et al. ?). As an example, ? studied the kinematics and excitation mechanism of the NIR emission lines in a sample of Seyfert 1 galaxies, including [FeII] $\lambda 1.257$ and $\lambda 1.645 \mu\text{m}$. They found 3 out of 22 sources in which the [FeII] lines were the broadest, even broader than the coronal [SiX] $\lambda 1.252 \mu\text{m}$. They suggest that in these objects [FeII] must arise from and additional source, partially formed in a region distinct from other low-ionization species, which they suggest to be associated with shock excitation from the radio jet. Blietz et al. (1995) also suggest that the [FeII] $\lambda 1.644$ emission in the Sy2

galaxy NGC1068 is preferentially emitted by gas which is interacting with the nuclear outflow/jet and is possibly located at the interface between the outflow and the dense circumnuclear molecular clouds.

4.4 Coronal lines

Detection of coronal emission is confirmed in the NIR spectrum, but not in the optical (Sect. 3.6). In the optical, the highest ionization lines are those emitted by Fe^{+6} , which on the other hand are fainter relative to the low ionization lines than typically found in AGN with strong coronal emission. Coronal lines are generally detected between just a few parsecs and a few hundred parsecs (e.g. Mazzalay et al. 2010;2014). They have been proposed to be formed in a region located at an intermediate distance between the classical NLR and the broad-line region (BLR) (Müller Sánchez et al. 2006). Alternatively, the coronal region might reside in the inner wall of the dusty torus (Murayama & Taniguchi 1998, Rose et al. 2013) or in the low density surface layer of the radiation pressure confined NLR clouds (S14).

The detection of coronal emission in the NIR and its absence in the optical spectrum of MRK 477 suggests that this region is heavily reddened, partially hidden from our line of sight. Alternatively, since the coronal lines are usually highly nucleated (<1 kpc from the nucleus) and much more concentrated than lower ionization NLR features (Rodríguez Ardila et al. 2011), it is also possible that they are diluted by the strong continuum contribution within the large aperture of the SDSS fibre (radius ~ 1 kpc).

The [SiVI] and [SiX] lines have luminosities 40.0 and 39.8 in log and erg s^{-1} respectively. Rodríguez Ardila et al. (2011) found that these lines display a narrow range in luminosity in Seyfert 1 and 2, with values for most objects located in the interval $\log(L)=39\text{--}40$. MRK 477 is at the high end of this range. We measure $[\text{SiVI}]/[\text{SiX}]=1.6\pm 0.1$, also within the range measured for Seyfert 1 and 2 galaxies by those authors.

4.5 Detection of narrow optical Fe^{+} emission

We have identified more than 10 *narrow optical* Fe^{+} emission lines in the SDSS spectrum of MRK 477. The NIR [FeII] lines at 1.26 and 1.64 μm are routinely detected in type 2 Seyferts (e.g. Ramos Almeida, Pérez García & Acosta-Pulido 2009). However, to our knowledge MRK 477 is the first type 2 AGN with optical Fe^{+} line detections.

To explain the absence of narrow optical Fe^{+} in type 2 AGN, while being prevalent in type 1 objects, Dong et al. (2010) proposed that such emission is confined to a disk-like geometry in the innermost region of the NLR on physical scales of parsecs. This would be smaller than the obscuring torus and within the dust sublimation radius. Iron, which is a refractory element which easily condenses on to dust grains, is in the gaseous phase in the absence of dust (Laor & Draine 1993). In this scenario the narrow optical Fe^{+} emitting region is visible along our line of sight in type 1 objects, but obscured by the extent of the dusty torus in type 2 counterparts.

It is however difficult to picture a geometry such as this to explain the optical Fe^{+} emission in MRK 477. Broad

permitted hydrogen and helium lines would be expected, unless a BLR did not exist, but this is not the case (Sect. 1).

We explore next whether the optical Fe^{+} features can be naturally explained by the intrinsic NLR emission. For this, we compare the measured integrated flux relative to $\text{H}\beta$ of all Fe^{+} lines in the blue (4000–6000 Å) and red (6000–7800 Å) bands with the ratios predicted by photoionization models appropriate for the NLR conditions. We use the radiation-pressure-confined NLR models described in S14. The assumption of these models implies that the line emission is essentially independent of the ionization parameter at the slab surface U_0 , as long as $U_0 \gg 0.03$ (see Fig. 3 in S14; see also Dopita et al. 2002 and Groves et al. 2004). A continuous distribution of dusty gas as a function of the distance from the nucleus R is considered. This distribution is characterized by η , the power-law index of the dusty gas covering factor as a function of logarithmic unit of R . These types of models are successful in explaining various observations of the NLR (see Dopita et al. 2002 and S14).

We assume solar metallicity, an ionization slope of $\alpha_{\text{ion}} = -1.6$ typical of luminous quasars (Telfer et al. 2002), and $\eta = 0$, i.e. a constant covering factor per log R , as suggested by the strong line ratios and the IR spectral energy distribution (S14). The photoionization models are calculated using CLOUDY (Ferland et al. 2013), assuming hydrostatic equilibrium. Depletion of refractory elements on to dust grains is taken into account, and the Fe^{+} ion is calculated according to the model described in Verner et al. (1999).

The results for the $\text{FeII}(4000 - 6000)/\text{H}\beta$ and $\text{FeII}(6000 - 7800)/\text{H}\beta$ are shown in Table 6 considering both dusty and non-dusty models. The uncertainties on the observed ratios account for the errors on the flux measurements and the possible misidentification of some features (Table 1). The models can reproduce successfully both the blue and the red bands. Therefore, the narrow optical Fe^{+} lines in MRK 477 can be naturally explained by the intrinsic NLR emission due to AGN photoionization. The high luminosity and/or closeness of MRK 477 could be the reason why these lines have been detected, unlike other type 2 AGN.

The effect of dust depletion produces ratios approximately twice lower than those derived from dust-free models. Taking into account the uncertainties, the Fe^{+} band ratios do not discriminate between the two scenarios. Model predictions of lines emitted by other refractory elements such as Mg, Ni, Ca (also detected for MRK 477, see Sect. 3.5) would help to infer whether the NLR contains dust-free gas. Only the [CaII] $\lambda 7291$ is predicted by CLOUDY. Calcium is much more sensitive to dust depletion than iron. The dust-free models predict $[\text{Ca II}]\lambda 7291/\text{H}\beta=0.08$, while the line would be 1000 times fainter (and thus completely undetectable) in the dusty case (see also Villar Martín et al. 1966). The [Ca II] line is tentatively detected with a ratio of 0.02 ± 0.01 relative to $\text{H}\beta$ (Fig. 12, third panel). If this detection is confirmed, it would imply that at least a fraction of the NLR gas is dust free.

| (1) $\left(\frac{\text{FeII}_{4000-6000}}{\text{H}\beta}\right)^{\text{obs}}$ | (2) $\left(\frac{\text{FeII}_{4000-6000}}{\text{H}\beta}\right)^{\text{int}}$ | (3) Model ₁ Dust | (4) Model ₂ Dust-free | (5) $\left(\frac{\text{FeII}_{6000-7800}}{\text{H}\beta}\right)^{\text{obs}}$ | (6) $\left(\frac{\text{FeII}_{6000-7800}}{\text{H}\beta}\right)^{\text{int}}$ | Model ₁ Dust | Model ₂ Dust-free |
|--|--|-----------------------------------|--|--|--|----------------------------|---------------------------------|
| 0.14±0.03 | 0.15±0.1 | 0.15 | 0.36 | 0.06±0.03 | 0.05±0.04 | 0.031 | 0.053 |

Table 6. Comparison between the measured integrated flux relative to H β of all Fe⁺ lines in the blue (4000–6000 Å) and red (6000–7800 Å) bands with the ratios predicted by photoionization models appropriate for the NLR conditions (see the text). The superscripts ^{obs} and ^{int} refer to the observed and reddening corrected ratios. Dusty and dust-free model predictions are shown. The narrow optical Fe⁺ lines in MRK 477 can be naturally explained by the intrinsic NLR emission due to AGN photoionization.

5 CONCLUSIONS

We perform a detailed spectral analysis and characterization of the NLR of the nearest obscured quasar (QSO2) MRK 477 at $z=0.037$ based on the SDSS optical and NIR H+K spectra obtained with WHT-LIRIS.

(i) We confirm the hybrid nature (AGN+starburst) proposed by Heckman et al. 1997, based on the new detection of the “red bump” at ~ 5800 Å due to WR stars, which implies that the system has undergone recent star formation.

(ii) The optical spectrum of MRK 477 is rich in emission lines, with ~ 90 detected features. In spite of the lack of spatial information of the SDSS spectrum, the spectral decomposition of numerous lines has allowed us to characterize, at least partially, the spatial structure and the gradients in the physical and kinematic properties of the gas.

(iii) Gas densities within the range $\sim 1945^{+670}_{-460}$ and up to $>10^4 \text{ cm}^{-3}$ are confirmed in the NLR of MRK 477.

(iv) As in many other active galaxies (AGN), a significant correlation is found between the lines FWHM and the critical density $\log(n_{\text{crit}})$. We propose that this correlation is caused by the outflow and its impact on the gas kinematics. This could be the case in other AGNs.

(v) MRK 477 is an example of a radio quiet powerful AGN where negative feedback (the nuclear outflow) can be dominated by the radio structures. The outflow emission has been isolated in many emission lines covering a large range of ionization potentials and critical densities (from [OI] $\lambda 6300$ to [FeVII] $\lambda 6086$). The outflowing gas, which is concentrated within $R \sim \text{few} \times 100$ pc from the central engine, is >13 times denser ($n \gtrsim 8,000 \text{ cm}^{-3}$) than the ambient non perturbed gas ($n \sim 400\text{--}630 \text{ cm}^{-3}$). Marginal evidence is found for densities as high as $n \gtrsim 5 \times 10^5 \text{ cm}^{-3}$ in the outflowing gas. It is possible that the density enhancement is due to the gas compression produced by jet induced shocks. This is supported by the correlation between the radio and [OIII] $\lambda 5007$ morphologies found by Heckman et al. (1997).

Alternatively, the density enhancement is not related to the jet. Instead, it might be a reflection of the NLR intrinsic density gradient, consequence of the gas being compressed by radiation pressure. In this scenario, and based on the comparative study between the density and the kinematic properties of the outflowing and the ambient gas, we conclude that the outflow has been generated at $\lesssim 220$ pc (possibly at $\lesssim 30$ pc) from the AGN. We find evidence of how its effects weaken as it propagates outwards, following the NLR density gradient. Beyond the radio jet edge, the gas emission is dominated by ambient less dense, non perturbed gas. This adds further support to the idea that the radio jet has triggered the outflow.

(vi) The [FeII] $\lambda 1.644 \mu\text{m}$ line presents a very different behaviour than the rest of the emission lines. It shows the most extreme effects of the outflow, with an underlying broad component of $\text{FWHM} = 4770 \pm 830 \text{ km s}^{-1}$. Its properties suggest that its emission is enhanced by shocks induced by the nuclear outflow/jet and is preferentially emitted at a different, less reddened spatial location, maybe the interface between the outflow and the dense circumnuclear molecular clouds.

(vii) More than 10 narrow optical Fe⁺ emission lines have been detected in the SDSS spectrum of MRK 477. To our knowledge, this is the first type 2 AGN with such a detection. We show that these lines can be explained as the natural emission from NLR gas photoionized by the AGN. Emission lines associated with other highly refractory elements (Mg, possibly Ni) are also detected. If the tentative detection of the [Ca II] $\lambda 7291$ line is confirmed, this would imply that at least part of the NLR gas is dust-free.

(viii) Coronal line emission is confirmed in the NIR, but not in the optical SDSS spectrum. The coronal region might be heavily reddened, partially hidden from our line of sight. Alternatively its optical emission might be diluted due to the large SDSS fibre aperture. The coronal region also participates in the outflow.

(ix) Pa α is spatially extended along the radio and [OIII] emission axes, up to a maximum radial extension of ~ 1.5 kpc from the AGN.

ACKNOWLEDGMENTS

Thanks to an anonymous referee for useful comments on the paper.

This work has been funded with support from the Spanish Ministerio de Economía y Competitividad through the grant AYA2012-32295. JS acknowledges financial support from the Alexander von Humboldt foundation. CRA is supported by a Marie Curie Intra European Fellowship within the 7th European Community Framework Programme (PIEF-GA-2012-327934). RGD acknowledges support through the grant AYA2010-15081.

This work is partially based on data obtained as part of the WHT service programme. The WHT and its service programme are operated on the island of La Palma by the Isaac Newton Group in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. We thank the WHT staff for performing the observations.

The work is also based on data from Sloan Digital Sky Survey. Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating

Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS website is <http://www.sdss.org/>.

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

REFERENCES

- Acosta Pulido J. et al. *ING Newsl.*, 7, 15
- Appenzeller I., Oestreich R., 1988, *AJ*, 95, 45
- Bennert N., Jungwiert B., Komossa S., Haas M., Chini R., 2006, *A&A*, 456, 953
- Bennert N., Jungwiert B., Komossa S., Haas M., Chini R., 2006, *A&A*, 459, 55
- Binette L., Wang J., Villar-Martín M., Martin P., Magris G., 1993, *ApJ*, 414, 535
- Blietz M., Cameron M., Drapatz S., Genzel R., Krabbe A., van der Werf P., Sternberg A., Ward M., 1994, *ApJ*, 421, 92
- De Robertis M.M., Ostrbrock D.E., 1984, *ApJ*, 286, 171, 953
- De Robertis M.M., Ostrbrock D.E., 1986, *ApJ*, 301, 727
- De Robertis M.M., 1987, *ApJ*, 316, 597
- Dong X. B., Ho L., Wang J.G., Want T.G., Wang H., Xiaohui F., Hongyan Z., 2010, *ApJL*, 721, L143
- Dopita M. A., Groves B. A., Sutherland R. S., Binette L., Cecil G., 2002, *ApJ*, 572, 753
- Dougados C., Bacciotti F., Cabrit S., Nisini B., 2010, in *Jets from Young Stars IV*, Lecture Notes in Physics, Volume 793. ISBN 978-3-642-02290-6. Springer-Verlag Berlin Heidelberg, 2010, p. 213
- Espey B., 1994, in Shlosman I., ed., *Proc. Conf. Mass-Transfer Induced Activity in Galaxies*. Cambridge Univ. Press, Cambridge, p.121
- Falcón-Barroso J., Ramos Almeida C., Böker T., Schinnerer E., Knapen J. H., Lancon A., Ryde, S., 2014, *MNRAS*, 438, 329
- Ferland G. J. et al., 2013, *Rev. Mex. Astron. Astrofis.*, 49, 137
- González Delgado R., Heckman T., Leitherer C., Meurer G., Krolik J., Wilson A., Kinney A., Koratkar A., 1998, *ApJ*, 505, 174
- Groves B. A., Dopita M. A., Sutherland R. S., 2004, *ApJS*, 153, 75
- Guseva N., Izotov Y., Thuan T., 2000, *ApJ*, 531, 776
- Heckman T. M., González Delgado R., Leitherer C., Meurer G.R., Krolik J., Wilson A.S., Koratkar A., Kinney A., 1997, *ApJ*, 482, 114
- Holt J., Tadhunter C. N., Morganti R., Emonts B., 2011, *MNRAS*, 410, 1527
- Keenan F. P., Aller L. H., Bell K. L., Hyung S., McKenna F. C., Ramsbottom C. A., 1996, *MNRAS*, 271, 1073
- Koss M., Mushotzky R., Treister E., Veilleux S., Vasudevan R., Tripp M., 2012, *ApJ*, 746, 22
- Krug H., Rupke D., Veilleux S., 2010, *ApJ*, 708, 1145
- Lamastra A., Bianchi S., Matt G., Perola G.C., Barcons X., Carrera F.J., 2009, *A&A*, 504, 73
- Laor A., Draine B., 1993, *ApJ*, 402, 441
- Luridiana V., Morisset C., Shaw R., 2015, *A&A*, 573, 42
- Manchado A., et al. 2004, *Proc. SPIE*, 5492, 1094
- Mazzalay X., Rodríguez-Ardila A., Komossa S., 2010, *MNRAS*, 405, 1315
- Mazzalay X. et al. 2013, *MNRAS*, 428, 2389
- Mazzalay X. et al., 2014, *MNRAS*, 438, 2036
- Müller Sánchez F., Davies R. I., Eisenhauer F., Tacconi L. J., Genzel R., Sternberg A., 2006, *A&A*, 454, 481
- Müller Sánchez F., Prieto M. A., Hicks E. K. S., Vives-Arias H., Davies R. I., Malkan M., Tacconi L. J., Genzel R., 2011, *ApJ*, 739, 69
- Murayama T., Taniguchi Y., 1998, *ApJ*, 503, L115
- Osterbrock D.E., 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei*. University Science Books, Mill Valley, CA
- Penston M.V., Fosbury R.A.E., Boksenberg A., Ward M.J., Wilson A.S., 1984, 208, 347
- Pier E., Voit G.M., 1994, *ApJ*, 450, 628
- Ramos Almeida C., Prez Garca A. M., Acosta-Pulido J. A., 2009, *AJ*, 137, 179
- Rodríguez Ardila A., Prieto M.A., Viegas S., Grwenland R., 2006, *ApJ*, 653, 109
- Rodríguez Ardila A., Pastoriza M. G., Viegas S., Sigut T. A., Pradhan A. K., 2004, *A&A*, 425, 457
- Rodríguez Ardila A., Prieto M.A., Portilla J.G., Tejeiro J.M., 2011, *ApJ*, 743, 100
- Rose M., Tadhunter C. N., Holt J., Rodríguez Zaurín J., 2013, *MNRAS*, 432, 2150
- Schaerer D., Vacca W. D. W., 1998, *ApJ*, 497, 618
- Shen Y. et al., 2011, *ApJS*, 194, 45
- Shu X., Wang J., Jiang P., Fan L., Wang T., 2007, *ApJ*, 657, 167
- Shuder J., Osterbrock D., 1981, *ApJ*, 250, 55
- Stern J., Laor A., 2012, *MNRAS*, 426, 2703
- Stern J., Laor A., Baskin A., 2014, *MNRAS*, 438, 901 (S14)
- Telfer R. C., Zheng W., Kriss G. A., Davidsen A. F., 2002, *ApJ*, 565, 773
- Tran H., Miller J., Kay L., 1992, *ApJ*, 397, 452
- Tran H., 1995, *ApJ*, 440, 578
- Veilleux S., 1988, *AJ*, 95, 1695
- Verón-Cetty M.P., Joly M., Véron P., 2004, *A&A*, 417, 515
- Verón-Cetty M.P., Véron P., Joly M., Kollatschny W., 2013, *A&A*, 475, 487
- Verner E. M., Verner D. A., Korista K. T., Ferguson J. W., Hamann F., Ferland G. J., 1999, *ApJS*, 120, 101
- Villar Martín M., Binette L., 1996, *A&A*, 309, 97
- Villar Martín M., Tadhunter C., Morganti R., Axon D., Koekemoer A., 1999, *MNRAS*, 307, 24

Villar Martín M., Emonts B., Humphrey A., Cabrera
Lavers A., Binette L., 2014, MNRAS, 440, 3202
Whittle M., 1992, ApJS, 79, 49
Wright E. L., 2006, PASP, 118, 1711
York D. G. et al., 2000, AJ, 120, 1579
Zakamska N. L. et al. 2003, AJ, 126, 2125
Zhang S.H., Bian W.H., Huang K.L., 2008, MNRAS, 488,
113
Zwicky F., 1966, ApJ, 143, 192