

## Lab Resource: Stem Cell Line

## Generation of a human iPSC line, IISHDOI002-A, with a 46, XY/47, XYY mosaicism and belonging to an African mitochondrial haplogroup



María del Carmen Ortuño-Costela<sup>a,b,c,d</sup>, Ana Moreno-Izquierdo<sup>d,e</sup>,  
Rafael Garesse<sup>a,b,c,d</sup>, M. Esther Gallardo<sup>a,b,c,d,\*</sup>

<sup>a</sup> Departamento de Bioquímica, Facultad de Medicina, Universidad Autónoma de Madrid, Spain

<sup>b</sup> Instituto de Investigaciones Biomédicas “Alberto Sols”, UAM-CSIC, Madrid, Spain

<sup>c</sup> Centro de Investigación Biomédica en Red (CIBERER), Madrid, Spain

<sup>d</sup> Instituto de Investigación Sanitaria, Hospital 12 de Octubre (i+12), Madrid, Spain

<sup>e</sup> Servicio de Genética, Hospital 12 de Octubre, Madrid, Spain

## ARTICLE INFO

## Article history:

Received 7 November 2017

Received in revised form 7 February 2018

Accepted 11 February 2018

Available online 14 February 2018

## ABSTRACT

We have generated a human iPSC line, IISHDOI002-A, from commercial primary normal human dermal fibroblasts belonging to an African mitochondrial haplogroup (L3), and with a 46, XY/47, XYY mosaicism. For this purpose, reprogramming factors Oct3/4, Sox2, Klf4 and cMyc were delivered using a non-integrative methodology that involves the use of Sendai virus.

© 2018 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Resource table.

Unique stem cell line identifier	IISHDOI002-A
Alternative name(s) of stem cell line	FCE-FiPS4F-7
Institution	Instituto de Investigación Sanitaria Hospital 12 de Octubre, i + 12
Contact information of distributor	Dr. M. Esther Gallardo <a href="mailto:egallardo.imas12@h12o.es">egallardo.imas12@h12o.es</a>
Type of cell line	iPSC
Origin	Human
Additional origin info	Sex: Male
Cell source	Normal human neonatal dermal fibroblasts (Lonza, CC2509)
Clonality	Mixed
Method of reprogramming	Sendai virus
Genetic modification	NO
Type of modification	N/A
Associated disease	N/A
Gene/locus	N/A
Method of modification	N/A
Name of transgene or resistance	N/A
Inducible/constitutive system	N/A
Date archived/stock date	May 2017
Cell line repository/bank	N/A

## Ethical approval

Fibroblasts were obtained after signing informed consent. This study was reviewed and approved by the Institutional Ethical Committee of the “Instituto de Investigaciones Biomédicas Alberto Sols”, CSIC-UAM, 406329 1.

## Resource utility

Functional studies using iPSCs generated from patients with mitochondrial diseases (MD) require the use of the most appropriate controls. An isogenic control is not always possible. The availability of the iPSC line, IISHDOI002-A, will be very useful for functional studies of MD patients belonging to a L3 haplogroup.

## Resource details

The establishment of a human iPSC line, IISHDOI002-A, was performed using non-integrative methodology that involves the use of Sendai viruses containing the reprogramming factors Oct3/4, Sox2, Klf4 and c-Myc (Takahashi et al., 2007). For this purpose, commercially available fibroblasts from a healthy neonatal donor were acquired in Lonza (CC2509). Subsequently, mitochondrial haplogroup determination was carried out in the IISHDOI002-A line and in the original fibroblasts by PCR amplification of mitochondrial DNA (mtDNA) regions containing the nucleotide positions that define the L3 haplogroup (m.10400 and m.3594) followed by direct sequencing of the generated amplicons (Table 1). The identification of the mtDNA nucleotide variations m.10400C and m.3594 T showed that fibroblasts and the generated iPSC line IISHDOI002-A belong to

\* Corresponding author.

E-mail address: [egallardo.imas12@h12o.es](mailto:egallardo.imas12@h12o.es) (M.E. Gallardo).

the African mitochondrial haplogroup L3 (Fig. 1A). IISHDOI002-A iPSC colonies displayed a typical ES-like colony morphology and growth behavior (Fig. 1B) and they stained positive for alkaline phosphatase (AP) activity (Fig. 1C). The endogenous expression of the pluripotency associated transcription factors *OCT4*, *SOX2*, *KLF4*, *NANOG*, *CRIPTO* and *REX1* was evaluated by quantitative real time polymerase chain reaction (qPCR) (Fig. 1D). Immunofluorescence analysis revealed expression of the pluripotency-associated transcription factors *OCT4*, *SOX2* and *NANOG*, and surface markers SSEA3, SSEA4, TRA1-81 and TRA1-60 (Fig. 1E). We also confirmed the clearance of the vectors and the exogenous reprogramming factor genes by RT-PCR after nine culture passages (Fig. 1F). The iPSC line has been adapted to feeder-free culture conditions and a karyotype analysis after more than 20 culture passages has been carried out. This analysis displayed a 46, XY (28)/47, XYY (2) mosaicism in the iPSC line (Fig. 1G). We also confirmed the presence of abnormal cells at 4–4.5% in two different aliquots of the original fibroblasts by FISH analysis of 200 interphase cells [nuc ish (DYZ1x2, DYZ1x1) [9/200] (Fig. 1G). These data suggest that this mosaicism could be a causal finding in the fibroblasts and have a somatic origin. The level of mosaicism in the fibroblasts has been maintained around 4.5% during more than 30 passages. In any case, males with an extra copy of the Y chromosome typically have no unusual physical features although they could have fertility problems. Regarding the case reported here, the identified mosaicism, to our knowledge, is not related with any noteworthy phenotypic trait present in the carrier. We also verified by DNA fingerprinting analysis that the line IISHDOI002-A was derived from the patient's fibroblasts (Supplementary Fig. S1). In addition, the line was confirmed by PCR analysis to be mycoplasma-negative (Fig. 1H). Finally, the capacity of the IISHDOI002-A iPSC line to differentiate into the three germ layers (endoderm, mesoderm and ectoderm) was evaluated *in vitro* using an embryoid body based assay (Fig. 1I).

## Materials and methods

### iPSCs generation

Healthy human neonatal fibroblasts (Lonza, CC2509) were reprogrammed using the CytoTune-iPS 2.0 Sendai reprogramming kit following the instructions of the manufacturer. IISHDOI002-A was expanded as described in Galera et al., 2016.

### AP analysis

The iPSC line was seeded on a feeder layer plate. After one week, direct AP activity was determined using the AP blue membrane substrate solution kit (Sigma, AB0300).

### Mitochondrial haplogroup

DNAs from fibroblasts and IISHDOI002-A were haplogrouped by PCR and direct sequencing, using the primers described in Table 2. The mtDNA nucleotides, m.10400C and m.3594 T, defining the African haplogroup L3, have been verified.

### qPCR analysis

Total mRNA was isolated using TRIZOL and cDNA using the Quantitect RT cDNA synthesis kit. The expression of the endogenous pluripotency associated genes (*OCT4*, *SOX2*, *KLF4*, *NANOG*, *CRIPTO* and *REX1*) was quantified by qPCR. Primers are listed in Table 2 (Aasen et al., 2008). All the expression values were normalized to the *GAPDH* gene. Plots are representative of at least three independent experiments.

### Karyotype and FISH


Cells were treated with 10 µg/mL of Colcemid (Gibco) for 90 min at 37 °C, trypsinized, treated with KCl 0.075 M, and fixed with Carnoy's fixative for karyotype analysis and fixed three times in methanol/acetic acid (3:1) for FISH. Cells were then dropped on a microscope glass slide and dried. Metaphase cells were G banded using Wright staining. At least 30 metaphases were karyotyped. For FISH, CEP probe for chromosomes Y (DYZ1) and slides were denatured at 72 °C for 2 min and incubated at 37 °C (16 h) for hybridization. Slides were washed in SSC with 0.1% Tween 20. Coverslips were mounted on the slides with DAPI. 200 interphase cells were analyzed with a Nikon fluorescent microscope. Digital images were acquired with a monochrome CCD camera linked to Metasystem software.

### Immunofluorescence analysis

Cells were grown on 0.1% gelatin-coated 35 mm culture plates, fixed with 4% paraformaldehyde 30 min at RT and permeabilized using TBS+

**Table 1**  
Characterization and validation.

Classification	Test	Result	Data
Morphology	Photography	Normal	Fig. 1, panel B
Phenotype	Qualitative analysis: immunocytochemistry	Positive for the pluripotency markers: SSEA3, SSEA4, TRA-1-81, TRA-1-60, OCT4, NANOG, SOX2	Fig. 1, panel E
	Qualitative analysis: alkaline phosphatase activity	Positive	Fig. 1, panel C
	Quantitative analysis: gene expression (qPCR)	Positive for the pluripotency markers <i>OCT4</i> , <i>KLF4</i> , <i>SOX2</i> , <i>CRIPTO</i> , <i>NANOG</i> , <i>REX1</i>	Fig. 1, panel D
Genotype	Karyotype (G-banding) and resolution	46, XY (28)/47, XYY (2) Resolution 450–500	Fig. 1, panel G
Identity	Microsatellite PCR (mPCR)	N/A	
	STR analysis	8 loci, all matched (D2S1338, D7S820, D8S1179, D13S317, D19S433, D21S11, VWA, amelogenin)	Supplementary Fig. S1
Mutation analysis (if applicable)	Sequencing	N/A	
Haplogroup analysis	Southern Blot OR WGS	N/A	
Microbiology and virology	Sequencing	Confirmation of the L3 haplogroup: m.10400C and m.3594 T	Fig. 1, panel A
	Mycoplasma	Negative	Fig. 1, panel H
	Sendai virus silencing	Virus silenced	Fig. 1, panel F
Differentiation potential	Embryoid body formation and directed differentiation	Positive for: smooth muscle actin (SMA), $\beta$ -tubulin (Tuj1) and alpha-fetoprotein (AFP)	Fig. 1, panel I
Donor screening (optional)	HIV 1+2 Hepatitis B, Hepatitis C	N/A	
Genotype additional info (optional)	Blood group genotyping	N/A	
	HLA tissue typing	N/A	



**Fig. 1.** Molecular and functional characterization of the IISHDOI002-A iPSC line.

(0.1% Triton X-100 in Tris-buffered saline, TBS) for 45 min. Then the cells were incubated in TBS++ (3% donkey serum, 0.3% Triton X-100 in TBS) for 2 h at RT. Primary antibodies were applied overnight at 4 °C. Secondary antibodies, 2 h at RT. Nuclei were stained with DAPI (Sigma, 28718-90-3). All the antibodies are listed in Table 2.

#### *In vitro* differentiation assay

The *in vitro* pluripotency capacity of the line IISHDOI002-A was tested by spontaneous embryoid body differentiation. The protocol we have used has been described in detail by Galera et al. (2016).

#### DNA fingerprinting

The markers D13S317, D7S820, VWA, D8S1179, D21S11, D19S433, D2S1338 and amelogenin for sex determination were amplified by PCR and analyzed by ABI PRISM 3100 Genetic analyzer and Peak Scanner v3.5 (ThermoFisher) (Table 2).

#### *Mycoplasma* detection

Mycoplasma detection was performed by PCR (primers specified in Table 2) using 1 mL of the cell culture supernatant (3 days culture at 90% confluence). The 300 bp band represents that the sample is positive for mycoplasma. The band at 570 bp is an internal control to discard the inhibition of the polymerase.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scr.2018.02.009>.

#### Acknowledgments

We are very grateful to Marta García López and Nathalie Rodríguez Mancera for their help with iPS cell generation and characterization. This work was supported by grants from: 1) the “Fondo de Investigación Sanitaria, Instituto de Salud Carlos III”: PI10/0703, PI13/00556 and PI16/00789 to RG and PI15/00484 to MEG cofunded by FEDER, 2) from the “Centro de Investigación Biomédica en Red en Enfermedades Raras” (CIBERER): grants 13-717/132.05 and

**Table 2**  
Reagents details.

Antibodies used for immunocytochemistry/flow-cytometry			
	Antibody	Dilution	Company Cat # and RRID
Pluripotency markers	Mouse anti-TRA-1-81	1:150	Millipore Cat# MAB4381, RRID: <a href="#">AB_177638</a>
	Mouse anti-TRA-1-60	1:150	Millipore Cat# MAB4360, RRID: <a href="#">AB_11211864</a>
	Rabbit anti-SOX2	1:100	Thermo Fisher Scientific Cat# PA1-16968, RRID: <a href="#">AB_2195781</a>
	Mouse anti-SSEA4	1:10	Millipore Cat# MAB4304, RRID: <a href="#">AB_177629</a>
	Rat anti-SSEA3	1:20	Abcam Cat# ab16286, RRID: <a href="#">AB_882700</a>
	Goat anti-NANOG	1:25	R and D Systems Cat# sc-5279, RRID: <a href="#">AB_628051</a>
	Mouse anti-OCT4	1:100	Santa Cruz Biotechnology Cat# sc-5279, RRID: <a href="#">AB_628051</a>
Differentiation markers	Mouse anti-β3 tubulin isotype III	1:300	Sigma-Aldrich Cat# T8660, RRID: <a href="#">AB_528427</a>
	Mouse anti-AFP	1:300	Sigma-Aldrich Cat# WH0000174M1, RRID: <a href="#">AB_1839587</a>
	Mouse anti-SMA	1:400	Sigma-Aldrich Cat# A2547, RRID: <a href="#">AB_476701</a>
Secondary antibodies	Cy <sup>TM</sup> 2-conjugated AffiniPure Donkey Anti-Goat IgG (H+L)	1:50	Jackson ImmunoResearch Labs Cat# 705-225-147, RRID: <a href="#">AB_2307341</a>
	Cy <sup>TM</sup> 2-conjugated AffiniPure Goat Anti-Mouse IgG, Fcγ Subclass 2b specific	1:50	Jackson ImmunoResearch Labs Cat# 115-225-207, RRID: <a href="#">AB_2338749</a>
	Cy <sup>TM</sup> 2-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L)	1:50	Jackson ImmunoResearch Labs Cat# 111-225-144, RRID: <a href="#">AB_2338021</a>
	Cy <sup>TM</sup> 3-conjugated AffiniPure Goat Anti-Rat IgM, μ chain specific	1:250	Jackson ImmunoResearch Labs Cat# 112-165-075, RRID: <a href="#">AB_2338249</a>
	Cy <sup>TM</sup> 3-conjugated AffiniPure Goat Anti-Mouse IgG, Fcγ Subclass 3 specific	1:250	Jackson ImmunoResearch Labs Cat# 115-165-209, RRID: <a href="#">AB_2338698</a>
	Cy <sup>TM</sup> 3-conjugated AffiniPure Donkey Anti-Mouse IgM, μ chain specific	1:250	Jackson ImmunoResearch Labs Cat# 715-165-020, RRID: <a href="#">AB_2340811</a>
	Goat anti-mouse IgG (H+L), Alexa Fluor 488	1:500	Thermo Fisher Scientific Cat# A-11029, RRID: <a href="#">AB_2534088</a>
Primers			
	Target	Forward/reverse primer (5'-3')	
Pluripotency markers (qPCR)	<i>Endo-KLF4</i>	AGCCTAAATGATGGTGCTTGGT/TTGAAAACCTTGGCTTCCTTGT	
	<i>Endo-OCT4</i>	GGGTTTTTGGGATTAAGTTCTTCA/GCCCCACCCCTTTGTGT	
	<i>Endo-SOX2</i>	CAAAAATGGCCATGCAGGT/AGTTGGGATCGAACAAAAGCTATT	
	<i>REX1</i>	CCTGCAGGCGAAATAGAAC/GCACACATAGCCATCACATAAGG	
	<i>CRIP1</i>	CGGAAGTGTGAGCAGCATGT/GGGCAGCCAGGTGTCATG	
House-Keeping Genes (qPCR)	<i>NANOG</i>	ACAAGTGGCCGAAGAATAGCA/GGTTCCAGTCGGGTTTAC	
	<i>GAPDH</i>	GCACCGTCAAGGCTGAGAAC/AGGGATCTCGCTCTCGAA	
Haplogroup analysis	mtDNA region: from m.9977 to m.10858	TCTCCATCTATTGATGAGGGTCT/CAACACCCACAGCCTAATT	
	mtDNA region: from m.3150 to m.3980	TACTTCAAAAAGCGCTTCC/CCCTTCGCCTATTCTTCAT	
	SeV	GGATCACTAGGTGATATCGAGC/ACCAGACAAGAGTTTAAGAGATATGTATC	
Virus silencing	KOS	ATGCACCGCTACGAGCTGAGCG/ACCTTGACAATCCTGATGTGG	
	Klf4	TTCCTGCATGCCAGAGGAGCCC/AATGTATCGAAGGTGCTCAA	
STR analysis	c-Myc	TAACTGACTAGCAGGCTTGTG/TCCACATACAGTCTGGATGATGATG	
	D2S1338	[6-FAM] CCACTGGATTGGAAACAGA/ACCTAGCATGGTACCTGCAG	
	D7S820	[6-FAM] TGTTCATAGTTTGAACGAACAAACG/CTGAGGTATCAAAAACCTCAGAG	
	D8S1179	[6-FAM] TTTTGTATTTCATGTGTACATTCG/CTAGCTATAATTAGTTCATTTTCA	
	D13S317	[6-FAM] ACAGAAAGTCTGGGATGTGGA/GCCCAAAAAGACAGACAGAA	
Mycoplasma detection	D19S433	[6-FAM] CCTGGGCAACAGAAATAGAT/TAGGTTTTTAAGGAACAGGTGG	
	D21S11	[6-FAM] GTGAGTCAATCCCCAAG/GTTGTATTAGTCAATGTTCTCC	
	VWA	[6-FAM] CCCTAGTGGATGATAAGAATAATC/GGACAGATGATAAATACATAGGATGGATGG	
	Amelogenin	[6-FAM] CCCTGGGCTCTGTAAAGAATAGT/ATCAGAGCTTAACTGGGAAGCTG	
	GPO-3/MGSO	GGGAGCAACAGGATTAGATACCT/TGCACCATCTGTCACTCTGTAACTCTC	



ER16P3AC717 to RG and 3) “Comunidad Autónoma de Madrid” (grant number S2010/BMD-2402 to RG). MDCOC receives grant support from the Ministerio de Educación, Cultura y Deporte, grant number FPU16/03895 and MEG is supported by the “Miguel Servet Program” (CP16/00046) from Instituto de Salud Carlos III (Spanish Ministry of Health).

#### Author disclosure statement

There are no competing financial interests in this study.

#### References

- Aasen, T., Raya, A., Barrero, M.J., Garreta, E., Consiglio, A., Gonzalez, F., Vassena, R., Bilić, J., Pekarik, V., Tiscornia, G., Edel, M., Boué, S., Izpisua Belmonte, J.C., 2008. [Efficient and rapid generation of induced pluripotent stem cells from human keratinocytes. Nat. Biotechnol.](#) 26 (11), 1276–1284.
- Galera, T., Zurita, F., González-Páramos, C., Moreno-Izquierdo, A., Fraga, M.F., Fernández, A.F., Garesse, R., Gallardo, M.E., 2016. [Generation of a human iPSC line from a patient with Leigh syndrome. Stem Cell Res.](#) 16 (1), 63–66.
- Takahashi, K., Tanabe, K., Ohnuki, M., Narita, M., Ichisaka, T., Tomoda, K., Yamanaka, S., 2007. [Induction of pluripotent stem cells from adult human fibroblasts by defined factors. Cell](#) 131 (5), 861–872.