

## Short communication

Social avoidance and altered hypothalamic-pituitary-adrenal axis in a mouse model of anxious depression: The role of LPA<sub>1</sub> receptor

R.D. Moreno-Fernández<sup>a,\*</sup>, P. Sampedro-Piquero<sup>b</sup>, F.J. Gómez-Salas<sup>c</sup>, A. Nieto-Quero<sup>c,d</sup>,  
G. Estivill-Torrús<sup>d</sup>, F. Rodríguez de Fonseca<sup>d,e</sup>, L.J. Santín<sup>c,d</sup>, C. Pedraza<sup>c,d,\*\*</sup>

<sup>a</sup> Facultad de Educación y Psicología. Universidad Francisco de Vitoria, Spain

<sup>b</sup> Departamento de Psicología Biológica y de la Salud. Facultad de Psicología. Universidad Autónoma de Madrid. Madrid, Spain

<sup>c</sup> Departamento de Psicobiología y Metodología en las CC, Instituto de Investigación Biomédica de Málaga (IBIMA), Universidad de Málaga, Málaga, Spain

<sup>d</sup> Instituto de Investigación Biomédica de Málaga y Plataforma en Nanomedicina (IBIMA Plataforma BIONAND), Málaga, Spain

<sup>e</sup> Unidad Clínica de Neurociencias, Hospital Regional Universitario de Málaga, Spain

## ARTICLE INFO

## Keywords:

LPA<sub>1</sub> receptor  
HPA-axis  
Social preference  
Dexamethasone  
Corticosterone  
Depression

## ABSTRACT

Anxious depression is a prevalent disease with devastating consequences. Despite the lack of knowledge about the neurobiological basis of this subtype of depression, recently our group has identified a relationship between the LPA<sub>1</sub> receptor, one of the six characterized G protein-coupled receptors (LPA<sub>1-6</sub>) for lysophosphatidic acid, with a mixed depressive-anxiety phenotype. Dysfunctional social behaviors, which have been related to increased activation of the hypothalamus-pituitary-adrenal (HPA) axis, are key symptoms of depression and are even more prominent in patients with comorbid anxiety and depressive disorders. Social behavior and HPA functioning were assessed in animals lacking the LPA<sub>1</sub> receptor. For these purposes, we first examined social behaviors in wild-type and LPA<sub>1</sub> receptor-null mice. In addition, a dexamethasone (DEX) suppression test was carried out. *maLPA<sub>1</sub>*-null mice exhibited social avoidance, a blunted response to DEX administration and an impaired circadian rhythm of corticosterone levels, which are features that are consistently dysregulated in many mental illnesses including anxious depression. Here, we have strengthened the previous experimental evidence for *maLPA<sub>1</sub>*-null mice to represent a good animal model of anxious depression, providing an opportunity to explore new therapeutic targets for the treatment of mood disorders, particularly this subtype of depression.

## 1. Introduction

Depression is a major health concern in modern societies. Major depressive disorder is a heterogeneous illness with various subtypes. One of these subtypes is depression disorder with comorbid anxiety, also known as anxious depression, that involves more disability, more resistance to treatment, greater risk of suicide and is associated with more severe psychological, physical and social impairments than either condition alone [1]. The association between depression and impaired social functioning is well recognized. Social withdrawal and lack of social support are key features involved in both the onset and maintenance of the depressive state and can be a relevant target for intervention [2]. Dysfunctional social behaviors are even more prominent in patients with comorbid anxiety and depressive disorders than either

disease alone [3].

From a neurobiological point of view, social avoidance behavior has been related to increased activation of the hypothalamus-pituitary-adrenal (HPA) axis [4]. In this sense, relatively high HPA-axis activity has been related to increases in behavioral inhibition and increased social avoidance. In fact, there is a direct link between increased cortisol stress-responsiveness and social avoidance behavior [4]. Because HPA-axis dysfunction is a well-known biological marker of depression, its function in depressed patients has been extensively studied using a sensitive pharmacological challenge test, the dexamethasone (DEX) suppression test (DST) [5]. Although there are conflicting results, DEX nonsuppression has been linked to depression. Most likely, the explanation for the conflict lies in the existence of different subtypes of this disease. Patients with anxious depression show DST nonsuppression

\* Corresponding author.

\*\* Corresponding author at: Departamento de Psicobiología y Metodología en las CC, Instituto de Investigación Biomédica de Málaga (IBIMA), Universidad de Málaga, Málaga, Spain.

E-mail addresses: [romandario.moreno@ufv.es](mailto:romandario.moreno@ufv.es) (R.D. Moreno-Fernández), [mdpedraza@uma.es](mailto:mdpedraza@uma.es) (C. Pedraza).

<https://doi.org/10.1016/j.bbr.2023.114681>

Received 21 June 2023; Received in revised form 3 September 2023; Accepted 20 September 2023


Available online 21 September 2023

0166-4328/© 2023 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

rates that indicate a dysregulation in the HPA-axis [6].

Recently, our group has been the first to propose a possible relationship between the LPA<sub>1</sub> receptor, one of the six G protein-coupled receptors through which lysophosphatidic acid (LPA, 1-acyl-2-sn-glycerol-3-phosphate) acts, and the mixed depressive-anxiety phenotype, which may shed light on the unknown neurobiological basis of this

subtype of depression [7,8]. In fact, based on the validity criteria and considering the results of accumulated data with studies using animals lacking the LPA<sub>1</sub> receptor, maLPA<sub>1</sub>-null mice have been proposed as an animal model of the mixed depressive-anxiety phenotype [7]. Moreover, absence of the LPA<sub>1</sub> receptor induces exaggerated endocrine responses to emotional stimuli [8,9], causes hypoactivity and impairs adaptation



**Fig. 1.** Assessment of Social behavior revealed dysfunctional social behavior in the absence of the LPA<sub>1</sub> receptor. (A) The amount of time (the mean  $\pm$  S.E.M) or (B) a representative picture of time mice spent in each of the three chambers revealed social avoidance in the null mice. (C) Data are corroborated by the number of entries into each chamber. (D) Duration (s) of grooming, (E) freezing and (F) rearing in each compartment. (G) Percentage of time in relation to the total time spent in the 'social compartment' by each genotype sniffing the juvenile mice, rearing, grooming or freezing. (H) The distance traveled and (I) the velocity measured during the three compartments test revealed no differences between the genotypes. (J) Principal component analysis of social behavior measures. Interpretable factor loadings are in bold. The results revealed evidence both for a factor relating to social behavior and for another factor that relates to exploratory behavior. (K) Variables with negative scores are inversely related to the factor; therefore, absence of the LPA<sub>1</sub> receptor is negatively related with social behavior. Rotation method: Varimax with Kaiser normalization. Rotation converged in 3 iterations. KMO = 0.51;  $\chi^2 = 34.69$ ;  $p < 0.001$ . (L) Scree plot of the principal components analysis. Data are represented as the mean  $\pm$  standard error of the mean (S.E.M). \* $p < 0.05$ , \*\* $p < 0.01$  and \*\*\* $p < 0.001$  significant differences between genotypes. # $p < 0.05$  significant difference with the central compartment.  $s$   $p < 0.05$  significant difference with social compartment.  $n = 7$  wild-type mice and 7 maLPA<sub>1</sub>-null mice.

of the HPA-axis after chronic stress [10], which are factors that have also been strongly correlated with depression [11]. While suggestive of a dysfunctional HPA-axis, this issue has not yet been systematically evaluated in *maLPA<sub>1</sub>*-null mice. Thereby, the aims of this study were to elucidate the involvement of the *LPA<sub>1</sub>* receptor in two separate depression related characteristics: HPA-axis functioning and in social behavior. For this purpose, social behavior was evaluated using the three compartments test in wild-type and *maLPA<sub>1</sub>*-null mice. Moreover, in both genotypes, DST was carried out in order to measure the corticosterone response.

## 2. Methods

### 2.1. Animals

For this study we used the Malaga variant of null mice for the *LPA<sub>1</sub>* receptor, termed as "*maLPA<sub>1</sub>*-null" [12], derived from the original colony of Contos et al. [13]. Except for 1-month-old juvenile mice used for the Social Preference Test, 3-month-old male mice were used. For the Social Preference Test 14 mice were used. An additional group of 15 animals were subjected to the DST (Figs. 1 and 2).

Animals were housed in groups of 4 in a cycle of 12 h light/dark (lights on at 07:00 h) with water and food available *ad libitum*. The experiments were carried out between 9:00 and 15:00 h.

The procedures were approved by the Ethics Committee of the University of Malaga (CEUMA: 1-2015-A, 08-7-15-273) and complied with the European Animal Research Laws (EU Directive 2010/63/EU, EU, 90/219 / EEC and Regulation [EC] N° 194/2003) and the Spanish Guidelines on Animal Experimentation and Use of Genetically Modified Organisms (Royal Decrees: 53/2013, 178/2004; Laws 32/2007, 9/2003 and Decree 320/2010).

### 2.2. Social preference test

The social preference test was performed in a rectangular box formed by three chambers (LE894 Panlab Harvard Apparatus, Spain). The left and right compartments were equipped with wire cups that contained either an unfamiliar male juvenile mouse (the 'social compartment') or an unfamiliar object. The time spent in each chamber, locomotion (distance travelled [cm]) and velocity (cm/s) were registered using a video tracking system (Ethovision XT, Noldus, Wageningen, The Netherlands), whereas ethological parameters (time sniffing the juvenile [social target], freezing, rearing or grooming) were scored by two trained observers who were blind to the genotype of the animals.

### 2.3. Dexamethasone suppression test and corticosterone assay

For plasma corticosterone level determination, two samples of blood were collected from the lateral tail vein at 09:00 and 15:00 h. Immediately after the first blood collection, mice were administered IP vehicle

(saline) or 0.1 mg/kg DEX (Sigma Aldrich, D0720000).

The tubes containing the blood samples in presence of EDTA were centrifuged, and the supernatant was stored at  $-80^{\circ}\text{C}$ . Plasma corticosterone levels were determined in duplicate using a commercially available Enzyme Immunoassay Kit (sensitivity ca. 27.0 pg/ml) following the manufacturer's instructions (Assay Designs/Stressgen, Ann Arbor, Michigan, USA).

### 2.4. Statistical analysis

Data are expressed as the mean  $\pm$  standard error of the mean (S.E. M.). Comparisons between groups (wild-type mice vs *maLPA<sub>1</sub>*-null mice) in social behavior variables (time spent in each chamber, sniffing, rearing, freezing and grooming) were analyzed by Student's t-test for independent samples together with effect size Cohen's d. Moreover, we also performed t-test for dependent samples to compare the time spent for each group in the different compartments of the maze (centre, social and inanimate). Power analysis ( $1-\beta$ ) was also performed to control the probability of Type 1 error in our analysis. Furthermore, to determine the relationship between behavioral variables in the social chamber and to reduce them to a smaller set of dimensions that could underlie the genotype effects [14], a principal components analysis (PCA) was carried out. PCA was tested for sampling adequacy by the Bartlett sphericity and the Kaiser-Meyer-Olking (KMO) tests and followed by a varimax orthogonal rotation, which ensures that the extracted factors are independent of one another. The resulting factors with an eigenvalue  $> 1$  were selected, and the contribution of each variable to a factor (i.e., 'factor loading') was considered significant when it was more than 0.60 in absolute value.

For corticosterone measures, an ANCOVA using the basal CORT levels (measured at 9:00 h) as the covariable was utilized, thus controlling the different basal levels between genotypes.


Only probabilities  $\leq 0.05$  were considered significant. For the benefit of clarity and brevity, only relevant results of these statistical analyses are reported.

## 3. Results

### 3.1. Social behavioral results

The data obtained in the social preference test are shown in Fig. 1. The statistical analysis showed that wild-type animals spent more time in the 'social compartment' in comparison to the amount of time spent in the 'inanimate compartment'. However, the animals lacking the *LPA<sub>1</sub>* receptor spent significantly less time in the 'social compartment' ( $t_{12} = -2.75$ ,  $p = 0.017$ ,  $(1-\beta) = 0.84$ ,  $d = 0.15$ ) and more time in the 'central compartment' ( $t_{12} = 3.22$ ,  $p = 0.007$ ,  $(1-\beta) = 0.87$ ,  $d = 0.17$ ; Fig. 1A and B).

In addition, *maLPA<sub>1</sub>*-null mice had significantly fewer entries in the 'social compartment' than those of the wild-type animals ( $t_{12} = -5.99$ ,



**Fig. 2.** HPA-axis response to dexamethasone is altered in mice lacking the *LPA<sub>1</sub>* receptor. CORT release from 9:00–15:00 in both genotypes indicated an inverted pattern in the *maLPA<sub>1</sub>*-null mice in comparison with that in the wild-type mice (A). After controlling for the baseline values of CORT, the data indicated that *maLPA<sub>1</sub>*-null mice exhibited nonsuppression of CORT levels at 6 h after dexamethasone administration (B). Data are represented as the mean  $\pm$  standard error of the mean (SEM). \*  $p \leq 0.05$  and \*\*  $p \leq 0.01$  compared to the wild-type mice-Veh; #  $p < 0.05$  compared to the wild-type mice-DEX. N: wild-type mice-Veh = 5; wild-type mice-DEX = 5; *maLPA<sub>1</sub>*-null mice-Veh = 5; *maLPA<sub>1</sub>*-null mice-DEX = 5).

$p < 0.001$ ,  $(1-\beta) = 1$ ,  $d = 0.32$ ; Fig. 1C).

Regarding the ethological behaviors, mice lacking the LPA<sub>1</sub> receptor remained in the freezing state in the 'central compartment' for a longer time ( $t_{12} = -2.41$ ,  $p = 0.032$ ,  $(1-\beta) = 0.60$ ,  $d = 0.13$ ; Fig. 1E), while they spent significantly less time rearing in the 'social compartment' ( $t_{12} = -2.39$ ,  $p = 0.034$ ,  $(1-\beta) = 0.61$ ,  $d = 0.13$ ) and relatively less time (in relation to the total time in the 'social compartment') rearing ( $t_{12} = -2.43$ ,  $p = 0.032$ ,  $(1-\beta) = 0.69$ ,  $d = 0.13$ ) and sniffing ( $t_{12} = -4.33$ ,  $p < 0.001$ ,  $(1-\beta) = 0.97$ ,  $d = 0.23$ ; Fig. 1F and G). Importantly, we did not observe any differences between groups in either the velocity or the distance measurements (Fig. 1H and I).

PCA extraction resulted in Factor 1 consisting of time (0.85), rearing (0.90) and sniffing (0.94) in the 'social chamber' and Factor 2 consisting of freezing (0.86) and grooming (0.76) in the 'social chamber' ( $KMO = 0.51$ ;  $\chi^2 = 34.69$ ;  $p < 0.001$ ; Fig. 1J). Additionally, mice lacking the LPA<sub>1</sub> receptor had significantly lower PC scores in Factor 1 compared to those of the wild-type mice ( $t_{12} = -3.47$ ,  $p = 0.005$ ,  $(1-\beta) = 0.95$ ,  $d = 0.19$ ; Fig. 1K).

### 3.2. CORT response to the dexamethasone suppression test

The normal pattern of daytime release of CORT was inverted in the *maLPA<sub>1</sub>*-null mice, which displayed higher levels at 9:00 h and lower levels at 15:00 h in comparison with the levels of the wild-type mice (Fig. 2A). Related to DST, the ANCOVA analysis revealed a significant interaction in the analysis of covariance between genotype and treatment ( $F[1,15] = 5.39$ ;  $p < 0.05$ ; Fig. 2B), showing nonsuppression by *maLPA<sub>1</sub>*-null mice in the DST.

## 4. Discussion

Social behavior is fundamental for emotional regulation [15]. Conversely, social isolation can have negative consequences on an individual's physical, cognitive and mental health [16]. In fact, loss of social contact has a significant impact on cardiovascular health and life expectancy, cognitive impairment and mood [17–20]. Dysfunctional social behaviors are particularly prominent in patients with comorbid anxiety and depressive disorders than either disease alone (Saris et al., 2017). Understanding the neurobiological mechanisms responsible for social isolation, which in turn are related to the development of several psychological diseases, may have important implications for the treatment or prevention. For this reason, we have further characterized the involvement of the LPA<sub>1</sub> receptor in social behavior. This receptor has been linked to problems in emotional regulation [9,21] and the development of a mixed depressive-anxiety phenotype [7,8].

On the other hand, social behavioral disturbances have been associated with dysregulation of the HPA-axis [4,22]. In fact, there is an association between relatively high HPA-axis activity and social avoidance, indicating the importance of an intact and functional HPA-axis for exhibiting successful social behaviors [4]. Different data suggest the participation of the LPA<sub>1</sub> receptor in the regulation of the HPA axis. Thus, on the one hand, exogenous administration of LPA induces a reduction of CORT levels at baseline but potentiates CORT release following chronic stressor application [23]. On the other hand, animals lacking the LPA<sub>1</sub> receptor show signs of dysregulation of the HPA axis [9,10]. However, this axis has not been specifically evaluated in *maLPA<sub>1</sub>*-null animals. The dexamethasone suppression test is commonly used to evaluate the dysregulation of the HPA axis particularly in depressive patients [24,25] and may be a useful tool to study the viability of the HPA axis in animal models [26,27].

Considering these data and with the aim of better characterizing the null animal as an animal model of depression, we have studied on the one hand the social behavior and on the other hand the response of the HPA axis to the application of an exogenous glucocorticoid in both wild type and *maLPA<sub>1</sub>*-null mice.

The present study has allowed us to know the involvement of the

LPA<sub>1</sub> receptor in social behavior. Thus, while wild-type mice showed a tendency to socialize, the lack of the LPA<sub>1</sub> receptor was associated with increased social avoidance tendencies during social exposure. Since no differences in either the velocity or distance travelled were observed between the groups, social avoidance may not be attributed to abnormalities in locomotor activity. Hence, the substantial decrease in vertical activity, a validated indicator in rodent depression [28] and anxiety [29] models, along with a reduced time spent sniffing the juvenile animal, was associated with a lower score in the sociability factor in PCA analyses. This result seems to suggest that the absence of the LPA<sub>1</sub> receptor led to impaired social behaviors [30].

Regarding the endocrinological factors, DST revealed a blunted response in null mice to an exogenous corticosteroid administration. Moreover, the circadian rhythm of corticosterone, a domain that is consistently dysregulated in many mental illnesses including anxious depression [31], seems to be disturbed in *maLPA<sub>1</sub>*-null mice. The dysregulation of the HPA axis evidenced by DST could be related to alterations in social behavior in *maLPA<sub>1</sub>*-null animals. However, further studies are needed to verify this association and to provide enough detail regarding CORT levels over time to allow adequate assessment of circadian changes in CORT levels in the absence of the LPA<sub>1</sub> receptor.

Over the past decade, it has been shown that the LPA<sub>1</sub> receptor can be targeted by antidepressants, such as tricyclic, tetracyclic, desipramine and fluoxetine [7], causing ERK1/2 stimulation and cellular responses [32]. The LPA<sub>1</sub> receptor could also be involved in catecholamine secretion [33]. For instance, gintonin, a ligand for LPA receptors, has anti-inflammatory effects against neurodegenerative conditions like Alzheimer's, Parkinson's, anxiety, and depression [34]. Research suggests that gintonin's release of catecholamine, acetylcholine, and glutamate in the brain may alleviate depression-like effects, providing insights for potential use in managing neurodegenerative conditions. Regarding the anti-inflammatory effects, recently Nagata et al. (2023) [35] reported that LPA treatment prevents microglial activation, suppresses IL-1 $\beta$  and TNF- $\alpha$  expression in the hippocampus, improving depression-like behaviors. Interestingly, another recent study found that amitriptyline, a typical tricyclic, mediates antidepressant effects such as normalizing the CORT-induced decrease in sucrose preference by binding to LPA<sub>1</sub> receptor [36]. Thus, the use of LPA<sub>1</sub> receptor agonists could potentially be a new family of antidepressants with anxiolytic profiles, having a buffering effect on HPA axis and stress hormones release. However, to date no drugs targeting LPA receptors have been approved by any regulatory agency.

Overall, the results presented here have provided, for the first time, data for the involvement of the LPA<sub>1</sub> receptor in social behavior, implicating the absence of the LPA<sub>1</sub> receptor in social avoidance. Moreover, altered HPA-axis functioning and possibly circadian rhythms, factors involved in the etiology of depression and especially the anxious depression subtype, have been verified in *maLPA<sub>1</sub>*-null mice. However, we also want to mention certain limitations of our study. On the one hand, the sample size is small, so it has been necessary to statistically control that the observed differences are not due to Type 1 errors. Thus, the resulting effect sizes could be considered as small and medium. On the other hand, it would have been very interesting to include females in our study, because it would have enhanced the extrapolation and interpretation of our results. In conclusion, these results extend the previous evidence on the participation of the LPA<sub>1</sub>-receptor in the development of depression-like features and expanding the opportunity for developing drugs to target the LPA<sub>1</sub> receptor that may be useful for the treatment of mood disorder.

### CRediT authorship contribution statement

Conceptualization: C.P., F.RdF., G.E.-T., L.J.S. and R.D.M.-F.; Data curation: A.N.-Q., F.G.-S., P.S.-P. and R.D.M.-F.; Formal analysis: C.P., P.S.-P. and R.D.M.-F.; Funding acquisition: C.P. and G.E.-T.; Investigation: A.N.-Q., C.P., F.G.-S. and R.D.M.-F.; Methodology: P.S.-P. and



R.D.M.-F.; Supervision: C.P., F.RdF. and L.J.S.; Writing – original draft: C.P., F.G.-S. and R.D.M.-F.; Writing – review & editing: C.P., F.G.-S., P. S.-P. and R.D.M.-F.

## Declaration of Competing Interest

Authors declare no conflicts of interest. All authors have agreed with the submission in its present form

## Data availability

Data will be made available on request.

## Acknowledgements

We are grateful to Renata Ottilia Marvanykövi for her collaboration in the assessment of the behavioral tasks. We thank the University of Malaga for kindly providing the mice used in this study. This research was funded by the Andalusian Ministry of Knowledge, Research and University (P20\_00460 to CP); the Spanish Ministry of Economy and Competitiveness (PID2020-117464RB-I00 to CP) and Universidad Francisco de Vitoria (UFV2022-41 to RDM-F).

## References

- H.J. Möller, B. Bandelow, H.-P. Volz, U.B. Barnikol, E. Seifritz, S. Kasper, The relevance of “mixed anxiety and depression” as a diagnostic category in clinical practice, *Eur. Arch. Psychiatry Clin. Neurosci.* 266 (2016) 725–736, <https://doi.org/10.1007/s00406-016-0684-7>.
- E. Nagy, S. Moore, Social interventions: an effective approach to reduce adult depression? *J. Affect. Disord.* 218 (2017) 131–152, <https://doi.org/10.1016/j.jad.2017.04.043>.
- I.M.J. Saris, M. Aghajani, S.J.A. van der Werff, N.J.A. van der Wee, B.W.J. H. Penninx, Social functioning in patients with depressive and anxiety disorders, *Acta Psychiatr. Scand.* 136 (2017) 352–361, <https://doi.org/10.1111/acps.12774>.
- K. Roelofs, J. van Peer, E. Berretty, P. de Jong, P. Spinhoven, B.M. Elzinga, Hypothalamus–pituitary–adrenal axis hyperresponsiveness is associated with increased social avoidance behavior in social phobia, *Biol. Psychiatry* 65 (2009) 336–343, <https://doi.org/10.1016/j.biopsych.2008.08.022v>.
- B.J. Carroll, Use of the dexamethasone suppression test in depression, *J. Clin. Psychiatry* 43 (1982) 44–50.
- E.A. Young, J.L. Abelson, O.G. Cameron, Effect of comorbid anxiety disorders on the Hypothalamic–Pituitary–Adrenal axis response to a social stressor in major depression, *Biol. Psychiatry* 56 (2004) 113–120, <https://doi.org/10.1016/j.biopsych.2004.03.017>.
- R.D. Moreno-Fernández, M. Pérez-Martín, E. Castilla-Ortega, C. Rosell-Valle, M. I. García-Fernández, J. Chun, G. Estivill-Torrús, F. Rodríguez de Fonseca, L. J. Santín, C. Pedraza, malPA1-null mice as an endophenotype of anxious depression, *Transl. Psychiatry* 7 (2017), e1077, <https://doi.org/10.1038/tp.2017.24>.
- S. Tabbaï, R.D. Moreno-Fernández, E. Zambrana-Infantes, A. Nieto-Quero, J. Chun, M. García-Fernández, G. Estivill-Torrús, F. Rodríguez de Fonseca, L.J. Santín, T. G. Oliveira, M. Pérez-Martín, C. Pedraza, Effects of the LPA1 receptor deficiency and stress on the hippocampal LPA species in mice, *Front. Mol. Neurosci.* 12 (2019), 146, <https://doi.org/10.3389/fnmol.2019.00146>.
- C. Pedraza, J. Sánchez-López, E. Castilla-Ortega, C. Rosell-Valle, E. Zambrana-Infantes, M. García-Fernández, F. Rodríguez de Fonseca, L.J. Santín, G. Estivill-Torrús, Fear extinction and acute stress reactivity reveal a role of LPA(1) receptor in regulating emotional-like behaviors, *Brain Struct. Funct.* 219 (2014) 1659–1672, <https://doi.org/10.1007/s00429-013-0592-9>.
- E. Castilla-Ortega, C. Hoyo-Becerra, C. Pedraza, J. Chun, F. Rodríguez de Fonseca, G. Estivill-Torrús, L.J. Santín, Aggravation of chronic stress effects on hippocampal neurogenesis and spatial memory in LPA<sub>1</sub> receptor knockout mice, *PLoS One* 6 (2011), e25522, <https://doi.org/10.1371/journal.pone.0025522>.
- A.A. Boucher, J.C. Arnold, G.E. Hunt, A. Spiro, J. Spencer, I.S. McGregor, M.R. Bennett, M. Kassiou, Resilience and reduced c-Fos expression in P2×7 receptor knockout mice exposed to repeated forced swim test, *Neuroscience* 189 (2011) 170–177, <https://doi.org/10.1016/j.neuroscience.2011.05.049>.
- G. Estivill-Torrús, P. Liebrez-Zayas, E. Matas-Rico, L.J. Santín, C. Pedraza, I. De Diego, I. Del Arco, P. Fernández-Llebrez, J. Chun, F. Rodríguez de Fonseca, Absence of LPA1 signaling results in defective cortical development, *Cereb. Cortex* 18 (2008) 938–950, <https://doi.org/10.1093/cercor/bhm132>.
- J.J.A. Contos, N. Fukushima, J.A. Weiner, D. Kaushal, J. Chun, Requirement for the lpA1 lysophosphatidic acid receptor gene in normal suckling behavior, *Proc. Natl. Acad. Sci. USA* 97 (2000) 13384–13389, <https://doi.org/10.1073/pnas.97.24.13384>.
- E. Castilla-Ortega, C. Rosell-Valle, E. Blanco, C. Pedraza, J. Chun, F. Rodríguez de Fonseca, G. Estivill-Torrús, L.J. Santín, Reduced wheel running and blunted effects of voluntary exercise in LPA1-null mice: the importance of assessing the amount of running in transgenic mice studies, *Neurosci. Res.* 77 (2013) 170–179, <https://doi.org/10.1016/j.neures.2013.09.004>.
- T. English, L. Eldesouky, We're not alone: understanding the social consequences of intrinsic emotion regulation, *Emotion* 20 (2020) 43–47, <https://doi.org/10.1037/emo0000661>.
- N. Pai, S.L. Vella, The physical and mental health consequences of social isolation and loneliness in the context of COVID-19, *Curr. Opin. Psychiatry* 35 (2022) 305–310, <https://doi.org/10.1097/YCO.0000000000000806>.
- A. Hajek, H.H. König, Do lonely and socially isolated individuals think they die earlier? The link between loneliness, social isolation and expectations of longevity based on a nationally representative sample, *Psychogeriatrics* 21 (2021) 571–576, <https://doi.org/10.1111/psyg.12707>.
- H. Li, N. Xia, The role of oxidative stress in cardiovascular disease caused by social isolation and loneliness, *Redox Biol.* 37 (2020), 101585, <https://doi.org/10.1016/j.redox.2020.101585>.
- A. Orben, L. Tomova, S.J. Blakemore, The effects of social deprivation on adolescent development and mental health, *Lancet Child. Adolesc. Health* 4 (2020) 634–640, [https://doi.org/10.1016/S2352-4642\(20\)30186-3](https://doi.org/10.1016/S2352-4642(20)30186-3).
- M. Ahmed, I. Cerda, M. Maloof, Breaking the vicious cycle: The interplay between loneliness, metabolic illness, and mental health, *Front Psychiatry* 14 (2023), 1134865, <https://doi.org/10.3389/fpsyt.2023.1134865>.
- L.J. Santín, A. Bilbao, C. Pedraza, E. Matas-Rico, D. López-Barroso, E. Castilla-Ortega, J. Sánchez-López, R. Riquelme, I. Varela-Nieto, P. de la Villa, M. Suardíaz, J. Chun, F. Rodríguez de Fonseca, G. Estivill-Torrús, Behavioral phenotype of malPA1-null mice: increased anxiety-like behavior and spatial memory deficits, *Genes Brain Behav.* 8 (2009) 772–784, <https://doi.org/10.1111/j.1601-183X.2009.00524.x>.
- B.A. Gasser, J. Kurz, W. Senn, G. Escher, M.G. Mohaupt, Stress-induced alterations of social behavior are reversible by antagonism of steroid hormones in C57/BL6 mice, *Naunyn. Schmiede Arch. Pharmacol.* 394 (2021) 127–135, <https://doi.org/10.1007/s00210-020-01970-7>.
- R.D. Moreno-Fernández, C. Rosell-Valle, A. Bacq, O. Zanoletti, M. Cifuentes, M. Pérez-Martín, A.L. Gavito, M.I. García-Fernández, G. Estivill-Torrús, F. Rodríguez de Fonseca, L.J. Santín, C. Sandi, C. Pedraza, LPA1 receptor and chronic stress: effects on behaviour and the genes involved in the hippocampal excitatory/inhibitory balance, *Neuropharmacology* 164 (2020), 107896, <https://doi.org/10.1016/j.neuropharm.2019.107896>.
- A. Karssen, O. Meijer, E. De Kloet, Corticosteroids and the blood–brain barrier, *Tech. Behav. Neural Sci.* (2005) 329–340.
- L.L. Koorneef, M. van der Meulen, M.S. Kooijman, E. Sánchez-López, J. F. Scheerstra, M.C. Voorhoeve, A.N.N. Ramesh, P.C.N. Rensen, M. Giera, J. Kroon, O.C. Meijer, Dexamethasone-associated metabolic effects in male mice are partially caused by depletion of endogenous corticosterone, *Front. Endocrinol.* 13 (2022), 960279, <https://doi.org/10.3389/fendo.2022.960279>.
- S. Lurie, C. Kuhn, J. Bartolome, S. Schanberg, Differential sensitivity to dexamethasone suppression in an animal model of the DST, *Biol. Psychiatry* 26 (1989) 26–34, [https://doi.org/10.1016/0006-3223\(89\)90005-x](https://doi.org/10.1016/0006-3223(89)90005-x).
- S. Ridder, S. Chourbaji, R. Hellweg, A. Urani, C. Zacher, W. Schmid, M. Zink, H. Hörtnagl, H. Flor, F.A. Henn, G. Schütz, P. Gass, Mice with genetically altered glucocorticoid receptor expression show altered sensitivity for stress-induced depressive reactions, *J. Neurosci.* 25 (2005) 6243–6250, <https://doi.org/10.1523/JNEUROSCI.0736-05.2005>.
- U. Skupio, M. Tertli, M. Sikora, S. Golda, A. Wawrzczak-Bargiela, R. Przewlocki, Behavioral and molecular alterations in mice resulting from chronic treatment with dexamethasone: relevance to depression, *Neuroscience* 286 (2015) 141–150, <https://doi.org/10.1016/j.neuroscience.2014.11.035>.
- G.-X. Liu, G.-Q. Cai, Y.-Q. Cai, Z.J. Sheng, J. Jiang, Z. Mei, Z.G. Wang, L. Guo, J. Fei, Reduced anxiety and depression-like behaviors in mice lacking GABA transporter subtype 1, *Neuropsychopharmacology* 32 (2007) 1531–1539, <https://doi.org/10.1038/sj.npp.1301281>.
- O. Kaidanovich-Beilin, T. Lipina, I. Vukobradovic, J. Roder, J.R. Woodgett, Assessment of social interaction behaviors, *J. Vis. Exp.* (48) (2011) 2473, <https://doi.org/10.3791/2473>.
- S. Rao, S. Zisook, Anxious depression: clinical features and treatment, *Curr. Psychiatry Rep.* 11 (2009) 429–436, <https://doi.org/10.1007/s11920-009-0065-2>.
- M.C. Olanas, S. Dedoni, P. Onali, Antidepressants activate the lysophosphatidic acid receptor LPA(1) to induce insulin-like growth factor-I receptor transactivation, stimulation of ERK1/2 signaling and cell proliferation in CHO-K1 fibroblasts, *Biochem. Pharmacol.* 95 (2015) 311–323, <https://doi.org/10.1016/j.bcp.2015.04.002>.
- T. Maruta, T. Yanagita, K. Matsuo, Y. Uezono, S. Satoh, T. Nemoto, N. Yoshikawa, H. Kobayashi, M. Takasaki, A. Wada, Lysophosphatidic acid-LPA1 receptor-Rho kinase-induced up-regulation of Nav1.7 sodium channel mRNA and protein in adrenal chromaffin cells: enhancement of 22Na<sup>+</sup> influx, 45Ca<sup>2+</sup> influx and catecholamine secretion, *J. Neurochem.* 105 (2008) 401–412, <https://doi.org/10.1111/j.1471-4159.2007.05143.x>.

- [34] M. Ikram, R. Ullah, A. Khan, M.O. Kim, Ongoing research on the role of gintonin in the management of neurodegenerative disorders, *Cells* 9 (2020) 1464, <https://doi.org/10.3390/cells9061464>.
- [35] W. Nagata, A. Koizumi, K. Nakagawa, S. Takahashi, M. Gotoh, Y. Satoh, T. Ishizuka, Treatment with lysophosphatidic acid prevents microglial activation and depression-like behaviours in a murine model of neuropsychiatric systemic lupus erythematosus, *Clin. Exp. Immunol.* 212 (2023) 81–92, <https://doi.org/10.1093/cei/uxad010>.
- [36] N. Kajitani, M. Okada-Tsuchioka, A. Inoue, K. Miyano, T. Masuda, S. Boku, K. Iwamoto, S. Ohtsuki, Y. Uezono, J. Aoki, M. Takebayashi, G protein-biased LPAR1 agonism promotes prototypic antidepressant effects, *BioRxiv* (2022), <https://doi.org/10.1101/2022.11.02.514841>.