

COMMUNICATION

View Article Online
View Journal | View IssueCite this: *Dalton Trans.*, 2024, **53**, 8550Received 18th December 2023,
Accepted 16th April 2024

DOI: 10.1039/d3dt04247b

rsc.li/dalton

Ni(2,2':6',2''-terpyridine)₂: a high-spin octahedral formal Ni(0) complex†Natalia Cabrera-Lobera,^a Estefanía del Horno,^a M. Teresa Quirós,^b Elena Buñuel,^a Magali Gimeno,^{c,d} William W. Brennessel,^{b,c} Michael L. Neidig,^{b,d} José Luis Priego^e and Diego J. Cárdenas^{b,a*}

We have synthesised and characterised the complex Ni(tpy)₂ (tpy = 2,2':6',2''-terpyridine). This formally Ni(0) complex is paramagnetic both in the solid state and in solution (*S* = 2). The crystal structure shows an octahedral geometry, with molecules arranged in independent dimers involving π -stacking between pairs of complexes. Magnetic measurements and DFT calculations suggest the existence of temperature-dependent intermolecular antiferromagnetic coupling in the solid state.

The importance of Ni complexes as catalysts for synthetically useful reactions has experienced a renaissance in recent years.¹ The accessibility of a wide variety of oxidation states has allowed the discovery of novel activation pathways of simple substrates and the development of multicomponent reactions for the ready preparation of complex organic substrates.² Especially relevant are Ni(I) derivatives, which have been proposed as intermediates in C–C bond formation reactions such as cross-couplings of alkyl and aryl halides with organomagnesium and organozinc reagents,³ and haloalkane carboxylation reactions.⁴ The occurrence of several oxidation states along the catalytic cycles allows complex cascade processes leading to the formation of several C–C and C–B bonds, such as cyclization-couplings,⁵ difunctionalization of alkenes with alkyl and aryl halides,⁶ and diborylative cyclization of enynes.⁷ Part of the success of these reactions relies on the use

of redox-active ligands capable of stabilizing different oxidation states.⁸ A general feature of such ligands in Ni-based catalytic systems is their capability to tune their reactivity by delocalizing electrons from the metal. Thus, the spin density in an open shell complex will not be localized on the metal centre but it will be partially delocalized on the ligands. This effect may confer special stability and modulate the reactivity of radical metal complexes. 2,2':6',2''-Terpyridine (tpy) is an important ligand in synthetically useful Ni-catalyzed reactions,⁹ and it may stabilize low-valent Ni complexes. In fact, it participates in the first isolated Ni(I) organometallic derivative (CH₃)Ni(tpy), a 17-electron square-planar complex.¹⁰ Calculations show the unpaired electron lying mainly on the π^* orbitals of tpy, rendering an anionic ligand coordinated to a Ni(II) center. However, the actual behavior depends on the rest of the ligands on the coordination sphere. Thus, the analogous Br derivative shows a metal centered SOMO instead (Fig. 1).¹¹

Polydentate pyridine-based ligands have been extensively used in Ni-catalyzed synthetically useful reactions involving organometallic intermediates.¹² The accessibility of Ni(I) is especially relevant in this context, and the formation and the electronic structure of these kinds of complexes have been studied both experimentally and computationally.^{4c,10,13} Some years ago, preparation of a complex with Ni(tpy)₂ formula was published, and the electronic structure of different putative geometries for this molecular formula were studied by DFT methods.¹⁴ Nevertheless, no spectroscopic or crystal structure data were reported. On the basis of its diamagnetic character

^aDepartment of Organic Chemistry, Facultad de Ciencias, Universidad Autónoma de Madrid, Institute for Advanced Research in Chemical Sciences (IAdChem), Red ORFEO-CINQA, Av. Francisco Tomás y Valiente 7, Campus de Cantoblanco, 28049 Madrid, Spain. E-mail: diego.cardenas@uam.es

^bDepartment of Organic Chemistry and Inorganic Chemistry, Facultad de Farmacia, Universidad de Alcalá de Henares, Campus Universitario, 28871 Madrid, Spain

^cDepartment of Chemistry, University of Rochester, Rochester, New York 14627, USA

^dInorganic Chemistry Laboratory, Department of Chemistry, University of Oxford, South Parks Road, Oxford OX1 3QR, UK

^eDepartment of Inorganic Chemistry, Universidad Complutense de Madrid, Ciudad Universitaria, 28040 Madrid, Spain

† Electronic supplementary information (ESI) available. CCDC 1891622. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d3dt04247b>




Fig. 1 Terpyridine may act as a redox-active ligand in Ni(I) complexes.

in the solid state and calculations, it was proposed to have the tetrahedral arrangement that could be expected for a common Ni(0) complex. For an alternative octahedral arrangement, calculations suggested electron delocalization on both terpyridine ligands, as it could be expected from previous studies on formally Ni(I) organometallic complexes with this and related ligands.

We became interested in the actual structure of Ni(0)-tpy complexes, since they may presumably be involved as precatalysts in the Ni-catalyzed diborylative cyclization of enynes we have previously reported.⁷ In addition, determination of crystal structures may help to shed light on their electronic structure. Thus, we tried to isolate and characterize species formed when Ni(0) precatalysts react with tpy in the absence of other reagents. In contrast to previous report,¹⁴ the reaction of Ni(cod)₂ with tpy at room temperature in toluene led to a dark blue solution from which crystals of Ni(tpy)₂ (**1**) were obtained (Fig. 2).¹⁵ It is important to mention that these conditions differ from those previously reported for the preparation of the diamagnetic complex with the same empirical formula tetrahedral isomer (cyclohexane, 12 h, 80 °C). This emphasizes that solvent and reaction conditions are critical in isolating the octahedral compound.

After extensive experimentation, we found that slow vapor diffusion of pentane at −25 °C (24 h) over a solution of a mixture of Ni(cod)₂ and tpy afforded dark blue crystals suitable for X-ray diffraction. Crystal structure determination showed an unexpected octahedral Ni(tpy)₂ complex with the asymmetric unit containing two [Ni(tpy)₂]⁰ molecules in general positions (Fig. 3).

The average Ni–N bond lengths of 2.127(7) [outer] and 1.985(4) [inner] Å slightly differ from the ones found for Ni^{II}(tpy)₂Cl₂ (2.02–2.14) Å,¹⁶ and are similar those found for dicationic [Ni(ftpy)₂]²⁺ (ftpy = 4'-(furan-2-yl)-tpy) of 2.107(8) [outer] and 1.991(5) [inner] Å.¹⁷ More telling, however, are the average C–C bond lengths between pyridine rings, which is 1.456(12) Å for our structure and 1.493(12) Å for [Ni(ftpy)₂]²⁺,¹⁷ the latter which is consistent with that of free tpy at 100 K.¹⁸ This trend is seen in the Fe system in which the average C–C bond lengths between pyridine rings shorten in going from [Fe(tpy)₂]²⁺ to [Fe(tpy)₂]⁰, 1.467(4) to 1.452(4) Å, which Wieghardt attributes to reduction of the ligands.¹⁹ Moreover, Ni⁰(bpy)₂ shows shorter Ni–N distances, around 1.92 Å,¹⁴ and Ni^I(tpy)R (R = Br, Me) range between 1.92–2.04.^{10,11} Thus, our complex




Fig. 2 Preparation of Ni(tpy)₂ complexes.




Fig. 3 Anisotropic displacement ellipsoid plot at the 50% probability level with hydrogen atoms omitted.

is best described as [Ni²⁺(tpy[−])₂]⁰, a nickel(II) center with two radical anionic tpy ligands.

Inspection of the crystal packing shows that molecules of the complex interact by stacking of tpy ligands with some of their neighbours, whereas other close molecules show just edge-to-face interaction between the ligands. Thus, three kinds of dimeric units can be considered in this structure (Fig. 4).

The ¹H-NMR in spectrum of the isolated crystals dissolved in *d*⁸-THF shows the paramagnetic character of this compound, with signals expanding between 160 and −140 ppm (see ESI).²⁰ On the other hand, quantitative EPR shows a signal corresponding to a complex with *S* = 1/2. However, this species only represents 1% of the total sample which suggests that the major species has *S* = 1 or 2. Following the Evans method, which uses difference in the NMR chemical shift in a solvent caused by the presence of a paramagnetic species,¹⁷ we measured the effective magnetic moment (*μ*_{eff}) in the solution of complex **1**, obtaining a value of 4.9(2)*μ*_B in THF and 5.2(2)*μ*_B in toluene. These results are consistent with a *S* = 2 ground-state for **1** in solution.

This complex is also paramagnetic in the solid state, in contrast to previously reported species which is diamagnetic.¹⁴ Its




Fig. 4 Three different arrangements for neighbouring molecules of Ni(tpy)₂. I and II show stacking pyridine ligands and different edge-to-face interactions; III presents just edge-to-face interactions, without stacking rings.

magnetic moment decreases from $3.8\mu_B$ at 298 K to 1.4 at 2 K (Fig. 5). In contrast, the analogous $M^0(\text{tpy})_2$ complexes of Ti,²¹ Cr,²² Mo,²² and W,²² are diamagnetic ($S = 0$) despite they show very similar crystal structures. On the other hand, Fe and Ru derivatives show $S = 1$,¹⁹ showing same unit cell and packing as **1**, as well.

We studied the electronic structure of the octahedral complex by DFT calculations at M06-2X/6-31G(d) (C,H,N), Ni (LANL2DZ) level, functional for different spin multiplicities.²³

For isolated $\text{Ni}(\text{tpy})_2$ molecules of complex **1**, which are the expected to be in solution, the quintet ground state is far more stable compared with the singlet and triplet states (Table 1).²⁴

Moreover, the quintuplet structure shows an octahedral arrangement, whereas the calculated triplet has a distorted tetrahedral structure, with two non-coordinated N atoms (one of each ligand) and the singlet being highly distorted. The high stability of the quintet is in accord with the Evans measurement in solution, which suggested $S = 2$ for complex **1**. On the other hand, the singly-occupied molecular orbitals for the four unpaired electrons are all ligand-centered (Fig. 6) which further supports a Ni(II) metal center coordinated to two anionic tpy ligands, as suggested by the crystal structure.

Then we extended our study to pairs of molecules, and considered the three kinds of arrangements, which can be distinguished in the crystal structure (Fig. 4), as models.

We computed the energy of these dimers for different spin states by broken-symmetry (BS) DFT calculations,²⁵ which allow to assign different spin multiplicities to the Ni complexes in the dimer to determine the energy for complexes with possible antiferromagnetic coupling. We took the atomic coordinates from the crystal structure and calculated the energy for the three models. As it can be seen in Table 1 (center), the most stable arrangement corresponds to the nonaplet dimer **I** showing π -stacking interaction between pyridines of two different complexes. The alternative nonaplet species are slightly less stable (2.2 and 2.0 kcal mol⁻¹ for **II** and **III**, respectively). These results indicate that π -stacking does not stabilize nonaplet spin states significantly. On the other hand, singlet states for dimer **III**, which lacks stacking

Table 1 Calculated relative energies for $\text{Ni}(\text{tpy})_2$ and the dimeric sub-units (kcal mol⁻¹). $m = 2S + 1$ (spin multiplicity). M06-2X/6-31G(d) (C,H,N) LANL2DZ (Ni)

Ni(tpy) ₂ optimized structures			
	$S = 0, m = 1$ (singlet)	$S = 1, m = 3$ (triplet)	$S = 2, m = 5$ (quintet)
Relative E	0	-28.9	-44.2
Relative G (298 K)	0	-31.8	-46.8
Relative energy for different [Ni(tpy)₂]₂ Single-point calculation on crystal structure fragments.			
	$S = 4, m = 9$ (nonaplet)	$S = 0$ (two triplets)	$S = 0$ (two quintets)
I	0	28.7	4.0
II	2.2	43.1	4.2
III	2.0	30.5	39.4
[Ni(tpy) ₂] ₂ optimized structures			
	$S = 4, m = 9$ (nonaplet)	$S = 0$ (two triplets)	$S = 0$ (two quintets)
Relative E	0	5.7	+1.7 (-0.6) ^a
Relative G (298 K)	0	10.9	2.3

^a Using SCF = QC as option in Gaussian optimization. See ref. 24.




Fig. 5 Temperature dependence of the molar susceptibility χ_M (red circles) and μ_{eff} (blue triangles) for solid $\text{Ni}(\text{tpy})_2$ at 0.5 T.

interactions, show much higher energies, regardless they are formed by either two triplet or quintet complexes (30.5 and 39.4 kcal mol⁻¹, respectively). On the other hand, regarding the dimers with π -stacked pyridines (**I** and **II**), the singlet (two-triplet) states are also considerably unstable compared to the nonaplets (28.7 and 43.1 kcal mol⁻¹, respectively). The results are in accord with the high magnetic moment measured in the solid state at room temperature. In contrast and noticeably, the corresponding singlet dimers composed of two quintets for both **I** and **II** lie only 4.0 and 4.2 kcal mol⁻¹ above the nonaplet, in sharp contrast with the energy obtained for the edge-to-face dimer without stacking interactions (**III**) which is much more unstable ($\Delta E = 39.4$ kcal mol⁻¹). These results suggest that antiferromagnetic coupling is feasible for the stacked molecules, considering the error inherent to the computational approach. The relative energy values are significant in this case to derive these conclusions.




Fig. 6 Calculated singly-occupied molecular orbitals for $\text{Ni}(\text{tpy})_2$ in the quintuplet state in decreasing order of energy from SOMO1 to SOMO 4.

Then, we performed structure optimizations for the π -stacked dimer in the three different spin states. This leads to structures different from those found in the crystal but may provide further structural information. As shown in Table 1 (bottom), the most stable structure at room temperature corresponds to the nonaplet dimer (in terms of G). However, the energy of the antiferromagnetic singlet dimer in which both $\text{Ni}(\text{tpy})_2$ units are quintets might be more stable at low temperature, according to the calculated electronic energy.²⁴ In any case, the energy difference compared to the nonaplet is very small and within the error of the calculations. Population of the high-spin state accounts for the overall magnetic moment observed at room temperature. Inspection of the molecular orbitals of the dimeric units for the singlet dimers containing two quintet units (HOMO to HOMO-5), show that external electrons are located on the ligands π^* orbitals and not on the metal centers (Fig. 7).

Comparison of the distances between centroids of pyridine rings in the crystal structure between **1** and the Fe and Ru analogues suggests that this kind of interaction could also operate in these derivatives (see ESI†).

In conclusion, coordination of two terpyridine ligands to Ni(0) induces the electron transfer to the π^* orbitals of the ligands, affording an octahedral quintet complex which should be considered as a $\text{Ni}(\text{II})$ species with two anionic tridentate ligands. In the solid state, close molecules may interact through ligand stacking and may couple antiferromagnetically, being the global magnetic moment dependent on the temperature. The high stability of the singlet dimers composed of quintets compared to other spin states, along with the difference of energy calculated on models for the different arrangements of molecules found in the crystal structure supports this proposal. This could be also the case for complexes of other transition metals. We can envision that incorporation of functional groups in the terpyridine ligands, or the use of




Fig. 7 Calculated two-electron occupied molecular orbitals for singlet dimeric units containing two quintuplet complexes **1**. HOMO-2 and HOMO-5 are responsible for the interaction between subunits.

other polydentate ligands may lead to new structures with extended electron delocalization. that may lead to new interesting solid-state structures or novel Ni precatalysts.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank the support by MICIU (PID2019-109088GB-I00 to D. J. C. and a Juan de la Cierva fellowship to M. T. Q) and INVESTIGO-CAM program (for a fellowship to E. d. H). We acknowledge the National Institutes of Health (R01GM111480 to M. L. N.). We also thank the Centro de Computaci3n Cientifica-UAM for computation time.

References

- (a) X. Qi and T. Diao, *ACS Catal.*, 2020, **10**, 8542; (b) K. E. Poremba, S. E. Dibrell and S. E. Reisman, *ACS Catal.*, 2020, **10**, 8237; (c) S. Z. Tasker, E. A. Standley and T. F. Jamison, *Nature*, 2014, **509**, 299.
- (a) C.-Y. Lin and P. P. Power, *Chem. Soc. Rev.*, 2017, **46**, 5347; (b) X. Hu, *Chem. Sci.*, 2011, **2**, 1867; (c) V. B. Phapale and D. J. C3rdenas, *Chem. Soc. Rev.*, 2009, **38**, 1598–1607.
- (a) G. M. Schwarzwald, C. D. Matier and G. C. Fu, *Angew. Chem., Int. Ed.*, 2019, **58**, 3571; (b) Z.-C. Cao, S.-J. Xie, H. Fang and Z.-J. Shi, *J. Am. Chem. Soc.*, 2018, **140**, 13575;

- (c) R. Soler-Yanes, I. Arribas-Álvarez, M. Guisán-Ceinos, E. Buñuel and D. J. Cárdenas, *Chem. – Eur. J.*, 2017, **23**, 1584; (d) R. Soler-Yanes, M. Guisán-Ceinos, E. Buñuel and D. J. Cárdenas, *Eur. J. Org. Chem.*, 2014, 6625; (e) P. M. P. Garcia, T. Di Franco, A. Epenoy, R. Scopelliti and X. Hu, *ACS Catal.*, 2016, **6**, 258; (f) O. Vechorkin, A. Godinat, R. Scopelliti and X. Hu, *Angew. Chem., Int. Ed.*, 2011, **50**, 11777; (g) O. Vechorkin and X. Hu, *Angew. Chem., Int. Ed.*, 2009, **48**, 2937; (h) A. J. Oelke, J. Sun and G. C. Fu, *J. Am. Chem. Soc.*, 2012, **134**, 2966; (i) S. W. Smith and G. C. Fu, *Angew. Chem. Int. Ed.*, 2008, **47**, 9334; (j) S. W. Smith and G. C. Fu, *J. Am. Chem. Soc.*, 2008, **130**, 12645; (k) V. B. Phapale, M. Guisán-Ceinos, E. Buñuel and D. J. Cárdenas, *Chem. – Eur. J.*, 2009, **15**, 12681.
- 4 (a) F. Juliá-Hernández, T. Moragas, J. Cornella and R. Martin, *Nature*, 2017, **545**, 84; (b) T. Yanagi, R. J. Somerville, K. Nogi, R. Martin and H. Yorimitsu, *ACS Catal.*, 2020, **10**, 2117; (c) R. Somerville, C. Odena, M. Obst, N. Hazari, K. Hopmann and R. Martin, *J. Am. Chem. Soc.*, 2020, **142**, 10936.
- 5 (a) M. Guisán-Ceinos, R. Soler-Yanes, D. Collado-Sanz, V. B. Phapale, E. Buñuel and D. J. Cárdenas, *Chem. – Eur. J.*, 2013, **19**, 8405; (b) V. B. Phapale, E. Buñuel, M. García-Iglesias and D. J. Cárdenas, *Angew. Chem., Int. Ed.*, 2007, **46**, 8790; (c) J. C. Nieto-Carmona, R. San Román, E. Buñuel and D. J. Cárdenas, *Eur. J. Org. Chem.*, 2022, e202200992.
- 6 (a) X. Wei, W. Shu, A. García-Domínguez, E. Merino and C. Nevado, *J. Am. Chem. Soc.*, 2020, **142**, 13515; (b) Y. Jin, H. Yang and C. Wang, *Org. Lett.*, 2020, **22**, 2724; (c) W. Shu, A. García-Domínguez, M. T. Quirós, R. Mondal, D. J. Cárdenas and C. Nevado, *J. Am. Chem. Soc.*, 2019, **141**, 13812.
- 7 N. Cabrera-Lobera, M. T. Quirós, W. W. Brennessel, M. L. Neidig, E. Buñuel and D. J. Cárdenas, *Org. Lett.*, 2019, **21**, 6552.
- 8 Redox-active ligands in catalysis: O. R. Luca and R. H. Crabtree, *Chem. Soc. Rev.*, 2013, **42**, 1440.
- 9 Y. H. Budnikova, D. A. Vicic and A. Klein, *Inorganics*, 2018, **6**, 18.
- 10 T. J. Anderson, G. C. Jones and D. A. Vicic, *J. Am. Chem. Soc.*, 2004, **126**, 8100–8101. see also Erratum: T. J. Anderson, G. D. Jones and D. A. Vicic, *J. Am. Chem. Soc.*, 2004, **126**, 11113.
- 11 J. T. Ciszewski, D. Y. Mikhaylov, K. V. Holin, M. K. Kadirov and Y. H. Budnikova, *Inorg. Chem.*, 2011, **50**, 8630–8635.
- 12 C. Wei, Y. He, X. Shi and Z. Song, *Coord. Chem. Rev.*, 2019, **385**, 1.
- 13 (a) C. S. Day, A. Rentería-Gómez, S. T. Jon, A. R. Gogoi, O. Gutiérrez and R. Martin, *Nat. Catal.*, 2023, **6**, 244; (b) M. T. Quirós, D. Collado-Sanz, E. Buñuel-Magdalena and D. J. Cárdenas, *Chem. Commun.*, 2021, **57**, 2424;
- (c) M. T. Quirós, D. Collado-Sanz, E. Buñuel-Magdalena and D. J. Cárdenas, *J. Phys. Chem. A*, 2018, **122**, 2250.
- 14 M. Wang, J. England, T. Weyhermüller and K. Wieghardt, *Eur. J. Inorg. Chem.*, 2015, 1511–1523.
- 15 Experimental procedure for the synthesis of complex **1**: In a glove-box, a mixture of Ni(cod)₂ (74 mg, 0.27 mmol) and terpyridine (125 mg, 0.57 mmol) were dissolved in toluene (3 mL) at room temperature. The resulting dark blue solution is stirred for 10 min, and pentane (5 mL) is added. The resulting precipitate is filtered to give a dark blue solid which is washed with pentane (76 mg, 54%).
- 16 (a) M. I. Arriortua and T. Rojo, *Bull. Soc. Chim. Belg.*, 1982, **91**, 337–338; (b) E. C. Constable, J. Lewis, M. C. Liptrot and P. R. Raithby, *Inorg. Chim. Acta*, 1990, **178**, 47.
- 17 B. Z. Momeni, S. K. Anari, M. Torrei and J. Janczak, *Appl. Organomet. Chem.*, 2021, **35**, e6179.
- 18 K. F. Bowes, I. P. Clark, J. M. Cole, M. Gourlay, A. M. E. Griffin, M. F. Mahon, L. Ooi, A. W. Parker, P. R. Raithby, H. A. Sparkes and M. Towrie, *CrystEngComm*, 2005, **7**, 269.
- 19 J. England, C. C. Scarborough, T. Weyhermüller, S. Sproules and K. Wieghardt, *Eur. J. Inorg. Chem.*, 2012, 4605.
- 20 (a) D. F. Evans, *J. Chem. Soc.*, 1959, 2003; (b) E. M. Schubert, *J. Chem. Educ.*, 1992, **69**, 62; (c) C. Piguet, *J. Chem. Educ.*, 1997, **74**, 815.
- 21 M. Wang, T. Weyhermüller, J. England and K. Wieghardt, *Inorg. Chem.*, 2013, **52**, 12763.
- 22 M. Wang, J. England, T. Weyhermüller, S.-L. Kokatam, C. J. Pollock, S. DeBeer, J. Shen, G. P. A. Yap, K. H. Theopold and K. Wieghardt, *Inorg. Chem.*, 2013, **52**, 4472.
- 23 Calculations with other density functionals based on M06 with different correlation-exchange ratios failed. Density functional wB97XD did not work either. On the other hand, as expected, B3LYP did not afford correct structures for the dimeric complexes.
- 24 Calculated energies (*E* in Table 1) correspond to potential energy minima. Energy values for the singlet dimer composed of two quintet subunits depend on the method used for self-consistent field optimization (QC or XQC). Free energy (*G*) values are calculated at 298 K. Calculated *G* values for dimers other than nonaplet could only be computed with SCF = XQC option of Gaussian 09 package.
- 25 (a) L. Noodleman, *J. Chem. Phys.*, 1981, **74**, 5737; (b) L. Noodleman, J. G. Norman, J. H. Osborne, A. Aizman and D. Case, *J. Am. Chem. Soc.*, 1985, **107**, 3418; (c) L. Noodleman and E. R. Davidson, *Chem. Phys.*, 1986, **109**, 131; (d) L. Noodleman, D. A. Case and A. J. Aizman, *J. Am. Chem. Soc.*, 1988, **110**, 1001; (e) L. Noodleman, C. Y. Peng, D. A. Case and J. M. Monesca, *Coord. Chem. Rev.*, 1995, **144**, 199.